

Fertility suppression of some wildlife species in southern Africa – A review

HJ Bertschinger^{1,*}, P Caldwell²

¹Veterinary Population Management Laboratory, Faculty of Veterinary Science, University of Pretoria, South Africa;

²Old Chapel Veterinary Clinic, Villieria, Pretoria, South Africa

*** Author's address (for correspondence):**

Henk Bertschinger, Veterinary Population Management Laboratory, Faculty of Veterinary Science, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa. E-mail: henkbert@tiscali.co.za

Contents

Generally speaking, southern Africa's wildlife populations in small to medium-sized protected game reserves (10 000 to 65 000 ha), reproduce at rapid rates which often leads overpopulation of certain species. Most commonly these are large predators such as lions, African wild dogs and cheetahs, and elephants. Overpopulation of large predators leads to depletion of prey species, break-outs into neighbouring communities and increased risks for disease transmission. An overabundance of elephants leads to habitat destruction which is to the detriment of not only other herbivores but also certain bird species. By far the most acceptable and effective method of population control is contraception. Another problem, particularly in South Africa, is the large numbers of large predators that are held in zoos, wildlife sanctuaries or captive breeding facilities. Once again, there is a need for contraception to control the rate of reproduction. In this review we discuss the methods that have been most commonly used for some wildlife species in southern Africa. The methods include hormonal control and

Wildlife fertility suppression southern Africa – a review immunocontraception. We also address the problem of androgen-related aggressive behaviours in elephant bulls and giraffe males and present solutions that have been used to ameliorate such behaviours.

Short history of contraception

According to Jöchle (2008), for thousands of years the only form of contraception practiced on domestic animals was surgical castration. Male castration dates back to 7-6000 BC. The early development of orchidectomy is not surprising, given that the male gonads of domestic species are situated extra-abdominally. More surprisingly, ovariectomy is quoted in Aristotle's writings as early as 384-322 BC (Jöchle 2008). In Europe, it is interesting to note that from the 15th to the 19th century neutering of male and female domestic animals was performed by professionals with a special license, but not veterinarians. It was only in the 18th and 19th centuries that veterinarians slowly entered the castration business. The possible use of intra-uterine devices in animals may date back as far as 3000 years ago, when nomads placed pebbles into the uteri of camels to prevent conception during their long treks through the desert (Museum of Contraception, University of Montpellier, France).

It was only during the second half of the 20th century, however, that non-surgical methods for contraception of house pets were widely adopted (Jöchle 2008). The first oral contraceptive for dogs, medroxy-progesterone acetate, became available in 1963. The development of 'modern' contraceptive methods such as the GnRH super-agonists and immunological tools such as porcine zona pellucida (pZP) and GnRH vaccines did not begin until 20-30 years later.

Contraception to control the rate of reproduction is an interesting and potentially practical means of population control in wildlife; both captive and free-ranging

Wildlife fertility suppression southern Africa – a review
populations. Broadly speaking, the following methods can be used for contraception of
animals:

- Surgical (gonadectomy, vasectomy and salpingectomy)
- Hormonal (oral contraceptives, depot-injections or slow-release implants)
- Immunocontraception

The ideal wildlife contraceptive should fulfil most of the following requirements:
Effective; allow remote delivery; be reversible, although this depends on requirements
for individual species and conditions; have little or no effect on social behaviours or
organization of groups or herds; have no deleterious short or long-term health effects;
should not pass through the food chain; be safe to use during pregnancy; have
affordable production and application costs (Bertschinger et al. 2008b).

Table 1 lists contraceptive methods that could be used and the extent to which they
comply with the properties of an ideal wildlife contraceptive.

This paper will concentrate on hormonal (also referred to as chemical) and
immunological methods. In many wildlife species surgical methods are impractical,
irreversible and, in the case of gonadectomy, affect behaviour.

Endocrine control of reproduction

Central endocrine control of reproduction is organised via the hypothalamo-pituitary-
gonadal axis (Fig. 1). The hypothalamic liberin, GnRH, is central to this axis and is the
primary controller of reproduction in both female and male mammals. GnRH stimulates
the release of both FSH and LH. The gonadotrophic hormones control gametogenesis
and gonadal steroid hormone production. Gonadal steroids are responsible for
secondary sex characteristics, sexual behaviour and modulate GnRH, FSH and LH
release. During the oestrous cycle the differential release of these gonadotrophic
hormones is complex and depends on the pulsatile nature of GnRH release as well as

Table 1. Extent to which potential wildlife contraceptive agents conform to the ideal properties (modified from Bertschinger et al. 2008)

Property	Surgical methods		Steroids		GnRH agonists		Immunocontraception	
	Gonadect	Vasec/FallT	Oral	Implants	Injectable	Implants	pZP	GnRH
Contraceptive efficacy								
Females	100%	100%	100% ^a	100%	≤100%	≤100%	70-100%	70-100%
Males	100%	100% ^b	Poor	Poor	≤100% ^c	≤100% ^c	No	70-100%
Remote delivery	No	No	Only captive	No	Yes	No ^d	Yes	Yes
Reversible	No	No ^e	Yes	Yes	Yes	Yes	Yes	Yes ^f
Social behaviours or organization of groups or herds	Affects behaviour	Some in cats	Affects behaviour	Affects behaviour	No data	♀ anoestrus ♂ aggression↓	♀ continue to cycle	♀ anoestrus ♂ aggression↓
Deleterious short or long-term health effects	Obesity	None	Carnivores some serious	Carnivores some serious	None	None	Local swelling	Local swelling
Contraceptive passes through the food chain	No	No	Possible	Yes	No	No	No	No
Safe to use during pregnancy	NA	NA	No	No	Yes/no	Yes/no	Yes	Yes
Production and/or application costs	Expensive	Expensive	Expensive	Medium	Medium	Medium	Medium	Medium

^bMales can remain fertile for a number of weeks; ^aOnly if given daily; ^cDoes not work in male ungulates; ^dRemote delivery system being developed (Herbert and Vogelnest 2007);

^eRequires microsurgery – not feasible under field conditions and especially not in mega-herbivores; ^fLong-acting formulations (GonaCon™) or longer term use in males or prepubertal use in both sexes may be irreversible

Gonadect = gonadectomy; Vasec = vasectomy; FallT = tying of fallopian tubes; NA = not applicable

feedback mechanisms from the ovary. The down-stream target cells of FSH and LH are the developing follicles (theca interna and granulosa cells). In the male FSH specifically binds to Sertoli cell membrane receptors, and is thus intricately involved in the regulation of spermatogenesis. Leydig cells on the other hand are targeted by LH which stimulates testosterone synthesis and secretion. Androgens also affect sperm production but more specifically the reduction division (spermatidogenesis) and spermiogenesis. The function of secondary reproductive organs such as the prostate gland, are stimulated by testosterone metabolites, notably dihydrotestosterone (Hewit 2001), but other hormones as well.




Fig. 1. Endocrine control of female reproduction and mechanism of action of progestins (A), GnRH-agonists (B), progesterone receptor inhibitors (C) and GnRH vaccines (D) (modified from Bertschinger et al. 2008b).

Recently new players in the control of reproduction have been discovered. Collectively known as kisspeptins and secreted by kisspeptin neurons in the hypothalamus these hormones control the activity of GnRH neurons. Furthermore, the kisspeptin neurons provide the links with the CNS, the environment, metabolic state and the reproductive system of an animal. Kisspeptins play a central role in puberty and

Wildlife fertility suppression southern Africa – a review

chronic administration of kisspeptin has been shown to induce precocious puberty in rats (Navarro et al. 2004). Kisspeptin administration in dogs (Albers-Wolthers et al. 2014) and ewes (Arreguin-Arevalo et al. 2007) resulted in LH peaks. Also, the photoperiod, which controls the release of melatonin from the pineal gland, is known to influence oestrous cycles of seasonal breeders. It is now established that melatonin exerts this effect via the kisspeptin neurons in the hypothalamus (Revel et al. 2007).

Hormonal methods of fertility suppression

Progestins or progestogens

Progestins have, for practical purposes, the same effect as progesterone on the hypothalamus and act by inhibiting the release of GnRH thus bringing about a downstream lack of FSH and LH release (Fig. 1A). As a result a new oestrous cycle is inhibited (anoestrus). A range of oral progestins were developed for use in the dog and cat (e.g. ovariid, megestrol acetate; Jöchle 2008) and the horse (altrenogest; Webel and Squires 1982). Some of these preparations are used alone, or in conjunction with other agents (e.g. deslorelin implants; Wright et al. 2001), to control reproduction of a variety of zoo species. Daily intake is required for to avoid a rebound release of GnRH. Thus the use in free-ranging wildlife is out of the question. Long-acting silicone implants or depot injections containing progestins like melengestrol acetate, have been extensively used to down-regulate female reproduction in wild felids such as lions and tigers, and some wild canid species. Although highly efficacious as contraceptive agents, their prolonged use resulted in a number of serious side-effects such as pyometra, which were life-threatening (Munson et al. 2005). Provided no serious pathology had occurred, the effect of the progestins could be reversed by removing the implant – a process which required immobilisation. Similar products regularly used for the contraception of females in a range of primate species with far less side effects.

GnRH super agonists

During the late 1990's, a new hormonal method of contraception became available. The GnRH analogue deslorelin, a so-called GnRH super-agonist, was manufactured in a slow-release formulation that released deslorelin for periods of 6 to 12 months, or even longer, and was designed for use in dogs and cats (Trigg et al. 2001; Munson et al. 2001). High continuous release of deslorelin from the subcutaneous biocompatible implant (Suprelorin, Virbac, France) acts by down-regulating the release of FSH and LH from the anterior pituitary (Fig. 1B). This in turn leads to down-regulation of gonadal activity and has the potential to be effective in both males and females. Following the successes achieved in male and female dogs and cats, preliminary trials were conducted in lionesses, male and female cheetahs, African wild dogs and leopards (Bertschinger et al. 2002). With the exception of wild dog females, a contraceptive efficacy of 100% was achieved. It is now the most commonly used method of contraception for captive and free-ranging carnivores and can be used in many primate and a range of other species. Following the administration of the implant an initial spike of FSH and LH is experienced before the pituitary gonadotrophs are down-regulated. In females this may induce a fertile oestrus in dogs (Trigg et al. 2001) and wild dogs (Bertschinger et al. 2001), but in large cats such as cheetahs and lions, only a short rise in oestrogens and progesterone, which does seem to be compatible with fertility, occurs (Bertschinger et al. 2002 and 2008). Nevertheless, many zoos will advocate suppression of oestrus with daily oral progestins starting 7 days prior to and continuing to 7 days after administering the implant (recommendations by the European Group on Zoo Animal Contraception). This regimen was first described for the domestic dog (Wright et al. 2001). In free-ranging cheetahs, lions and leopards, where this is not possible, we have never observed pregnancies following placement of the implants (Bertschinger et al. 2002 and 2008). Reconception after single treatments with 1 x 4.7 mg and 2 x 4.7 mg

Wildlife fertility suppression southern Africa – a review
implants occurs on average after 24 and 30 months in cheetahs (unpublished data) and lions (Bertschinger et al. 2008), respectively. Female lions, but not cheetahs, are prone to develop obesity after repeated treatments. We also demonstrated two reversals in one free-ranging female leopard following treatment with 1x 4.7 mg. Reconception occurred after 24 and 26 months respectively (unpublished data).




Fig. 2. Effects of annual 1 x 4.7 mg Suprelorin implants on mean testicular volume from day of first treatment (Year 0) up to Year 9. Numbers in brackets indicate number of observations for each year.

We have treated 66 male cheetah with 1 x 4.7 mg deslorelin implants on an annual basis and some for up to 9 years without any side effects being recorded. Down-regulation of fertility in males takes longer than in females with viable sperm being noted for up to six weeks post-treatment or longer (epididymal origin). During the first two years following annual treatment a rapid decrease in testicular volume to 59.6% and 47.5% of original volume, respectively is observed. This reflects a rapid decline in spermatogenesis. This was followed by a slower decline to reach between 29 to 24 % of the original volume after 5 to 9 years of annual treatment (Fig. 2). We have demonstrated return to fertility in one male treated annually for three years. He was released into the wild with two females and was able to mate successfully two and three

Wildlife fertility suppression southern Africa – a review
years after his last treatment. So far we have not been able to demonstrate reversal in
males treated annually for 5 years or longer (unpublished data).

Induction of abortion in some wild carnivores

Particularly when examining free-ranging carnivores for contraception, the pregnancy status of the animal will be unknown. If a female is found to be pregnant, we offer the owner the option of abortion, which can be combined with deslorelin treatment. We have used two methods in large carnivores: luteolytic prostaglandins and the progesterone receptor antagonist aglepristone (Fig. 1C).

Luteolytic prostaglandins

Initially we used dinoprost (Lutalyse, Zoetis, South Africa; Bertschinger et al. 2008) but now we have changed to cloprostenol (Estrumate, Intervet) which has much less effect on the smooth muscle of the GIT and is thus safer. Use one third of the cattle dose (8 mg dinoprost and 170 µg cloprostenol) for a lion or tiger. It is important to administer three doses with the same dose on three consecutive or alternate days (Table 2). The follow-up treatments can be administered with a drop-out dart. In African wild dogs the side effects of dinoprost are similar to those observed in domestic dogs. We thus advise against the use of this drug in wild dogs.

Table 2. Induction of abortion in lions and tigers with dinoprost (modified from Bertschinger et al. 2008).

Species	SPC nmol/l	Stage of gestation	Dinoprost treatment
Tiger	177.92	<3 weeks	7.5 mg on 3 alternate days; 2 weeks later
Lion	194.08	3 weeks	7.5 mg on 3 consecutive days from PD
Lion	107.29	3 weeks	7.5 mg on 3 consecutive days from PD
Tiger	63.70	70 days	7.5 mg on 3 alternate days from PD
Lion	212.80	80 days	7.5 mg on 3 consecutive days from PD

Aglepristone (Alizine®, Virbac, South Africa)

Alizine® is registered for use in domestic dogs in South Africa and the recommended dose to induce abortion is 1 ml/3 kg (30 mg aglepristone) repeated a day later. We used Alizine® as an extra-label drug to successfully induce abortion in African wild dogs and lions. Five three to four-year old captive wild dogs weighing 20 to 25 kg were each treated once with 8 ml Alizine® (approximately the same dose prescribed for the domestic dog). The Alizine® was injected intramuscularly in two sites using a pole-syringe while the dogs were trapped in a crush. Stage of pregnancy varied from 21 to 55 days. Abortion took place 5 to 7 days after treatment and no complications were observed.

Fourteen three to nine-year old lionesses weighing an estimated 150 to 210 kg and during the first to third trimester of gestation, were successfully aborted with Alizine®. The immobilised lionesses were given a single total dose of 35-45 ml each which is ≈70% of the dose recommended for the domestic dog. Abortion ensued after three to 5 days, no foetal remains were seen and only one female was seen with a serosanguineous vaginal discharge on Day 3 post treatment. All lionesses were immobilised 7 to 10 days after treatment for the purpose of contraception when they were all healthy.

Immunocontraception

Immunocontraception relies on carefully selecting peptides that are involved in critical steps of reproduction. Including such a peptide or peptides as antigens in a vaccine provokes the production of antibodies that neutralise the endogenous molecule or block a particular process. Examples of vaccines commonly in use are GnRH and native porcine zona pellucida (pZP) vaccines.

GnRH vaccines

The method involves the use of GnRH vaccines that stimulate the production of anti-GnRH specific antibodies. The antibodies neutralise endogenous GnRH and, in so doing, suppress down-stream endocrine mechanisms that control reproductive functions of both females (Fig. 1D) and males. GnRH immunocontraception was originally developed for the immunocastration of cattle (D'Occhio 1993) but one of the main reasons for further development of these vaccines was to use them as an alternative to surgical castration for the control of boar taint in pork (Dunshea et al. 2001) . GnRH vaccines have also, amongst others, been used to suppress fertility or reproductive behaviour in male feral pigs (Killian et al. 2006) and white tailed deer (Miller et al. 2000). In mares they have also been applied successfully to down-regulate fertility (Botha et al. 2008) and sex-related behaviour (Elhay et al. 2007).




Fig.3. A: Association between frequency of aggressive behaviours and faecal androgen metabolite concentrations in five African elephant bulls. B: Effect of three GnRH vaccinations (black arrows) on faecal androgen concentrations in an initially aggressive but non-musth African elephant bull (modified from De Nys et al. 2010).

The association between aggressive behaviour, especially during musth, and testosterone concentrations is clearly established in African elephant bulls. Bulls with raised testosterone levels and particularly during musth, when peak levels are reached, are very difficult to manage in captivity and also in smaller game reserves in South Africa. The positive relationship between the frequencies of aggressive behaviours and faecal androgen metabolites (FAM) is shown in Fig. 3A. In 2003 we initiated a study to see if testosterone and thus aggressive behaviour could be down-regulated in six bulls with GnRH vaccine treatments. The bulls that were aggressive (n=2) or in musth (n=1) at the start of treatment responded very well. Fig. 3B shows the effect of three treatments with a GnRH vaccine (Pepscan Systems, Lelystad, The Netherlands) on FAM concentrations of a problem bull (highly aggressive but not in musth) in one of the game reserves (De Nys et al. 2010). The bull in question, Thembo, is now in captivity and has been treated with a GnRH vaccine for the past 12 years. He is 32 years old and, since the start of treatment, has never shown musth or aggressive behaviour. Since our

Wildlife fertility suppression southern Africa – a review

initial trial in 2003 we have treated at least 40 bulls successfully to control aggressive behaviour and musth. In 2006 the GnRH vaccine Improvac® (Zoetis, South Africa) was registered for use in pigs in South Africa and, after we had established that it was also effective, it was routinely used to control aggressive behaviour and musth in elephant bulls (Bertschinger and Sills 2013).

Until four years ago, however, we did not know what effect GnRH vaccine would have on fertility of African elephant bulls. To investigate this we studied the effect of repeated Improvac® treatments on 11 captive and 2 free-ranging bulls (8 to 35 years of age) on semen quality and the internal reproductive organs. The bulls were examined and then treated twice with 5 ml (1000 µg) Improvac® (Day 0 and 35) and then boosted with the same dose every 5 months. Follow-up examinations were carried out at 6-monthly intervals for two to three years. After 6 months testicular size had reduced significantly and by the end of the study they were only 40% of the original area. Six months after the first vaccination ejaculates either contained dead sperm or were aspermic (Lueders et al. 2014). The trial clearly showed that the GnRH vaccine can be used as a contraceptive vaccine and at a fraction of the cost of vasectomy, which is a major surgical procedure in the elephant requiring highly specialised equipment. Currently we are monitoring some of the bulls to establish reversibility of the method on fertility. Furthermore, the method is being used in two game reserves for population control. Given the long intercalving interval in elephants (4-6 years), the results are not available yet.

During the period 2011-2015 we have also used Improvac® to control problem free-ranging male giraffes on six game reserves in South Africa. One adult bull (eight years old) and five three to four year-old pubertal males were treated. Typical problems associated with such males are that they attack motor vehicles, people on foot and giraffe calves that are sometimes their own offspring. Each male was darted

Wildlife fertility suppression southern Africa – a review
intramuscularly with two doses of Improvac® (800 µg each) five weeks apart. All males responded well with problem behaviour ceasing after the primary immunisation. So far, follow-up treatment has not been necessary in any of the males (unpublished data).

pZP immunocontraception

The zona pellucida capsule consists of 3 or 4 glycoproteins some of which are essential to the fertilisation process. Before a sperm can penetrate the zona capsule it must bind to a sperm receptor site. This process induces the acrosome reaction which, in brief, enables the penetration process. If a female is treated with the pZP vaccine she produces antibodies which bind to zona proteins surrounding the oocyte. The process is believed to block sperm-zona binding and thereby prevents fertilisation from taking place (Fig. 4). As long as antibody titres are sufficiently high, fertilisation will be




Fig. 4. Mechanism of action of the pZP vaccine

prevented but, because reproductive hormones are unaffected by this process, the female will continue to cycle normally. The origins of the pZP contraceptive vaccine

Wildlife fertility suppression southern Africa – a review date back to 1973 when researchers demonstrated that antibodies to pZP proteins could inhibit sperm-zona binding. In 1989 the first paper on contraception of domestic mares with pZP vaccine was published (Liu et al. 1989). Over the next 19 years, the pZP vaccine was used successfully as a contraceptive in feral mares and proved an effective means of limiting population growth. The success of the technique in wild horses led to its application to other wildlife species (≈ 100 species; Kirkpatrick et al. 2011). The vaccine can be delivered remotely and is both safe and reversible.

Although pZP immunocontraception was first tested on elephants in the Kruger National Park in 1996 (Fayer-Hosken et al. 1999; Fayer-Hosken et al. 2000) it was only from 2000 that the method was routinely used for population control of African elephants in The Makalali Game Reserve (Deslink et al. 2006; Deslink et al. 2007). When used in smaller reserves where elephant cows can be individually identified the method is 100% successful. It is safe to the cow, safe during pregnancy, has no effect on behaviour (Delsink et al. 2013) and is reversible, at least in the medium term (Bertschinger *et al.* 2012). The vaccine is made in our laboratory and is now supplied to 20 private and 5 national or provincial game reserves where, in total, about 700 cows are being treated for population control.

Conclusions

Contraception is an ideal solution for population control of free-ranging wildlife. Contrary to methods such as culling and removal of animals, both of which reduce population density, it does not stimulate reproductive rate. Here we presented one hormonal and two immunological methods that have been used for population and androgen-related behaviour control in some African species. The successes have mostly been good but it should be remembered that no method meets all the properties of an ideal contraceptive. Furthermore, although we have a range of effective methods, there

Wildlife fertility suppression southern Africa – a review is a need to critically investigate possible side effects in each species where they are applied. Apart from the obvious health issues, possible effects on social and territorial behaviour are very important and often neglected.

Selected References

- Albers-Wolthers KHJ, de Gier J, Kooistra HS, Rutten VPMG, van Kooten PJS, Leegwater PAJ, Millar RP, Schaefers-Okkens AC, 2014. Identification of a novel kisspeptin with high gonadotrophin stimulatory activity in the dig. Neuroendocrinology DOI: 10.1159/000364877: 1-11.
- Arreguin-Arevalo JA, Lents CA, Famerie TA, Nett TM, Clay CM, 2007. KiSS-1 peptide induces release of LH by a direct effect on the hypothalamus of ovariectomized ewes. Anim Reprod Sci **101**: 265-275.
- Bertschinger H J, Asa CS, Calle PP, Long JA, Bauman K, DeMatteo K, Jöchle W, Trigg TE, Human A, 2001. Control of reproduction and sex related behaviour in exotic wild carnivores with the GnRH analogue deslorelin: preliminary observations. J Reprod Fert **57**(Suppl.): 275–283.
- Bertschinger HJ, Trigg TE, Jöchle W. Human A, 2002. Induction of contraception in some African wild carnivores by down-regulation of LH and FSH secretion using the GnRH analogue deslorelin. Reproduction **60**: 41-52.
- Bertschinger HJ, Jago M, Nöthling JO, Human A, 2006. Repeated use of the GnRH analogue deslorelin to down-regulate reproduction in male cheetahs (*Acinonyx jubatus*). Theriogenology **66**: 1762-1767.
- Bertschinger HJ, De Barros Vaz Guimarães MA, Trigg TE, Human A, 2008a. The use of deslorelin implants for the long-term contraception of lionesses and tigers. Wildl Res **35**: 525-530.

Wildlife fertility suppression southern Africa – a review

- Bertschinger H, Delsink A, van Altena JJ, Kirkpatrick J, Killian H, Ganswindt A, Slotow R, Castley G, 2008b: Reproductive control of elephants. In: RJ Scholes RG and KG MennellKG (eds), Elephant management. A scientific assessment for South Africa. Wits University Press, 1 Jan Smuts Avenue, Johannesburg, pp. 357-328.
- Bertschinger HJ, Delsink AK, Kirkpatrick JF, van Altena JJ, Ahlers M, Dickerson T, Powrie D, Burger A, 2012. Porcine zona pellucida immunocontraception of African elephants (*Loxodonta africana*): beyond the experimental stage. In: Cain JW and Jason P. Marshal P (eds), IVth International Wildlife Management Congress, Durban, 9-12 July, pp. 95-102.
- Bertschinger HJ, Sills ES, 2013. Contraceptive applications of GnRH-analogs and vaccines for wildlife mammals of southern Africa: Current experience and future challenges. In: Sills ES (ed), Gonadotropin-releasing hormone (GnRH). Production, structure and function. Nova Science Publishers Inc., New York, ISBN: 978-1-62808-478-8 (eBook) 278 pp. 85-107
- Botha AE, Schulman ML, Bertschinger HJ, Guthrie AJ, Annandale CH, Hughes SB, 2008. The use of GnRH vaccine to suppress mare ovarian activity in a large group of mares under field conditions. *Wildl Res* **35**: 548- 554.
- Delsink AK, Kirkpatrick J, van Altena JJ, Bertschinger JJ, Ferreira SM, Slotow R, 2013. Lack of spatial and behavioral responses to immunocontraception application in African elephants (*Loxodonta africana*). *J Zoo Wildl Med* **44**(4S): S52–S74.
- Delsink AK, van Altena JJ, Grobler D., Kirkpatrick J., Bertschinger H, Slotow R, 2006. Regulation of a small, discrete African elephant population through immunocontraception in the Makalali Conservancy, Limpopo, South Africa. *SA J Sci* **102**: 403-405.

- Delsink AK, van Altena JJ, Grobler D, Bertschinger HJ, Kirkpatrick J, Slotow R, 2007. Implementing immunocontraception in free-ranging African elephants at Makalali Conservancy. *J SA Vet Ass* **78**: 25-30.
- De Nys HM, Bertschinger HJ, Turkstra JA, Colenbrander B, Palme R, Human AM, 2010. Vaccination against GnRH may suppress aggressive behaviour and musth in African elephant (*Loxodonta africana*) bulls – a pilot study. *J SA Vet Ass* **81**: 8-15.
- D’Occhio, MJ, 1993. Immunological suppression of reproductive functions in male and female mammals. *Anim Reprod Sci* **33**, 345–372.
- Dunsha FR, Colantoni C, Howard K, Mc Cauley I, Jackson P, Long KA, Lopaticki S, Nugent EA, Simons JA, Walker J, Hennessy DP, 2001. Vaccination of boars with GnRH vaccine (Improvac) eliminates boar taint and increases growth performance. *J Anim Sci* **79**, 2525–2535.
- Elhay M, Newbold A, Britton A, Turley P, Dowsett K, Walker J, 2007. Suppression of behavioural and physiological oestrus in the mare by vaccination against GnRH. *Aust Vet J* **85**, 39–45.
- Fayrer-Hosken RA, Bertschinger HJ, Kirkpatrick JF, Grobler D, Lambreski N, Honneyman G, Ulrcih T, 1999. Contraceptive potential of the porcine zona pellucida vaccine in the African elephant (*Loxodonta africana*). *Theriogenology* **52**: 835-846.
- Fayrer-Hosken, R.A., Grobler, D., van Altena JJ, Bertschinger HJ, Kirkpatrick JF, 2000. Immunocontraception of African elephants. *Nature* **407**: 149.
- Herbert, C.A., L. Vogelnest 2007. Catching kangaroos on the hop: development of a remote delivery contraceptive for marsupials. In: *Proceedings of 6th International Conference on Fertility Control for Wildlife*, 3–5 September, York, UK, p. 15.
- Hewit D 2001. Physiology and endocrinology of the male. In: Simpson G, England G, Harvey M (eds), *BSAVA Manual of small animal reproduction and neonatology*, BSAVA Ed. Pp. 62-69.

- Jöchle W, 2008. History of non-surgical contraception in dogs and cats. 2007. In: Schäffer J (ed), *Geschichte der Gynäkologie und Andrologie de Haustiere*, 14. Jahrestagung der DVG-Fachgruppe, 2-3 November, pp. 250-259.
- Killian G, Miller L, Rhyan J, Doten H, 2006. Immunocontraception of Florida feral swine with a single-dose GnRH vaccine. *Amer J Reprod Immun* **55**, 378–384.
- Kirkpatrick JF, Lyda RO, Frank KM, 2011. Contraceptive vaccines for wildlife: a review. *Amer J Reprod Immun* **66**: 40-50.
- Liu IKM, Bernoco M, Feldman M, 1989. Contraception in mares heteroimmunized with pig zonae pellucidae. *J Reprod Fert* **85**: 19-29.
- Lueders I, Young D, Botha S, Gray C, Luther I, Maree L, van der Horst G, Tindall B, Bertschinger HJ, 2014. GnRH vaccination of African elephant bulls for contraception and behavioural management. *Proceedings of International Conference on Diseases of Zoo and Wild Animals*, Warschau, Poland, 2014 p.10.
- Miller LA, Johns BE, Killian GJ, 2000. Immunocontraception of white-tailed deer with GnRH vaccine. *Amer J Reprod Immun* **44**, 266–274.
- Munson L, Bauman JE, Asa CS, Jöchle W, Trigg TE, 2001. Efficacy of the GnRH analogue deslorelin for suppression of oestrous cycles in cats. *J Reprod Fert* **57**: 269-273.
- Munson L, Moresco A, Calle PP, 2005. Adverse effects of contraceptives. In: Asa CS and Porton IJ, (eds), *Wildlife contraception*, The John Hopkins University Press, Baltimore: pp. 66-82.
- Navarro VM, Castellano JM, Fernandez-Fernandez R, Barreiro ML, Roa J, Sanchez-Criado JE, Aguilar E, Dieguez C, Pinilla L and TenaSempere M, 2004. Developmental and hormonally regulated messenger ribonucleic acid expression of KiSS-1 and its putative receptor, GPR54, in rat hypothalamus and potent luteinizing hormone-releasing activity of KiSS-1 peptide. *Endocrinology* **145**: 4565–4574.

Wildlife fertility suppression southern Africa – a review

- Revel FG, Ansel A, Klosen P, Saboureau M, Pévet P, Mikkelsen JD, Simonneaux V, 2007. Kisspeptin: A key to seasonal breeding. *Rev Endocr Metab Disord* **8** : 57-65.
- Trigg TE, Wright PJ, Armour AF, Williamson PE, Junaidi A, Martin GB, Doyle AG and Walsh J, 2001. Use of a GnRH analogue implant to produce reversible long-term suppression of reproductive function of male and female domestic dogs. *J Reprod Fert* **57**:255-261.
- Webel SK, Squires EL, 1982. Control of the oestrus cycle in mares with altrenogest. *J Reprod Fertil Suppl.*, **32**: 193-198.
- Wright PJ, Verstegen JP, Onclin K, Jochle W, Armour AF, Martin GB and Trigg TE, 2001. Suppression of the oestrous responses of bitches to the GnRH analogue