

**Characterizing the reproductive biology of the female pygmy hippopotamus  
(*Choeropsis liberiensis*) through non-invasive endocrine monitoring**

Gabriella L. Flacke <sup>a,b,\*</sup>, Franz Schwarzenberger <sup>c</sup>, Linda M. Penfold <sup>d</sup>, Susan L. Walker <sup>e</sup>,  
Graeme B. Martin <sup>a</sup>, Robert Peter Millar <sup>b,f,g</sup>, and Monique C. J. Paris <sup>a,b,f,h</sup>

<sup>a</sup> School of Animal Biology, University of Western Australia, 35 Stirling Highway, Crawley  
6009, Australia

<sup>b</sup> Institute for Breeding Rare and Endangered African Mammals (IBREAM), Edinburgh EH3  
6AT, United Kingdom

<sup>c</sup> Department of Biomedical Sciences, Unit of Physiology, Pathophysiology and  
Experimental Endocrinology, University of Veterinary Medicine (Vetmeduni Vienna),  
Veterinärplatz 1, 1210 Vienna, Austria

<sup>d</sup> South East Zoo Alliance for Reproduction & Conservation (SEZARC), 581705 White Oak  
Road, Yulee, FL 32097, USA

<sup>e</sup> Chester Zoo, Upton-by-Chester CH2 1LH, United Kingdom

<sup>f</sup> Mammal Research Institute and Centre for Neuroendocrinology, University of Pretoria,  
Department of Zoology and Entomology, Pretoria 0084, South Africa

<sup>g</sup> Institute for Infectious Diseases and Molecular Medicine, University of Cape Town, Cape  
Town 7925, South Africa

<sup>h</sup> College of Public Health, Veterinary and Medical Sciences, James Cook University,  
Townsville 4811, Australia

## Highlights

- Non-invasive endocrine monitoring demonstrated the estrous cycle to be an average of 31 days.
- The pygmy hippo under managed care is a spontaneous ovulator and a non-seasonally polyestrous species.
- Enzyme immunoassays that cross-react with pregnanediol can detect pregnancy from mid-gestation onward.
- Estrogen metabolite peaks were also observed during pregnancy and lactation, suggesting that follicular development occurs during both reproductive states.

**Corresponding author:**

Gabriella L. Flacke, DVM, MVSc, PhD; Department of Animal Health; Zoo Miami; 12400 SW 152<sup>nd</sup> Street; Miami, Florida; USA; Gabriella.Flacke@miamidade.gov

**Abstract**

The pygmy hippopotamus (*Choeropsis liberiensis*) is endangered in the wild and very little is known about its reproductive biology. In zoological facilities, this species experiences a number of reproductive issues that complicate breeding management, including a high rate of stillbirths and failure of many pairs to reproduce. We conducted a comprehensive study to evaluate reproductive cycles and pregnancy in this species using enzyme immunoassays (EIAs) for fecal hormone metabolite analysis. Fresh fecal samples were collected twice weekly for a one to three year period from 36 female pygmy hippos housed at 24 zoological institutions. Samples were analyzed in three separate laboratories. Three progestogen metabolite EIAs (*Pg-diol*: 5 $\beta$ -pregnane-3 $\alpha$ ,20 $\alpha$ -diol 3HS:BSA; *PdG*: pregnanediol-3-glucuronide R13904; *mono-P4*: Quidel clone 425) and three estrogen metabolite EIAs (*E2a*: estradiol-17 $\beta$ -OH 17-HS:BSA; *E2b*: estradiol 17 $\beta$  R0008; *E2c*: estradiol 17 $\beta$  R4972) accurately reflected reproductive events. Average estrous cycle length was  $31.8 \pm 7.4$  days based on estrogen metabolite peaks and  $30.9 \pm 7.3$  days based on nadir to nadir progestogen metabolite concentrations. Cyclical patterns in both estrogen and progestogen metabolites were detected throughout the year, indicating a lack of seasonality. Estrogen metabolite peaks were also observed during pregnancy and lactation, suggesting that follicular development occurs during both reproductive states. Pregnancy was most reliably demonstrated by elevation in progestogen metabolites (*Pg-diol* or *PdG*) in the second half of gestation. Average gestation length based on breeding to calving date was  $203 \pm 4$  days for 15

pregnancies. This comprehensive overview of the reproductive biology of the female pygmy hippo provides valuable data for guiding long-term breeding management for this endangered species and serves as a baseline for future studies addressing the potential influence of social structure, diet, body condition, and other husbandry factors on estrous cycling and reproduction.

**Keywords:** Enzyme-linked immunoassay (EIA); estrogen; estrous cycle; gestation; pygmy hippo; progestogens

## 1. Introduction

The pygmy hippopotamus (*Choeropsis liberiensis*) – hereafter referred to as pygmy hippo – is classified as endangered by the International Union for the Conservation of Nature [1] and is ranked 21<sup>st</sup> worldwide among mammals by Programme EDGE ([www.edgeofexistence.org](http://www.edgeofexistence.org)) as a priority for conservation action [2]. The pygmy hippo is endemic to the Upper Guinean Rainforest ecosystem in the West African countries of Côte d'Ivoire, Guinea, Liberia and Sierra Leone. Wild population size is uncertain but is estimated at < 2500 and is thought to be declining due to ongoing habitat loss and poaching [1,3]. Our understanding of the biology of this species in the wild is limited and data pertinent for developing effective conservation strategies are lacking. The first Conservation Strategy Action Plan was developed by the IUCN Pygmy Hippo Specialist Group in 2010 [3] and one of several research priorities was to characterize basic reproductive biology for both sexes.

There is no information concerning reproduction in wild pygmy hippos, but some general aspects of this species' reproductive biology are known from animals under managed care [4]. For example, both males and females reach sexual maturity between three and four years of age and can remain reproductively active into their third decade. The length of the estrous cycle, based on behavioral observations, ranges from 28 to 40 days. The gestation period is approximately 200 days after which a single calf is born; twin births are rare. There are no external signs of pregnancy except for enlargement of the mammary glands within a few days of parturition. The female is assumed to be polyestrous as births occur throughout the year in both northern and southern hemispheres, and conception is possible within a few weeks of perinatal mortality [5]. Historically, certain pairs have reproduced readily and often, and are therefore genetically over-represented in the managed population [4,5]. This trend continues

because several breeding pairs have failed to reproduce despite regular estrous behavior and/or mating being observed by husbandry staff. Other pairs have repeatedly experienced perinatal calf mortality or stillbirth [5]. These issues, together with a high neonatal mortality rate (> 30%), have limited the success of captive breeding and could reduce the long-term genetic diversity of the managed population, especially as imports from the wild ceased after 1982.

Finding solutions to these problems is hampered by a poor basic knowledge of the reproductive biology in this species, particularly the absence of hormone patterns. A clear way forward is to use enzyme immunoassays (EIAs) for the non-invasive assessment of endocrine processes that present insights into reproductive biology and improved breeding management [6]. These assays were originally established to quantify native hormones in serum or urine and exhibit varying degrees of cross-reactivity with the multitude of hormone metabolites excreted in the feces. As a result, measuring fecal hormone metabolites using these assays can prove challenging for some taxa, such as different species of tapir (J. Brown, personal communication, 2015), due to limited cross-reactivity between available antibodies and that species' particular repertoire of metabolites. In these scenarios collaborations between institutions are essential, especially where endangered species with limited numbers in the *ex situ* population are concerned, because not all endocrine laboratories will have access to the same selection of EIAs.

Endocrine assessment of reproductive events in female pygmy hippos is limited to one study, with only two females, that used a radio-immunoassay to analyze progesterone in skin secretions and saliva over a six-month period; the authors reported an average estrus cycle length of 26 days [7]. For the common hippo (*Hippopotamus amphibius*), on the other hand,

non-invasive methods for confirming pregnancy, monitoring the reproductive cycle, pinpointing estrus, and identifying the timing of puberty have already been established [8–10]. However, this information cannot be directly extrapolated to the pygmy hippo because the reproductive physiology and steroid hormone metabolites of even closely related species can differ markedly. For example, different species of rhinoceros, felids, and ursids exhibit different estrous cycle characteristics and produce variable types and amounts of fecal estrogen and progesterone metabolites [11]. Thus, non-invasive protocols for evaluating reproductive events in the pygmy hippo are needed to facilitate estrous cycle monitoring, pregnancy diagnosis and prediction of parturition, especially as repeated blood sample collection is difficult.

Therefore, our overall objective in the present study was to characterize the basic reproductive biology of female pygmy hippos under managed care using fecal hormone metabolite analysis to define endocrine profiles during the estrous cycle, pregnancy and lactation. Specific aims were to: 1) demonstrate biological relevance for measuring fecal metabolites of estrogen and progestogens via enzyme immunoassays; 2) establish a standardized method for pregnancy detection; 3) determine the length of the estrous cycle via physiologic rather than behavioral assessment; 4) test for seasonality of the estrous cycle.

## **2. Materials and Methods**

### *2.1. Animals and sample collection*

Thirty-six female pygmy hippos from 13 European and nine North American zoological institutions were included in this study; 33 were sexually mature ( $\geq 3$  y) at the time sampling

commenced (Appendix I). Fecal samples were collected twice weekly and stored frozen at –20°C until extraction and analysis. Occasionally, biweekly sampling was not possible and the interval between samples was up to two weeks. Reproductive events, including behavioral estrus, mating and parturition, were recorded in conjunction with the timing of fecal sample collection (Appendix I).

## 2.2. *Gastrointestinal transit time*

The lag time between patterns of steroid secretion in the blood and subsequent metabolite excretion in the feces is correlated with gastrointestinal transit time [12,13]. We determined gastrointestinal transit time for one male and one female pygmy hippo housed at the same facility and fed the same diet. We used an easily identifiable fecal marker (glitter; Sulyn Industries, Coral Springs, Florida, USA), as previously described in similar studies investigating transit time [14,15], mixed with grain and fed to each hippo. We recorded the time from ingestion until the first and last passage of glitter in the feces for both hippos.

## 2.3. *Reproductive hormone metabolite analysis*

Hormone analysis was conducted in three separate laboratories (designated Lab A, Lab B, and Lab C; see Appendix I) using previously established fecal hormone metabolite extraction and EIA techniques. Modifications to these techniques are described below.

### *2.3.1. Fecal hormone extraction*

Lab A (Vienna, Austria) performed fecal extraction as previously described [16]. Briefly, 0.5 g wet fecal material and 0.5 mL water were mixed with 4 mL reagent grade methanol and vortexed for 30 min. After centrifugation to remove solid fecal material, 1 mL of the methanol solution was transferred to a clean vial and mixed with 0.25 mL of a 5% NaHCO<sub>3</sub> solution and 5 mL diethyl ether. The mixture was then vortexed for 30 s, centrifuged at 1200 g for 10 min, and the extract supernatant held at -20°C for 30 min. Finally, the supernatant ether phase was placed in a clean vial, evaporated to dryness, and re-suspended in 0.5 mL assay buffer.

Lab B (Chester, UK) performed fecal extraction with methods adapted from Walker et al. [17]. Following manual homogenization of the sample, 0.5 g of wet fecal material was mixed with 4.5 mL methanol (reagent grade, Sigma-Aldrich, Dorset, UK) and 0.5 mL of distilled water. Samples were vortexed for 30 s, rotated continuously on an orbital shaker at room temperature overnight and centrifuged the following day for 20 min at 598g. The supernatant was air-dried in a water bath at 55°C in a fume cupboard and then re-suspended in 1 mL of methanol.

Lab C (SEZARC, Yulee, Florida, USA) performed fecal extraction using previously described methods [18]. Briefly, after manual homogenization of the sample, 0.5 g wet fecal material was mixed with 4 mL methanol (reagent grade, Fisher Scientific, Fair Lawn, NJ, USA) and 1 mL reverse osmosis-purified water. Samples were then shaken in a Glas-Col Large Capacity Mixer (Glas-Col LLC, Terre Haute, IN, USA) for 20 min at 90 rpm followed by centrifugation for 10 min at 3100 rpm. Then 200µL of the supernatant was transferred into new vials containing 800µL assay buffer and stored at -20°C until analysis.

### 2.3.2. *Enzyme immunoassays (EIAs) & validation*

Italicized terms are those we have used to refer to the individual hormone assays throughout the remainder of our study. Cross-reactivities for all of the assays are provided in Appendix II. All samples, controls, and standards were assayed in duplicate.

Lab A analyzed fecal extract aliquots using previously performed group-specific EIAs with rabbit-origin polyclonal antibodies against the following: *i*) 20-oxo-pregnane ( $5\alpha$ -pregnane- $3\beta$ -ol-20-one 3HS:BSA [19]); *ii*) pregnanediol (*Pg-diol*:  $5\beta$ -pregnane- $3\alpha,20\alpha$ -diol 3HS:BSA [20]); *iii*) total estrogen (*E2a*: estradiol- $17\beta$ -OH 17-HS:BSA [21]). The intra- and inter-assay coefficients of variation for all assays were  $< 10\%$  and  $< 15\%$ , respectively. The minimum assay sensitivities were 2.6 pg/well for 20-oxo-pregnane, 5.5 pg/well for Pg-diol, and 0.5 pg/well for E2a. Serial dilutions of fecal extracts yielded displacements curves parallel to the standards curves for all three EIAs. Depending on the concentration in the sample, the dilution factors for the samples were between 1:10,000 and 1:125,000 for 20-oxo-pregnane, Pg-diol and E2a. As the data for the 20-oxo-pregnane assay were not biologically relevant, it was not used for further sample analysis.

Lab B analyzed all fecal extract aliquots using a monoclonal progesterone antibody (*mono-P4*: Quidel clone 425 (CL425)) and an estrogen metabolite antibody (*E2b*: estradiol  $17\beta$  R4972) developed by Coralie Munro, University of California, Davis, USA [22]. The intra- and inter-assay coefficients of variation for all assays were  $< 10\%$  and  $< 15\%$ , respectively. The minimum assay sensitivities were 0.78 pg/well for mono-P4 and 1.25 pg/well for E2b. Assays were validated by demonstrating parallelism between standard curves and serial dilutions of fecal extracts: CL425 ( $r^2 = 0.980$ ,  $F_{1,7} = 293$ ,  $P < 0.001$ ); E2b ( $r^2 = 0.991$ ,  $F_{1,7} =$

809,  $P < 0.001$ ). Dilutions were run at 1:256, 1:512, 1:1024, 1:2048, 1:4096, 1:8192, 1:16,386 and 1:32,768 for mono P4 and E2b. There was no evidence of matrix interference, demonstrated by significant recovery of pooled fecal extract added to the standards: CL425 (observed =  $4.13 + 0.98$  [expected],  $r^2 = 0.999$ ,  $F_{1,7} = 8694$ ,  $P < 0.001$ ); E2b (observed =  $7.86 + 2.19$  [expected],  $r^2 = 0.986$ ,  $F_{1,7} = 514$ ,  $P < 0.001$ ). Both the mono-P4 and the E2b EIAs were biologically validated and used for sample analysis at Lab B.

Lab C analyzed fecal extracts using a monoclonal progesterone antibody (*mono-P4*: Quidel clone 425), a polyclonal progesterone antibody (*poly-P4*: R4859), a pregnanediol antibody (*PdG*: pregnanediol-3-glucuronide, R13904) and an estrogen metabolite antibody (*E2c*: estradiol 17 $\beta$  R0008), all developed by Coralie Munro, University of California, Davis, CA, USA. A double-antibody EIA protocol was used as previously described [18,23]. The intra- and inter-assay coefficients of variation for all assays were  $< 10\%$  (average 2.58%) and  $< 20\%$  (average 8.07%), respectively. The minimum assay sensitivities, determined at 90–95% binding, were 10.1 pg/well for mono-P4, 11.0 pg/well for poly-P4, 18.6 pg/well for PdG, and 4.04 pg/well for E2c. Serially diluted fecal extracts yielded parallel displacement curves to the standard curves in all antibodies tested: mono-P4 ( $r^2 = 0.801$ ,  $F_{1,7} = 14.1$ ,  $P < 0.001$ ); poly-P4 ( $r^2 = 0.844$ ,  $F_{1,7} = 19.0$ ,  $P < 0.001$ ); PdG ( $r^2 = 0.874$ ,  $F_{1,7} = 24.3$ ,  $P = 0.001$ ); E2c ( $r^2 = 0.831$ ,  $F_{1,7} = 17.3$ ,  $P = 0.002$ ). Dilutions were run at 1:30, 1:60, 1:120, 1:240, 1:480, 1:960 and 1:1920 for mono-P4 and poly-P4, and at 1:16; 1:32, 1:64, 1:128; 1:256, 1:512, and 1:1024 for PdG. An antibody accuracy check showed that recovery of known amounts of progesterone (0.19–100 ng/mL) added to pools of diluted extracted feces was 107% at 1:100 for PdG ( $y = 1.030x + 0.002$ ,  $r^2 = 0.998$ ,  $F_{1,17} = 5397$ ). Recovery of known amounts of estrogen (0.039–10.0 ng/mL) was 119% at 1:50 for E2c ( $y = 1.607x + 9.381$ ,  $r^2 = 0.992$ ,  $F_{1,15} = 714$ ). The mono-P4 and poly-P4 EIAs were not biologically validated as neither clearly

demonstrated a rise in progestogen metabolites during pregnancy. Thus, recovery was not assessed and neither EIA was investigated further at Lab C.

#### 2.4. *Data analysis*

We plotted hormone concentrations over the sampling period (1 to 3 y) for each pygmy hippo. We first calculated non-pregnant baseline values for each EIA by averaging the lowest 10% of the samples for each female. We subsequently calculated concentrations for each EIA as a percent of each female's baseline. We characterized progestogen metabolite profiles by examining luteal phases and estrogen metabolite profiles by observing estrogen peaks indicative of follicular phases. We defined a luteal phase as any period when progestogen immunoreactivity was at least 50% above baseline for >14 days. We defined the peak luteal values for each cycle as the highest progestogen metabolite level during the cycle, and the nadir luteal concentration as progestogen metabolite level closest to baseline on either side of the peak. We defined an estrogen metabolite peak as the highest value within a group of samples that was greater than 10 days apart and greater than 2.5-fold (250%) above baseline for E2a and greater than 2-fold (200%) above baseline for E2b and E2c. We then determined the length of the estrous cycle by *i*) the interval between two estrogen metabolite peaks or *ii*) the onset of one luteal phase until the onset of the next. For some profiles we used a 3-point moving averaging to reduce background noise and clarify peak and nadir patterns; however, the values for actual peak and nadir metabolite levels were calculated from the non-averaged data for each female. Due to the numerous sources of variability associated with measuring immunoreactive metabolites in fecal material, in some cases we made subjective observations to distinguish between true cyclic patterns and random fluctuations in the data. We also assessed estrous cycle length based on direct observations of mating or behavioral by

husbandry staff and compared these data (when available) with the hormone metabolite profiles for each hippo.

We calculated the length of gestation from the date of a confirmed mating event until parturition; for some pregnancies, the date of mating was unknown and we were unable to determine the length of gestation. We evaluated both progestogen and estrogen metabolites before, during and after gestation to determine if there is follicular activity during pregnancy, if there is a period of lactational anestrus, and when the first post-partum estrus occurs. For some study females, we could only generate partial pregnancy profiles as the sampling periods did not overlap with the full length of gestation. We calculated the mean and standard deviation ( $\pm$  SD) for estrogen and progestogen metabolite concentrations during pregnancy, in the non-pregnant state overall, and during peak luteal phase by combining data for all females. The data for pregnancy comprised all hormone concentrations from mating until parturition; we calculated the overall mean as well as a separate mean for the first and the second half of gestation (E2a & Pg-diol:  $n = 7$  pregnancies,  $n = 256$  samples; E2b & mono-P4:  $n = 3$  pregnancies,  $n = 96$  samples; E2c & PdG:  $n = 8$  pregnancies,  $n = 341$  samples). The data for non-pregnancy included all hormone concentrations throughout the estrous cycle and during periods of acyclicity (E2a & Pg-diol:  $n = 16$  hippos,  $n = 1683$  samples; E2b & mono-P4:  $n = 9$  hippos,  $n = 719$  samples; E2c:  $n = 8$  hippos,  $n = 321$  samples; PdG:  $n = 16$  hippos,  $n = 1237$  samples). The data for peak luteal phase comprised the highest concentration of progestogen metabolites during each luteal phase for each female (Pg-diol:  $n = 15$  hippos,  $n = 174$  samples; mono-P4:  $n = 7$  hippos,  $n = 58$  samples; PdG:  $n = 15$  hippos,  $n = 120$  samples). We first used Shapiro-Wilk tests for normality and then used either paired  $t$ -test analysis (two groups) or ANOVA (more than two groups) to compare estrogen and progestogen

immunoreactivity in the different reproductive states (pregnancy, non-pregnant, and peak luteal phase).

### **3. Results**

#### *3.1. Gastrointestinal transit time*

The time between consumption of the fecal marker and the first passage of glitter in the feces was 20 h for the female and 30 h for the male. Peak glitter excretion occurred between 30 and 46 h for the female and between 48 to 70 h for the male. Thus, gastrointestinal transit time for pygmy hippos can vary between individuals fed the same diet, but ranges from one to three days. Hormone metabolite levels measured in the feces are therefore likely to represent endocrine events that occurred in the previous 24 to 48 h. However, for simplification of results, reproductive events are reported as if they occurred on the same day as the feces were voided. Behavioral reports of estrus up to 72 h before endocrine events indicative of estrus were considered temporally associated.

#### *3.2. Reproductive patterns – gestation and the post-partum period*



Mean  $\pm$  SD, minimum and maximum hormone metabolite levels for each EIA are presented in Table 1 for both pregnant and non-pregnant pygmy hippos. Mean progestogen metabolite concentrations were higher during pregnancy than in the non-pregnant state for all three progestogen EIAs (Pg-diol,  $n = 16$  hippos,  $P < 0.001$ ; PdG,  $n = 16$  hippos,  $P < 0.001$ ; mono-P4,  $n = 9$  hippos,  $P < 0.001$ ). Compared to baseline for each female, concentrations during pregnancy were up to 144-fold higher for Pg-diol, up to 9-fold higher for PdG, and up to 12-

**Table 1.** Mean ( $\pm$  SD) values for minimum, maximum, and baseline concentrations (per g feces) of reproductive hormone metabolites in female pygmy hippos during pregnancy and the estrous cycle. Results are for six different EIAs assayed in three separate laboratories (A, B and C).

	<b>Lab A</b>	<b>Lab B</b>	<b>Lab C</b>	<b>Lab A</b>	<b>Lab B</b>	<b>Lab C</b>
	<b>Pg-diol</b>	<b>mono-P4</b>	<b>PdG</b>	<b>E2a</b>	<b>E2b (R4972)</b>	<b>E2c (R0008)</b>
	<b><math>\mu\text{g/g}</math></b>	<b><math>\text{ng/g}</math></b>	<b><math>\mu\text{g/g}</math></b>	<b><math>\text{ng/g}</math></b>	<b><math>\text{ng/g}</math></b>	<b><math>\text{ng/g}</math></b>
Pregnant	14.3 $\pm$ 9.4	1137 $\pm$ 319	5.3 $\pm$ 1.6	1.3 $\pm$ 0.4	121 $\pm$ 13.9	611 $\pm$ 341
Range	0.19 – 59.8	313 – 4187	1.16 – 19.8	0.01 – 17.5	15.6 – 218	132 – 2922
<i>n</i> pregnancies	7	3	8	7	3	8
Non-pregnant	3.2 $\pm$ 1.7	917 $\pm$ 271	3.2 $\pm$ 1.1	1.3 $\pm$ 0.7	74.7 $\pm$ 24.2	388 $\pm$ 186
Range	0.02 – 26.9	50 – 7084	0.66 – 13.9	0.09 – 43.8	17.1 – 261	43 – 1052
<i>n</i> hippos	16	9	16	16	9	8
Cycle length (days)	30.2 $\pm$ 7.4	31.8 $\pm$ 5.9	30.7 $\pm$ 8.7	31.2 $\pm$ 7.1	32.6 $\pm$ 7.6	–
Range	14 – 46	17 – 42	14 – 49	14 – 48	17 – 46	–
<i>n</i> cycles	157	50	80	147	34	–
Peak luteal	6.2 $\pm$ 3.5	2150 $\pm$ 682	4.7 $\pm$ 1.1	–	–	–
Range	0.97 – 26.9	683 – 7084	2.3 – 9.9	–	–	–
Nadir luteal	1.3 $\pm$ 0.8	498 $\pm$ 185	1.8 $\pm$ 0.5	–	–	–
Range	0.12 – 4.8	171 – 1180	0.66 – 3.18	–	–	–
Peak follicular	–	–	–	4.5 $\pm$ 2.9	142 $\pm$ 47.7	–
Range	–	–	–	0.55 – 43.8	57.0 – 262	–
Baseline	0.93 $\pm$ 0.61	354 $\pm$ 162	1.9 $\pm$ 0.6	0.30 $\pm$ 0.10	44.2 $\pm$ 14.6	252 $\pm$ 173
Range	0.27 – 2.8	152 – 733	1.1 – 3.2	0.14 – 0.56	25.5 – 62.1	99 – 618
<i>n</i> hippos	16	9	16	16	9	8

fold higher for mono-P4. Concentrations of Pg-diol and PdG excreted during the second half of pregnancy were significantly higher than those excreted during the first half (Fig. 1). Additionally, during the first half of gestation Pg-diol ( $P = 0.143$ ) and PdG ( $P = 0.131$ ) were similar to peak luteal phase concentrations, whereas during the second half of gestation they were significantly higher than peak luteal values for both Pg-diol (up to 9-fold higher,  $n = 7$  pregnancies,  $P < 0.001$ ) and PdG (up to 4-fold higher,  $n = 8$  pregnancies,  $P = 0.010$ ). On the other hand, no gestational difference was noted for mean concentrations of mono-P4 excreted throughout pregnancy, and mean values were significantly lower those excreted during peak luteal phase ( $n = 3$  pregnancies,  $P < 0.001$ ).



The average length of gestation for 15 pregnancies ( $n = 12$  hippos) was  $203 \pm 4$  days. Progesterone metabolite concentrations (Pg-diol and PdG) rose to more than 50% above baseline values within  $17 \pm 15$  days after mating and conception. Concentrations fluctuated considerably during the first half of gestation but overall exhibited a slow, steady increase followed by a discernable decrease mid-gestation, around 90 to 100 days post-conception. This decrease was followed by a prominent rise in progesterone metabolites throughout the second half of gestation, reaching peak levels (mean = 31-fold above baseline values) just before parturition followed by a rapid return to baseline values within  $13 \pm 8$  days after parturition (Fig. 1). These patterns were readily evident in some pregnancies (Fig. 2a), whereas in others there was considerable variation, complicating interpretation (Fig. 2b). In the latter situation a three-point moving average helped to clarify these patterns. The gestational patterns observed with Pg-diol and PdG were not as clearly discernable with mono-P4; however, this EIA was only used for two pregnancies (Fig. 4).



**Fig. 1.** Mean fecal progestogen metabolite concentrations before, during, and after gestation in the pygmy hippo; standard errors are not shown for clarity. Each data point comprised samples for all females spanning a 3 d time frame (between five and eight samples per mean). The data are aligned with parturition (Day 0); mating and conception occurred at an average of  $-203$  d.

**A.** PdG (pregnanediol-3-glucuronide); mean values for  $n = 8$  pregnancies.

**B.** Pg-diol ( $5\beta$ -pregnane- $3\alpha,20\alpha$ -diol 3HS:BSA); mean values for  $n = 7$  pregnancies. A nadir is noted at mid-gestation for both profiles, followed by a marked increase through the second half of gestation until parturition.



**Fig. 2.** Individual profiles for fecal PdG (pregnanediol-3-glucuronide; ●) and E2c (Estradiol 17 $\beta$  R0008; ○) immunoreactivity for two pygmy hippos; the data are aligned with parturition (Day 0).

**A.** SB No. 1063, “Chomel.” Example of a profile that demonstrates a gradual increase in progestogen metabolite concentrations after conception with a visible decline around mid-gestation (–100 d from parturition) followed by a substantial increase during the second half of gestation and a subsequent return to baseline shortly after parturition. Several estrogen metabolite peaks are also evident during the first half of gestation. The baseline for both PdG and E2c are indicated by the solid horizontal line. Gestation was 199 days.

**B.** SB No. 1169, “Violet.” Example of a pregnancy profile that does not demonstrate clear patterns in progestogen metabolites. Estrogen metabolite peaks are again noted during the first half of gestation. The baseline for PdG is indicated by a solid horizontal line; the baseline for E2c is indicated by a dotted horizontal line. Gestation was 210 days.

Estrogen values varied considerably during gestation, but some trends were evident. There was no difference between overall mean pregnant and non-pregnant concentrations of estrogen metabolites for E2a ( $n = 16$  hippos,  $P = 0.319$ ) whereas for E2b and E2c, overall concentrations were higher during pregnancy than in non-pregnant hippos ( $n = 10$  hippos,  $P < 0.001$ ). There was evidence of follicular activity as demonstrated by clear peaks in estrogen metabolite concentrations, especially during the first half of gestation, with a final peak just before or at the time of parturition in several females (e.g. Fig. 3a & b). In other females, clear peaks were not discernable, but values remained above baseline for extended periods during gestation, although in some cases the sampling interval was greater than twice weekly (e.g. Fig. 4b) and thus may not accurately reflect estrogen metabolite patterns. An estrogen metabolite peak prior to mid-gestation was evident in four of the five females with E2a data that spanned the mid-gestation time frame (e.g. Fig. 3a). For E2c pregnancy profiles, a similar peak just prior to mid-gestation was noted for three of seven females (e.g. Fig. 3c). Only two pregnancy profiles were analyzed with E2b and neither showed a mid-gestation peak (Fig. 4).



Average estrogen metabolite concentrations during gestation were not statistically different for male versus female calves for the E2a EIA ( $P = 0.087$ ); however, concentrations tended to be lower for pregnancies with male calves. Concentrations were significantly different during gestation for the E2c EIA ( $P = 0.002$ ); lower for male calves, higher for female calves. The

sex ratio and number of samples during pregnancy for male and female calves was similar for E2a (3 ♂, 4 ♀) and slightly skewed toward females for E2c (3 ♂, 5 ♀).

The first post-partum estrus, demonstrated by an estrogen metabolite peak indicative of follicular activity (e.g. Fig. 3b), was observed from 22 to 92 days (mean  $40 \pm 21$  d) after parturition with viable calves ( $n = 10$ ). However, only seven of these post-partum follicular peaks were followed by a luteal phase indicative of ovulation. It is unknown whether these were fertile cycles as the male was always kept separate from the female and calf. For two females, the first post-partum estrogen metabolite peak was followed by an additional period of lactational anestrus (61 d and 99 d); the other eight females continued to cycle after the first post-partum estrus event. Hormonal evidence of estrous cycling resumed well before the calf was weaned for all 10 females. Two calves experienced perinatal mortality; in these cases estrus and mating occurred at 14 and 16 days post-partum and neither female conceived.

### 3.3. *Reproductive patterns – estrous cycle*




Mean ( $\pm$  SD) values for nadir and peak luteal concentrations of progestogens and peak follicular concentrations of estrogens are presented in Table 1. The overall mean estrous cycle length based on the interval between follicular peaks was  $31.8 \pm 7.4$  days ( $n = 185$  cycles;  $n = 20$  hippos). Peak estrogen metabolite levels compared to baseline varied considerably between and among individual hippos. The overall mean estrous cycle length



**Fig. 3.** Individual profiles for fecal mono-P4 (Quidel clone 425; ●) and E2b (estradiol-17β R4972; ○) for two pygmy hippos before, during and after pregnancy; the data are aligned with parturition (Day 0). The baseline for mono-P4 is indicated by a solid horizontal line and for E2b by a dotted horizontal line.

**A.** SB No. 993, “Debby.” Progestogen metabolite concentrations rise above baseline immediately after mating with several initial peaks and remain elevated until parturition followed by an immediate return to baseline. Estrogen metabolite concentrations are already well above baseline at the time of mating and continue to increase until mid-gestation (–90 d) followed by a rapid decline and fluctuating lower levels until parturition. Gestation was 210 days.

**B.** SB No. 938, “Wendy.” Progestogen metabolite concentrations also begin to rise above baseline immediately after mating but return to baseline at mid-gestation (–125 d and –100 d), then remain elevated until parturition followed by a return to baseline within 25 days. Estrogen metabolite concentrations fluctuate throughout gestation but generally remain above baseline with a peak just before (–11 d) and at parturition (Day 0). Gestation was 200 days.



**Fig. 4.** Individual profiles for fecal progestogen and estrogen metabolite profiles during gestation for three pygmy hippos. The baseline for both progestogen and estrogen metabolites is indicated by the horizontal line; the data are aligned with parturition (Day 0).





**A.** SB No. 993, “Debby.” Pg-diol (pregnanediol-3-glucuronide; ●) and E2a (estradiol-17β-OH 17-HS:BSA; ○); progestogen metabolite concentrations remain above baseline from early gestation until parturition. An estrogen metabolite peak is noted at the time of mating (consistent with estrus), at parturition, and at various points throughout gestation, A peak occurs just prior to mid-gestation (−107 d) that temporally correlates with a marked decline in progestogen metabolites.

**B.** SB No. 1033, “Torpedo.” Pg-diol (pregnanediol-3-glucuronide; ●) and E2a (estradiol-17β-OH 17-HS:BSA; ○); a significant increase in progestogen metabolite concentrations above baseline does not occur until the second half of gestation. Estrogen metabolite peaks are again noted at the time of mating, throughout gestation including just before parturition, and at approximately at 22 days post-partum, indicative of post-partum estrus.

**C.** SB No. 1143, “Fitri.” PdG (pregnanediol-3-glucuronide 13904; ●) and E2c (Estradiol 17β R0008; ○); progestogen metabolite concentrations begin to rise at mid-gestation (−101 d) and remain elevated through parturition followed by a return to baseline by nine days post-partum. Estrogen metabolite concentrations fluctuate throughout gestation, with several prominent peaks.

based on the interval between luteal phases was  $30.9 \pm 7.3$  days ( $n = 287$  cycles,  $n = 30$  hippos). The overall mean estrous cycle length based on behavioral observations was  $32.4 \pm 6.3$  days ( $n = 42$  cycles,  $n = 13$  hippos). There was hormonal evidence of estrous cycling during all months of the year. Consecutive, year-round estrous cycles were observed for 13 females (age range 3 to 37 y), 10 from temperate climates where the animals were housed indoors during the colder months and three from sub-tropical climates where they were housed outdoors year-round. For the remaining 23 females in this study these trends could not be assessed, either because a portion of the data was collected during gestation and lactation, or the sampling time frame(s) did not include a contiguous 12-month period.

Follicular estrogen metabolite peaks or luteal progestogen metabolite nadirs were associated with observed estrous behavior in 31/46 cases (67%) when behavioral estrus was reported by the husbandry staff (e.g. Fig. 5b). In the other 15 cases, behavioral estrus was reported but hormone metabolite levels did not indicate estrus (e.g. Fig. 5c). Behavioral estrus was not reported for the majority of physiologic estrous cycles demonstrated by follicular (139/185) or luteal phases (241/287). In most cases, we were unable to determine if this disagreement was due to a lack of reporting or an actual absence of behavioral signs of estrus because behavioral data were not provided for most females (e.g. Fig 5d). However, for some females, the husbandry staff ensured us they had not noted any signs of behavioral estrus yet patterns for estrogen and/or progestogen metabolites indicated clear estrous cycling (e.g. Fig. 5a).



**Fig. 5.** Representative profiles from individual pygmy hippos demonstrating estrous cycles via peak estrogen metabolite concentrations or nadir progesterone metabolite concentrations. Data are presented as percentage of baseline (100%). The horizontal line indicates 250% baseline (2.5-fold increase) for estrogen metabolites; vertical arrows denote behavioral estrus events; horizontal arrows denote a luteal phase.

**A.** SB No. 1182, “Ashaki.” Estrous cycling indicated by peaks in E2a (estradiol-17 $\beta$ -OH 17-HS:BSA) immunoreactivity; there was no behavioral evidence of estrus during the sampling period.

**B.** SB No. 548, “Tana.” Estrous cycling indicated by peaks in E2a immunoreactivity; there are three points where behavioral estrus corresponds with an estrogen peak.

**C.** SB No. 864, “Kelsey.” Luteal phases indicated by nadirs in PdG (pregnanediol-3-glucuronide 13904) immunoreactivity; there are two behavioral/hormonal matches (arrow with \*) and the remaining behavioral estrus events do not correlate with a luteal nadir.

**D.** SB No. 1148, “Krakunia.” Luteal phases indicated by nadirs in mono-P4 (Quidel clone 425) immunoreactivity; no data were provided concerning behavioral estrus. This female was housed at a zoo without a male; the data support spontaneous ovulation in the absence of a male.

Our study included four hippos aged 30 to 38 y at the start of the sample collection period; all four showed evidence of either intermittent or continuous estrous cycling. There were three juvenile hippos aged 2 to 2.5 y at the start of the study period and each showed evidence of intermittent cycles from the age of three onward. Only five adult females were housed at a facility without a male; the remaining females were housed at least adjacent to a male and in many cases the male was continuously together with the female (Appendix I). In all five females housed in complete absence of a male, there was clear evidence of estrous cycling as indicated by luteal phases (e.g. Fig. 5d).

There was marked individual variation among hormonal patterns for non-pregnant, non-lactating, sexually mature females ( $n = 26$  hippos). Seventeen of these animals exhibited near-continuous estrous cycling throughout the study period. The remainder exhibited variable periods of acyclicity (8/26); one adult female (age 7 y) did not show any patterns consistent with estrous cycling throughout the 12-month sampling period and neither behavioral estrus nor mating was observed.

#### **4. Discussion**

We have demonstrated biological relevance for a number of EIAs for measuring fecal metabolites of estrogen and progestogens in pygmy hippos and have employed these techniques to characterize several important aspects of female reproductive biology.

#### 4.1. Gestation and the post-partum period

Pregnancy diagnosis was possible from mid-gestation onward (approximately 100 days before parturition) using an EIA that cross-reacts with  $5\beta$ -pregnanediol metabolites of progesterone (Pg-diol, PdG). Progestogen metabolite concentrations significantly exceeded peak luteal values throughout the second half of gestation, in contrast with the first half of gestation where values fluctuated considerably and were similar to luteal values. It is therefore necessary to analyze fecal samples for at least four months after an observed mating event to avoid a false negative pregnancy diagnosis. With the mono-P4 EIA, progestogen metabolite concentrations did not rise above peak luteal levels at any point during gestation and only exhibited a sustained rise above baseline in one female, so we do not recommend using this assay to diagnose pregnancy in pygmy hippos. This finding is in contrast to the common hippo, where mono-P4 immunoreactivity was elevated throughout gestation at levels significantly greater than during luteal phases [8]. We noted a general trend of increasing estrogen metabolite concentrations during the second half of gestation, but there was considerable variation between individuals. Based on our results, fecal estrogen metabolites cannot be used to reliably diagnose pregnancy in the pygmy hippo.

The decrease in progestogen metabolite concentrations around mid-gestation followed by a marked increase above luteal values until parturition suggests a change in the dynamics of progestogen secretion. There are two possible explanations for these patterns, although both need verification as a priori hypotheses. First, the follicular activity noted during gestation for several pygmy hippos in our study, especially during the first 100 days before the decline and subsequent rise of progestogen metabolite concentrations, may indicate the formation of accessory corpora lutea (CLs), similar to the domestic mare [24]. A fluctuating pattern in serum estrogen concentrations during the first half of gestation, possibly indicative of accessory follicle formation, has also been observed in some lowland tapirs (*Tapirus*

*terrestris*) [25]. Second, the marked increase in progestogens in the second half of pregnancy may be indicative of a luteal-placental shift. This phenomenon has been documented via non-invasive endocrine monitoring in a number of other large mammalian species, including the domestic mare [26]. Other well-studied examples are the three rhinoceros species most commonly kept in zoos (black *Diceros bicornis*, white *Ceratotherium simum* and greater one-horned *Rhinoceros unicornis*), and the okapi (*Okapia johnstoni*), where fecal progestogen metabolites remain at luteal phase concentrations during the initial part of pregnancy but increase dramatically between three and five months after conception, signifying a luteal-placental shift [16,20,27–30].

Estrogen levels during gestation tended to being higher for pregnancies with female calves, although we cannot conclusively state if there is a sex specific difference, as has been reported in other species such as orcas (*Orcinus orca*) [31] and red-fronted lemurs (*Eulemur fulvus rufus*) [32]. In contrast to the trend in pygmy hippos, estrogen levels during gestation were higher in mother's carrying male calves for both of these species.

There were 10 females in our study with viable calves and sufficient data to determine timing of the first post-partum estrous cycle; seven of these females had hormonal evidence of an estrous cycle approximately 30 days after calving. Although weaning age in the wild is unknown, if the female can become pregnant approximately one month after parturition and carry a second calf during lactation, then weaning in wild pygmy hippos may occur around eight months of age (one month post-partum estrus followed by a seven month gestation). Calving intervals from studbook data indicate that the female is able to conceive on the first post-partum estrus, within one month of parturition, when the calf experiences neonatal mortality [5]. In the common hippo, lactational anestrus averaged 34 weeks; however it was

not always observed and was inconsistent between and among individuals, possibly indicating the influence of external factors, such as photoperiod or body condition, on the timing of the first post-partum estrus [8].

#### 4.2. *Estrous cycle*

All of the estrogen metabolite EIAs we used in this study detected patterns indicative of follicular development. Similarly, all three progestogen assays (Pg-diol, PdG, mono-P4) demonstrated hormone metabolite patterns consistent with luteal activity. However, the mono-P4 EIA most clearly demonstrated luteal phases with minimal background fluctuations. Mean estrous cycle length was 31 to 32 days, slightly shorter than the mean cycle length of 35 days reported for the common hippo [8]. Cycle length ranged from 14 to 49 days and exhibited considerable variability both between and among individual animals.

For animals with longer cycles, our biweekly sampling protocol may have missed some estrogen peaks indicating follicular development. Although we were able to identify 185 follicular phases with this protocol, we could not determine the precise duration of these phases as the majority were represented by a single estrogen metabolite peak. However, the follicular phase likely spans at least three to four days or else biweekly sampling would not have detected the associated endocrine patterns. Nadir luteal concentrations may likewise have been missed with biweekly sampling as the progestogen metabolite nadir was typically also short, with only one or two points at or near baseline before the next luteal phase. On the other hand, the pygmy hippo may actually exhibit both short and long cycles, similar to the Malayan tapir (*Tapirus indicus*) [25], the white rhino [29,33,34] and the black rhino [34,35,36]. The only previous reproductive hormone study for the pygmy hippo reported an

average cycle length of 26 days based on salivary progesterone metabolites; the data were for a total of 10 cycles from two females [7]. However, these authors noted hormonal oscillations every nine days, leading them to hypothesize a normal cycle length of 18, 27, or 36 days.

Our ability to detect estrous cycles has allowed us to determine several aspects of the basic reproductive biology of the female pygmy hippo under managed care. We noted hormonal evidence of estrous cycling between 2.5 and 3 y for the three juvenile females, indicating that the onset of sexual maturity occurs within this time frame. This is slightly earlier than the 3 to 4 y reported for the onset of sexual maturity in common hippos based on non-invasive endocrine monitoring [10]. Our data also indicate that females continue to exhibit hormonal evidence of estrous cycles well into their 30s. Behavioral estrus and mating was observed for one aged (>30 y) pygmy hippo on several occasions and hormone profiles also provided evidence of estrous cycles; however, she never conceived. Within the overall Studbook population, the oldest multiparous female to successfully reproduce was 37 years old at the time of conception, but generally reproduction in females is uncommon beyond the age of 30 [5].

In accordance with the Studbook [5], and similar to the common hippo [8], our results also indicate the pygmy hippo to be non-seasonally polyestrous in both subtropical as well as temperate climates. Additionally, similar to domestic ruminants [37], pygmy hippos in the managed population exhibit spontaneous ovulation, as demonstrated by clear hormonal evidence of estrous cycles in females housed in facilities without a male. Although we have no comparative information from wild populations, the ecologic life history traits of the pygmy hippo largely support non-seasonally polyestrous reproduction. The species is endemic to a very limited region of tropical, equatorial West African rainforest where

temperature and photoperiod remain relatively constant throughout the year, although there is a seasonal difference in rainfall [38].

#### 4.3. *General discussion*

Interestingly, the mono-P4 EIA was able to detect luteal phases but not pregnancy. This assay predominately cross-reacts with 20-oxo-progesterone metabolites, so its spectrum of detection is notably different to the Pg-diol and PdG EIAs that primarily detect metabolites with 20 $\alpha$ -hydroxyl groups (Appendix II). In order for two EIAs with distinctly different cross-reactivities to accurately reflect two separate phases of gestation, namely the post-ovulatory CL and the feto-placental unit, the predominant progesterones produced by each must be structurally dissimilar. This phenomenon has most clearly been demonstrated in the mare where  $\Delta$ 4 progesterones are produced in the first part of gestation by the primary and accessory CLs, as described by Legacki et al. [39]. Thereafter, an increase in 5 $\alpha$ -dihydroprogesterone (a 5 $\alpha$ -reduced pregnane) marks the luteo-placental shift, and metabolites of 5 $\alpha$ -reduced pregnanes, produced by the feto-placental unit, predominate in mid to late gestation [39]. A similar transition in the principal type of progesterone produced by the CL versus the feto-placental unit in pygmy hippos may explain why Pg-diol and PdG immunoreactivity increased in the second half of gestation and mono-P4 did not, remaining at or below luteal levels.

Temporally-associated estrous behavior was not reported for the majority of estrous cycles identified via endocrine monitoring. Dathe and Kuckelkorn [7] also noted a lack of behavioral signs during periods of physiologic estrus for the two females in their study. These findings indicate that behavioral evaluation, even when performed by experienced zoo staff,

is not an infallible method of estrus detection. A comparable phenomenon of ‘silent’ estrus has been reported in other species, for example the greater one-horned rhino [40] and the Baird’s tapir (*Tapirus bairdii*) [41], further emphasizing the value of endocrine monitoring as an important tool to guide breeding programs for endangered species. Additionally, data from two sets of females ( $n = 4$  hippos) at the same zoo ( $n = 2$  zoos) suggest that pygmy hippos housed in close proximity possibly synchronize their cycles as follicular activity exhibited temporal association on a number of occasions. Thus, behavioral signs of estrus may be more difficult to identify in subdominant females when multiple females are housed together or in close proximity.

When behavioral signs of estrus are evident, they are likely to be a true indication of physiologic estrus as the majority (67%) of behavioral reports temporally coincided with an estrogen metabolite peak or progesterone metabolite nadir. Thus, husbandry staff should be consistently trained in recognizing and correctly classifying these behaviors, which include male vocalization, increased activity level, and direct interest in the female; females may exhibit a deep, audible breathing pattern [7,42]. Possible explanations for the 33% of cases where estrus-associated behaviors were reported but expected endocrine patterns were not detected include incorrect classification of behaviors or sampling frequency. The lack of an estrogen metabolite peak before all luteal phases could also be due to the biweekly sampling schedule occasionally missing these peaks. Reproductive behavior and mating in the absence of endocrine evidence of estrus has been reported in the white rhino [29] and was noted for one female in our study.

There were occasional estrogen metabolite peaks during luteal phases; these may represent follicular waves, similar to the patterns seen in domestic animals [37]. Alternatively, they

may be an indication of estrogen production by the CL; however this phenomenon has only been described in humans and non-human primates [43,44]. We also intermittently noted temporally coinciding estrogen and progesterone metabolite peaks; this finding may represent secretion of progesterones from a non-ovulatory follicle undergoing atresia. Some individuals had periods of regular estrous cycling with clear follicular peaks and luteal phases followed by periods where physiologically relevant patterns were not discernable. This phenomenon may represent periods of normal cyclicity interspersed with irregular ovarian activity, possibly associated with environmental factors, captivity-associated stress [45], underlying reproductive pathology, or persistent luteal activity, as recently characterized in the domestic mare [46].

Estrous cycle patterns exhibited considerable variability both among and between individuals. A similar phenomenon has been reported for other pachyderms, including elephants and rhinoceroses, with potential links to social stress and other captivity-related factors [11,35,47]. Certain husbandry variables, including diet, body condition, and social grouping might also influence estrous cycling and reproduction in pygmy hippos. Dietary differences not only affect the amount of undigested fiber and the water content of the feces, but phytoestrogens and soy-based pelleted feeds could also influence hormone levels and cause significant reproductive abnormalities, especially in herbivores [48]. Diet also has a direct influence on body condition, and a number of pygmy hippos in zoological collections are overweight [4]. In Asian elephants (*Elephas maximus*), obesity has been proposed as a primary cause of reproductive problems in captivity, including a high incidence of stillbirth [49]. In zoo-housed black rhinos, higher body condition may play a role in reduced reproductive success for some females [35]. Overweight pygmy hippos may also experience impaired reproductive function.

Direct and indirect (i.e., camera trap photos, footprint tracking) observation of wild pygmy hippos has repeatedly demonstrated these animals to be predominately solitary [50–54]. Separate housing except during breeding therefore more closely reflects this species' natural biology; however, logistical constraints often preclude this arrangement and many zoos keep their pygmy hippo pair together continuously (see Appendix I). Abnormal social structure in zoo environments has been linked to reproductive abnormalities in several species, for example the cheetah (*Acinonyx jubatus*) [55,56], African elephant (*Loxodonta africana*) [47], and red river hog (*Potamochoerus porcus*) [57]. Chronic stress is hypothesized to be the primary link between unnatural social structure and resulting reproductive abnormalities and has been shown to negatively influence reproductive parameters in dairy cows [22], domestic ewes [58] and male cheetah [59]. Poor reproductive success and abnormal estrous cycling in zoo-born white rhinos is hypothesized to be associated with stress and undetermined captivity-associated factors [11,29,34,60,61]. These findings raise concern for captivity-induced modulation of reproductive performance, a phenomenon that may also impact pygmy hippos.

#### 4.4. *Future research and recommendations*

Additional investigations are warranted to address a number of unanswered questions concerning the reproductive biology of the female pygmy hippo. We recommend performing ultrasound examination in conjunction with endocrine monitoring to correlate reproductive events with fecal hormone metabolite profiles and to further validate physiologic relevance. Ultrasound can also be used to determine if additional CLs are formed during gestation and to identify the timing of CL regression as an indication of luteal-placental shift. Further research

is also needed to establish if there is more than one distinct estrous cycle length in the pygmy hippo, if longer periods of luteal activity are abnormal, and if the failure of some pairs to reproduce despite endocrine evidence of (often irregular) ovarian activity is indicative of underlying reproductive pathology as has been documented for the elephant and white rhino [62,63]. Additionally, future studies can employ the techniques we have described for monitoring reproductive hormones to investigate potential links between zoo-specific husbandry variables, obesity, chronic stress, and reproduction in pygmy hippos.

Saliva sampling, piloted in two pygmy hippos by Dathe and Kuckelkorn [7] has been successfully employed for endocrine monitoring in numerous other wildlife species [28,64–66] and further research may demonstrate it to be a valuable tool in pygmy hippos as well. Determining the biochemical structure of immunoreactive metabolites of estrogen and progesterone in the feces would help further refine which EIAs most accurately reflect reproductive hormones for this species; possibilities include high performance liquid chromatography (HPLC) and mass spectrometry [67]. Radio-label studies would also be useful to determine what percentage of reproductive hormone metabolites are excreted via the urinary and fecal route and to more precisely determine lag time. However, this approach is not always practical or safe, and is often not justified in rare and endangered species. Thus, non-invasive endocrine monitoring paired with ultrasound validation of physiologic events is the next-best approach for tracking reproductive events.

## **5. Conclusions**

This landmark collaborative study sets a significant milestone in elucidating reproductive endocrinology of the female pygmy hippo and provides essential data for future research. We

describe endocrine patterns during pregnancy, lactation and the estrous cycle and have identified several EIAs with biological relevance for non-invasively monitoring reproductive hormone metabolites in the feces. Additionally, we have identified two progestogen metabolite EIAs (Pg-diol & PdG) that can detect pregnancy in the second half of gestation. We have also established that the pygmy hippo under managed care is a spontaneous ovulator and a non-seasonally polyestrous species. The length of the estrous cycle as determined by endocrine analysis was slightly shorter than the length reported from behavioral observations. Results were repeatable and consistent between laboratories using slightly different EIAs and techniques. Our data provide a vital resource for further investigations of reproductive physiology in the pygmy hippo and demonstrate how non-invasive endocrine monitoring can be a valuable tool for characterizing normal biology. These same technologies can potentially help diagnose reproductive abnormalities and improve future breeding management of this endangered species.

### **Acknowledgements**

We would like to express our gratitude to all of the institutions that participated in this multi-year study: Aalborg Zoo; Arnhem Royal Burger's Zoo; Baton Rouge Zoo; Bristol Zoo; Chicago Zoological Society, Brookfield Zoo; Colchester Zoo; Dierenpark Wissel, Epe; Dvůr Králové Zoo Safari; Edinburgh Zoo; Gaia Zoo, Kerkrade; Gladys Porter Zoo, Brownsville; Jackson Zoo; Louisville Zoo; Marwell Zoo; Omaha's Henry Doorly Zoo and Aquarium; Paris Zoological Park; Parken Zoo, Eskilstuna; Rotterdam Zoo; Rum Creek Center for Conservation of Tropical Ungulates; Tampa's Lowry Park Zoo; Zoo Basel; Zoo Berlin; Zoo Miami; and ZooParc Overloon. Our special appreciation goes out to the husbandry staff that collected the samples; without their help and dedication we would not have been able to

conduct this research. Additionally, we would like to thank Cayman Adams, Sarah Allred, Kim Daly-Crews, Thijs van den Houten, Saleha Khan, Elke Leitner, Lara Metrione, Erich Möstl, Paige Pickering, Kayla Weller, and the staff at the University Medical Center Groningen, The Netherlands for technical support.

Funding: This work was supported by the American Association of Zoo Veterinarians Wild Animal Health Fund; the Center for Conservation of Tropical Ungulates; Dr. Mervyn Jacobson; Givskud Zoo; Omaha's Henry Doorly Zoo and Aquarium; the Royal Zoological Society of Scotland; Zodiac Zoos; Zoo Basel; the University of Western Australia (UWA), Convocation Postgraduate Research Travel Award; UWA Postgraduate Student's Association Fieldwork and Data Collection Award; and a UWA Graduate Research School PhD Completion Scholarship.

## References

- [1] Ransom C, Robinson PT, Collen B. *Choeropsis liberiensis*. IUCN Red List Threat Species 2015 2015. doi:e.T10032A18567171. Accessed 22.12.2015.
- [2] Isaac NJB, Turvey ST, Collen B, Waterman C, Baillie JEM. Mammals on the EDGE: Conservation priorities based on threat and phylogeny. PLoS One 2007;2:e296. doi:10.1371/journal.pone.0000296.
- [3] Mallon D, Wightman C, De Ornellas P, Ransom C. Conservation Strategy for the Pygmy Hippopotamus. Gland, Switzerland & Cambridge, UK: IUCN Species Survival Commission; 2011.
- [4] Flacke GL, Chambers BK, Martin GB, Paris MCJ. The pygmy hippopotamus *Choeropsis liberiensis* (Morton, 1849): Bringing to light research priorities for the largely forgotten, smaller hippo species. Der Zool Garten NF 2015;84:234–65.
- [5] Steck B, editor. International Studbook for the Pygmy Hippopotamus 2015. 22nd ed. Basel: Zoo Basel, Switzerland; 2016.
- [6] Schwarzenberger F. The many uses of non-invasive faecal steroid monitoring in zoo and wildlife species. Int Zoo Yearb 2007;41:52–74.
- [7] Dathe HH, Kuckelkorn B. Progesteronnachweis in Sekreten des Zwergflußpferdes (*Choeropsis liberiensis* Morton, 1844). Der Zool Garten NF 1989;59:201–8.
- [8] Graham LH, Reid K, Webster T, Richards M, Joseph S. Endocrine patterns associated with reproduction in the Nile hippopotamus (*Hippopotamus amphibius*) as assessed by fecal progestagen analysis. Gen Comp Endocrinol 2002;128:74–81.
- [9] Smith TE, Richards M, Joseph S, Savage A. Endocrine determinants of pregnancy in the Nile hippopotamus (*Hippopotamus amphibius*). Proc. 2nd Annu. Symp. Zoo Res., Paignton Zoo Environmental Park, Paignton, Devon, United Kingdom: 2000, p. 187–90.

- [10] Wheaton CJ, Joseph S, Reid K, Webster T, Richards M, Savage A. Body weight as an effective tool for determination of onset of puberty in captive female Nile hippopotami (*Hippopotamus amphibious*). *Zoo Biol* 2006;25:59–71.
- [11] Schwarzenberger F, Brown JL. Hormone monitoring: An important tool for the breeding management of wildlife species. *Wien Tierarztl Monatsschr* 2013;100:209–25.
- [12] Palme R, Fischer P, Schildorfer H, Ismail MN. Excretion of infused <sup>14</sup>C-steroid hormones via faeces and urine in domestic livestock. *Anim Reprod Sci* 1996;43:43–63.
- [13] Schwarzenberger F, Möstl E, Palme R, Bamberg E. Faecal steroid analysis for non-invasive monitoring of reproductive status in farm, wild and zoo animals. *Anim Reprod Sci* 1996;42:515–26.
- [14] Hulsman A, Dalerum F, Ganswindt A, Muenscher S, Bertschinger HJ, Paris M. Non-invasive monitoring of glucocorticoid metabolites in brown hyaena (*Hyaena brunnea*) feces. *Zoo Biol* 2011;30:451–8.
- [15] Van der Weyde LK, Martin GB, Blackberry MA, Gruen V, Harland A, Paris MCJ. Reproductive hormonal patterns in pregnant, pseudopregnant and acyclic captive African wild dogs (*Lycaon pictus*). *Anim Reprod Sci* 2015;156:75–82.
- [16] Schwarzenberger F, Rietschel W, Vahala J, Holeckova D, Thomas P, Maltzan J, et al. Fecal progesterone, estrogen, and androgen metabolites for noninvasive monitoring of reproductive function in the female Indian rhinoceros, *Rhinoceros unicornis*. *Gen Comp Endocrinol* 2000;119:300–7.
- [17] Walker SL, Waddell WT, Goodrowe KL. Reproductive endocrine patterns in captive female and male red wolves (*Canis rufus*) assessed by fecal and serum hormone analysis. *Zoo Biol* 2002;21:321–35.

- [18] Metrione LC, Norton TM, Beetem D, Penfold LM. Seasonal reproductive characteristics of female and male Jackson's hartebeest (*Alcelaphus buselaphus jacksoni*). *Theriogenology* 2008;70:871–9.
- [19] Schwarzenberger F, Tomášová K, Holečková D, Matern B, Möstl E. Measurement of fecal steroids in the black rhinoceros (*Diceros bicornis*) using group-specific enzyme immunoassays for 20-oxo-pregnanes. *Zoo Biol* 1996;15:159–71.
- [20] Schwarzenberger F, Francke R, Göldenboth R. Concentrations of faecal immunoreactive progestagen metabolites during the oestrous cycle and pregnancy in the black rhinoceros (*Diceros bicornis michaeli*). *J Reprod Fertil* 1993;98:285–91.
- [21] Patzl M, Schwarzenberger F, Osmann C, Bamberg E, Bartmann W. Monitoring ovarian cycle and pregnancy in the giant anteater (*Myrmecophaga tridactyla*) by faecal progestagen and oestrogen analysis. *Anim Reprod Sci* 1998;53:209–19.
- [22] Walker SL, Smith RF, Jones DN, Routly JE, Dobson H. Chronic stress, hormone profiles and estrus intensity in dairy cattle. *Horm Behav* 2008;53:493–501.
- [23] Barnes SA, Teare JA, Staaden S, Metrione L, Penfold LM. Characterization and manipulation of reproductive cycles in the jaguar (*Panthera onca*). *Gen Comp Endocrinol* 2016;225:95–103.
- [24] Brinsko SP, Blanchard TL, Varner DD, Schumacher J, Love CC. *Manual of Equine Reproduction*. 3rd ed. Maryland Heights, Missouri: Mosby, Inc.; 2010.
- [25] Pukazhenti B, Quse V, Hoyer M, van Engeldorp Gastelaars H, Sanjur O, Brown JL. A review of the reproductive biology and breeding management of tapirs. *Integr Zool* 2013;8:18–34.
- [26] Schwarzenberger F, Möstl E, Bamberg E, Pammer J, Schmechlik O. Concentrations of progestagens and oestrogens in the faeces of pregnant Lipizzan, trotter and thoroughbred mares. *J Reprod Fertil Suppl* 1991;44:489–99.

- [27] Brown JL, Wasser SK, Wildt DE, Graham LH, Monfort SL. Faecal steroid analysis for monitoring ovarian and testicular function in diverse wild carnivore, primate, and ungulate species. *J Mamm Biol* 1997;62:27–31.
- [28] Czekala NM, Callison L. Pregnancy diagnosis in the black rhinoceros (*Diceros bicornis*) by salivary hormone analysis. *Zoo Biol* 1996;15:37–44.
- [29] Patton ML, Swaisgood RR, Czekala NM, White AM, Fetter GA, Montagne JP, et al. Reproductive cycle length and pregnancy in the southern white rhinoceros (*Ceratotherium simum simum*) as determined by fecal pregnane analysis and observations of mating behavior. *Zoo Biol* 1999;18:111–27.
- [30] Schwarzenberger F, Rietschel W, Matern B, Schaftenaar W, Bircher P, Van Puijenbroeck B, et al. Noninvasive reproductive monitoring in the okapi (*Okapia johnstoni*). *J Zoo Wildl Med* 1999;30:497–503.
- [31] Robeck TR, Steinman KJ, O'Brien JK. Characterization and longitudinal monitoring of serum androgens and glucocorticoids during normal pregnancy in the killer whale (*Orcinus orca*). *Gen Comp Endocrinol* 2017;247:116–29.
- [32] Ostner J, Heistermann M. Endocrine characterization of female reproductive status in wild redfronted lemurs (*Eulemur fulvus rufus*). *Gen Comp Endocrinol* 2003;131:274–83.
- [33] Schwarzenberger F, Walzer C, Tomasova K, Vahala J, Meister J, Goodrowe KL, et al. Faecal progesterone metabolite analysis for non-invasive monitoring of reproductive function in the white rhinoceros (*Ceratotherium simum*). *Anim Reprod Sci* 1998;53:173–90.
- [34] Brown JL, Bellem AC, Fouraker M, Wildt DE, Roth TL. Comparative analysis of gonadal and adrenal activity in the black and white rhinoceros in North America by noninvasive endocrine monitoring. *Zoo Biol* 2001;20:463–86.

- [35] Edwards KL, Shultz S, Pilgrim M, Walker SL. Irregular ovarian activity, body condition and behavioural differences are associated with reproductive success in female eastern black rhinoceros (*Diceros bicornis michaeli*). *Gen Comp Endocrinol* 2015;214:186–94.
- [36] Garnier JN, Holt W V., Watson PF. Non-invasive assessment of oestrous cycles and evaluation of reproductive seasonality in the female wild black rhinoceros (*Diceros bicornis minor*). *Reproduction* 2002;123:877–89.
- [37] Evans ACO. Characteristics of ovarian follicle development in domestic animals. *Reprod Domest Anim* 2003;38:240–6.
- [38] Critical Ecosystem Partnership Fund. Ecosystem Profile - Upper Guinean Forest Ecosystem of the Guinean Forests of West Africa Biodiversity Hotspot. 2000.
- [39] Legacki EL, Scholtz EL, Ball BA, Stanley SD, Berger T, Conley AJ. The dynamic steroid landscape of equine pregnancy mapped by mass spectrometry. *Reproduction* 2016;151:421–30.
- [40] Stoops MA, West GD, Roth TL, Lung NP. Use of urinary biomarkers of ovarian function and altrenogest supplementation to enhance captive breeding success in the Indian rhinoceros (*Rhinoceros unicornis*). *Zoo Biol* 2014;33:83–8.
- [41] Brown JL, Citino SB, Shaw J, Miller C. Endocrine profiles during the estrous cycle and pregnancy in the Baird's Tapir (*Tapirus bairdii*). *Zoo Biol* 1994;13:107–17.
- [42] Lang EM. Das Zwergflußferd. Wittenberg Lutherstadt: A. Ziemsen Verlag, DDR; 1975.
- [43] Baird DT, Baker TG, McNatty KP, Neal P. Relationship between the secretion of the corpus luteum and the length of the follicular phase of the ovarian cycle. *J Reprod Fertil* 1975;45:611–9.
- [44] Devoto L, Henríquez S, Kohen P, Strauss JF. The significance of estradiol metabolites

- in human corpus luteum physiology. *Steroids* 2017;123:50–4.
- [45] Mason GJ. Species differences in responses to captivity: Stress, welfare and the comparative method. *Trends Ecol Evol* 2010;25:713–21.
- [46] Santos VG, Bettencourt EM V., Ginther OJ. Long-term characteristics of idiopathic persistent corpus luteum in the mare. *Theriogenology* 2015;84:242–51.
- [47] Brown JL, Paris S, Prado-Oviedo N, Meehan C, Hogan JN, Morfeld K, et al. Reproductive health assessment of female elephants in North American zoos and association of husbandry practices with reproductive dysfunction in African elephants (*Loxodonta africana*). *PLoS One* 2016;11:e0145673.  
doi:10.1371/journal.pone.0145673.
- [48] Adams NR. Detection of the effects of phytoestrogens on sheep and cattle. *J Anim Sci* 1995;73:1509–15.
- [49] Taylor VJ, Poole TB. Captive breeding and infant mortality in Asian elephants: A comparison between twenty western zoos and three eastern elephant centers. *Zoo Biol* 1998;17:311–32.
- [50] Bülow W. Untersuchungen am Zwergflußpferd, *Choeropsis liberiensis* im Azagny - Nationalpark, Elfenbeinküste. Diplomarbeit, Zoologischen Institut Braunschweig, 1987.
- [51] Collen B, Howard R, Konie J, Daniel O, Rist J. Field surveys for the endangered pygmy hippopotamus *Choeropsis liberiensis* in Sapo National Park, Liberia. *Oryx* 2011;45:35–7.
- [52] Conway AL. Conservation of the Pygmy Hippopotamus (*Choeropsis liberiensis*) in Sierra Leone, West Africa. PhD Thesis, University of Georgia, Athens, 2013.
- [53] Hentschel KM. Untersuchung zu Status, Ökologie und Erhaltung des Zwergflusspferdes (*Choeropsis liberiensis*) in der Elfenbeinküste. Doktorand

- Dissertation, Technischen Universität Carolo-Wilhelmina, Braunschweig, 1990.
- [54] van Heukelum M. In search of the elusive Pygmy Hippo; Establishment of methods to determine population structure of Pygmy Hippos in Tai National Park, and assessment of their role in seed dispersal. MSc Thesis, Wageningen University, 2011.
- [55] Brown JL. Female reproductive cycles of wild female felids. *Anim Reprod Sci* 2011;124:155–62.
- [56] Wielebnowski NC, Ziegler K, Wildt DE, Lukas J, Brown JL. Impact of social management on reproductive, adrenal and behavioural activity in the cheetah (*Acinonyx jubatus*). *Anim Conserv* 2002;5:291–301.
- [57] Jarboe M, Adams C, Bahr JM, Penfold L, Newell-Fugate AE. Boar urine pheromone exposure modifies estrous cycle length and regularity in same-sex housed female red river hogs (*Potamochoerus porcus*). *Proc. 5th Int. Soc. Wildlife Endocrinol. Conf.*, Berlin, Germany: 2015, p. 10.
- [58] Pierce BN, Clarke IJ, Turner AI, Rivalland ETA, Tilbrook AJ. Cortisol disrupts the ability of estradiol-17 $\beta$  to induce the LH surge in ovariectomized ewes. *Domest Anim Endocrinol* 2009;36:202–8.
- [59] Terio KA, Marker L, Munson L. Evidence for chronic stress in captive but not free-ranging cheetahs (*Acinonyx jubatus*) based on adrenal morphology and function. *J Wildl Dis* 2004;40:259–66.
- [60] Carlstead K, Brown JL. Relationships between patterns of fecal corticoid excretion and behavior, reproduction, and environmental factors in captive black (*Diceros bicornis*) and white (*Ceratotherium simum*) rhinoceros. *Zoo Biol* 2005;24:215–32.
- [61] van der Goot AC, Martin GB, Millar RP, Paris MCJ, Ganswindt A. Profiling patterns of fecal 20-oxopregnane concentrations during ovarian cycles in free-ranging southern white rhinoceros (*Ceratotherium simum simum*). *Anim Reprod Sci* 2015;161:89–95.

- [62] Hermes R, Hildebrandt TB, Göritz F. Reproductive problems directly attributable to long-term captivity—asymmetric reproductive aging. *Anim Reprod Sci* 2004;82–83:49–60.
- [63] Hermes R, Hildebrandt TB, Walzer C, Göritz F, Patton ML, Silinski S, et al. The effect of long non-reproductive periods on the genital health in captive female white rhinoceroses (*Ceratotherium simum simum*, *C.s. cottoni*). *Theriogenology* 2006;65:1492–515.
- [64] Dathe HH, Kuckelkorn B, Minnemann D. Salivary cortisol assessment for stress detection in the Asian elephant (*Elephas maximus*): A pilot study. *Zoo Biol* 1992;11:285–9.
- [65] Heintz MR, Santymire RM, Parr LA, Lonsdorf E V. Validation of a cortisol enzyme immunoassay and characterization of salivary cortisol circadian rhythm in chimpanzees (*Pan troglodytes*). *Am J Primatol* 2011;73:903–8.
- [66] Illera J-C, Silván G, Cáceres S, Carbonell M-D, Gerique C, Martínez-Fernández L, et al. Assessment of ovarian cycles in the African elephant (*Loxodonta africana*) by measurement of salivary progesterone metabolites. *Zoo Biol* 2014;33:245–9.
- [67] Handelsman DJ, Wartofsky L. Requirement for mass spectrometry sex steroid assays in the *Journal of Clinical Endocrinology and Metabolism*. *J Clin Endocrinol Metab* 2013;98:3971–3.

## Supplementary files

**Appendix I.** Studbook (SB) number, demographics, sampling date range (Lab A, B or C), and reproductive events for 36 female pygmy hippos (*Choeropsis liberiensis*) from European and North American zoological institutions. Estrus behavior and mating are based on husbandry staff observations and were not reported for some study animals.

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
438	Betty	38	No	–	–	21 May 2014 26 Apr 2015	–	–	–	Suspected post-reproductive
441	Paula	30	Yes <sup>☼</sup>	14 Sep 2006 04 Oct 2007	07 Oct 2007 24 Aug 2009	–	14 Nov 2007 17 Sep 2008 01 Dec 2008 06 Mar 2009	04 Apr 2007 15 Apr 2008 16 Jun 2008 19 Jul 2008 03 Dec 2008 11 Jan 2009	–	
444	Agnes	38	No	–	–	14 Apr 2014 16 Apr 2015	–	–	–	Suspected post-reproductive
454	Adele	37	No	–	–	20 May 2014 21 May 2015	–	–	–	Suspected post-reproductive

Appendix I. Con't

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
548	Tana	27	Yes	06 Apr 2007 13 Jan 2009	–	–	08 Jun 2007 10 Jul 2007 08 Aug 2007	–	–	
626	Lise	23	Yes <sup>☆</sup>	14 Dec 2006 13 Aug 2007	–	–	22 Jan 2007 03 Mar 2007 03 Apr 2007 10 May 2007 11 Jun 2007 21 Jul 2007 24 Aug 2007	12 Dec 2006	26 Nov 2006 <sup>b</sup>	
687	Lalla	21	Yes <sup>△</sup>	02 Oct 2006 15 Sep 2008	–	–	–	2-3 Oct 2006 <sup>a</sup>	20 Jul 2006 <sup>b</sup> 23 Apr 2007	
758	Obesa	26	No	–	–	20 May 2014 21 May 2015	15 Jan 2015 17-19 Feb 2015	–	–	

Appendix I. Con't

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
772	Anais	19	Yes	05 Feb 2007 05 Oct 2007	–	–	–	–	05 Jun 2007	
817	Tootsie	24	Yes <sup>☼</sup>	–	–	14 Apr 2014 16 Apr 2015	–	–	23 Mar 2014	Housed with an adult male from 21 Aug 2014 to 15 Jan 2015
864	Kelsey	22	No	–	–	05 Mar 2014 23 Mar 2015	13 Mar 2014 14 May 2014 07 Jun 2014 03 Jul 2014 26 Aug 2014 18 Mar 2015	–	–	Housed separately from but adjacent to an adult male
904	Nelie	13	No	13 Oct 2006 20 Feb 2009	–	–	–	–	–	
913	Leah	13	Yes <sup>☼</sup>	–	28 Oct 2007 03 May 2009	–	–	–	–	Male arrived 21 May 2008

Appendix I. Con't

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
931	Hope	19	Yes <sup>Δ</sup>	–	–	05 May 2014 04 May 2015	–	–	28 Aug 2014	
938	Wendy	13	Yes <sup>☼</sup>	–	25 Mar 2008 23 May 2009	–	17 Feb 2009 21 Mar 2009	25 Apr 2008 <sup>a</sup>	11 Nov 2008	
941	Sabine	11	Yes <sup>☼</sup>	10 Oct 2006 06 Dec 2007	–	–	–	–	–	
993	Debby	9	Yes <sup>Δ</sup>	05 Sep 2006 05 Oct 2007	08 Oct 2007 02 Apr 2009	–	22 Sep 2006	24 Oct 2006 <sup>a</sup> 22 May 2008 <sup>a</sup>	17 May 2007 18 Dec 2008	
1028	Monica	8	Yes	02 Nov 2006 22 Jan 2009	–	–	15 Apr 2008 09 May 2008 02 Jun 2008 19 Aug 2008 31 Oct 2008 03 Dec 2008	01 Mar 2009 <sup>a</sup>	11 Dec 2006 19 Sep 2009	

Appendix I. Con't

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
1033	Torpedo	8	Yes <sup>Δ</sup>	25 Jul 2006 13 Nov 2009	–	–	25 Sep 2006 30 Jan 2007	17 Aug 2007 <sup>a</sup> 15 Apr 2009 <sup>a</sup>	17 Feb 2006 06 Mar 2008 28 Oct 2009	
1063	Chomel	14	Yes <sup>Δ</sup>	–	–	20 May 2014 18 May 2015	–	22 May 2014 <sup>a</sup> 21-22 Dec 2014 03-05 Feb 2015 07-09 Mar 2015 15 Mar 2015 <sup>a</sup>	07 Dec 2014 <sup>b</sup> 01 Oct 2015	
1068	Venus	7	Yes <sup>☆</sup>	21 Mar 2007 29 Sep 2007	25 Oct 2007 03 Aug 2009	–	27 May 2007 02 Jul 2007 22 Oct 2008 14 Apr 2009	05 May 2009 <sup>a</sup>	27 Nov 2009	Inconsistent sampling

Appendix I. Con't

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
1078	Tanquey	7	Yes	06 Apr 2007 13 Jan 2009	–	–	24 Apr 2007 25 May 2007 22 Jun 2007 27 Jul 2007 24 Aug 2007 11 Sep 2007 25 Oct 2007	30-Nov-07	–	
1082	Bagatelle	5	Yes <sup>☼</sup>	10 Oct 2006 30 Dec 2008	06 Jan 2009 10 Sep 2009	–	15 Mar 2009 28 Apr 2009 23 Aug 2009	28 May 2009	–	
1096	Haley	13	Yes <sup>☼</sup>	–	–	14 Apr 2014 16 Apr 2015	–	03 Nov 2014 <sup>a</sup>	30 May 2015	Housed with an adult male from 10 July 2014 to 04 Nov 2015
1143	Fitri	10	Yes <sup>Δ</sup>	–	–	28 Apr 2014 27 Apr 2015	–	–	12 Oct 2014	Mating and conception approximately 21 Mar 2014

Appendix I. Con't

SB No.	Name	Age* (years)	Housed with a ♂	Lab A	Lab B	Lab C	Estrus behavior	Mating	Births <sup>§</sup>	Notes
1147	Clover	10	Yes <sup>Δ</sup>	–	–	29 May 2014 28 May 2015	–	13-14 Oct 2014	24 Jun 2015	Mating and conception approximately 01 Dec 2014
1148	Krakunia	3	Yes <sup>☼</sup>	–	28 Oct 2007 03 May 2009	–	–	–	–	Male arrived 21 May 2008
1169	Violet	9	Yes <sup>☼</sup>	–	–	14 Apr 2014 16 Apr 2015	–	15-16 Apr 2014 <sup>a</sup>	12 Nov 2014	
1178	Kabibi	9	Yes <sup>☼</sup>	–	–	14 Apr 2014 16 Apr 2015	–	–	20 Aug 2014	Mating and conception approximately 01 Feb 2014
1182	Ashaki	5	Yes <sup>☼</sup>	19 Apr 2011 25 Nov 2013	–	–	–	25 Aug 2013 <sup>a</sup>	18 Mar 2014	
1185	Ellen	2	Yes <sup>Δ</sup>	30 Nov 2006 12 Oct 2007	18 Oct 2007 09 Jul 2009	–	–	20 Nov 2008 <sup>a</sup>	09 Jun 2009	Inconsistent sampling
1194	Tahiti	2	No	30 May 2008 13 Jan 2009	–	–	–	–	–	Juvenile

**Appendix I. Con't**

<b>SB No.</b>	<b>Name</b>	<b>Age* (years)</b>	<b>Housed with a ♂</b>	<b>Lab A</b>	<b>Lab B</b>	<b>Lab C</b>	<b>Estrus behavior</b>	<b>Mating</b>	<b>Births<sup>§</sup></b>	<b>Notes</b>
1201	Sirana	2	No	–	25 Mar 2008 17 Feb 2009	–	–	–	–	Inconsistent sampling, juvenile
1208	Isoke	7	Yes <sup>Δ</sup>	–	–	02 Jun 2014 23 May 2015	–	–	–	
1283	Clementine	5	Yes <sup>Δ</sup>	–	–	25 Jul 2014 06 Aug 2015	31 Oct 2014	–	–	Housed with a juvenile male from 04 Sept 2014
1338	Asali	3	No	–	–	05 Mar 2014 31 Oct 2014	13 Mar 2014 11 May 2014 07 Jun 2014 27 Aug 2014 26-28 Sep 2014	–	–	Housed separately from but adjacent to an adult male

\* Age at the beginning of the study period

§ Reported for 12 months before, during and 12 months after the study period for each hippo

⊛ Only put with a male for breeding or during estrus

<sup>Δ</sup> Housed together with a male when not with a calf

<sup>a</sup> Conceived after mating

<sup>b</sup> Stillbirth or neonatal mortality

**Appendix II.** Cross-reactivities for the eight EIAs used to assess reproductive hormone metabolites in female pygmy hippos.

Cross-Reacting Steroids	% Cross Reactivity							
	Estradiol-17 $\beta$ -OH	Estradiol 17 $\beta$ R4972	Estradiol 17 $\beta$ R0008	5 $\alpha$ -pregnane-3 $\beta$ -ol-20-one	Monoclonal progesterone quidel clone 425	Polyclonal progesterone R4849	5 $\beta$ -pregnane-3 $\alpha$ ,20 $\alpha$ -diol 3HS:BSA	Pregnanediol-3-glucuronide R13904
	<i>E2a</i>	<i>E2b</i>	<i>E2c</i>	<i>20-oxo-pregnane</i>	<i>Mono-P4</i>	<i>Poly-P4</i>	<i>Pg-diol</i>	<i>PdG</i>
Estrone	100.0	3.3	0.73	< 0.10	–	< 0.01	–	–
Estrone sulfate	–	< 0.01	< 0.01	–	–	–	–	–
Estradiol 17 $\alpha$	19.0	–	–	–	–	–	–	–
Estradiol 17 $\beta$	70.0	100.00	100.0	–	–	< 0.01	–	0.04
Estriol	129.0	–	–	–	–	–	–	–
1,3,5(10),7-estratetraen-3-ol-17-one	87.0	–	–	–	–	–	–	–
1,3,5(10),7-estratetraen-3,17 $\beta$ -diol	20.0	–	–	–	–	–	–	–
Progesterone	–	0.80	< 0.01	100.0	100.0	100.0	–	0.20
Pregnenolone	–	–	–	–	–	0.12	–	–
11 $\alpha$ -hydroxy-progesterone	–	–	–	–	–	40.0	–	–
17 $\alpha$ -hydroxy-progesterone	–	–	–	–	–	0.38	–	–
20 $\alpha$ -hydroxy-progesterone	–	–	–	<0.10	–	0.13	150.0	44.8
20 $\beta$ -hydroxy-progesterone	–	–	–	<0.10	–	0.13	2.1	3.2
4-pregnen-3 $\alpha$ -ol-20-one	–	–	–	20.0	188.0	–	–	–
4-pregnen-3 $\beta$ -ol-20-one	–	–	–	68.0	172.0	–	–	–
4-pregnen-11 $\alpha$ -ol-3,20 dione	–	–	–	–	147.0	–	–	–
4-pregnen-11 $\beta$ -ol-3,20 dione	–	–	–	–	2.7	–	–	–
5-pregnen-3 $\beta$ , 20 $\alpha$ -diol	–	–	–	–	–	–	16.6	–

Appendix II. Con't

Cross-Reacting Steroids	% Cross Reactivity							
	Estradiol-17β-OH	Estradiol 17β R4972	Estradiol 17β R0008	5α-pregnane-3β-ol-20-one	Monoclonal progesterone quidel clone 425	Polyclonal progesterone R4849	5β-pregnane-3α,20α-diol 3HS:BSA	Pregnanediol-3-glucuronide R13904
	<i>E2a</i>	<i>E2b</i>	<i>E2c</i>	<i>20-oxo-pregnane</i>	<i>Mono-P4</i>	<i>Poly-P4</i>	<i>Pg-diol</i>	<i>PdG</i>
5α-pregnane-3,20-dione	–	–	–	75.0	55.0	12.2	–	–
5α-pregnane-3α-ol-20-one	–	–	–	8.0	64.0	–	–	–
5α-pregnane-3β-ol-20-one	–	–	–	102.0	94.0	–	–	–
5α-pregnane-20α-ol-3-one	–	–	–	–	–	–	50.0	–
5α-pregnane-3α, 20α-diol	–	–	–	< 0.01	–	–	24.0	–
5α-pregnane-3β, 20α-diol	–	–	–	–	–	–	55.6	–
5α-pregnane-3α, 20β-diol	–	–	–	< 0.01	< 0.01	–	–	–
Pregnanediol	–	–	–	< 0.01	< 0.01	–	100.0	–
Pregnanediol-3-glucuronide	–	–	–	–	–	< 0.01	–	100.0
5β-pregnane-3,20-dione	–	–	–	151.0	8.0	–	–	–
5β-pregnane-3α-ol-20-one	–	–	–	20.0	2.5	–	–	–
5β-pregnane-3β-ol-20-one	–	–	–	36.0	12.5	–	–	–
5β-pregnane-20α-ol-3-one	–	–	–	–	–	–	176.0	–
5β-pregnane-3β, 20α-diol	–	–	–	–	–	–	100.0	–
Androstenedione	–	1.00	–	–	< 0.01	–	–	–
Testosterone	–	1.00	< 0.01	–	–	< 0.01	–	0.20
Cortisol	–	< 0.01	< 0.01	< 0.01	–	< 0.04	–	0.06
Corticosterone	–	< 0.01	< 0.01	–	< 0.01	–	–	–