

Effect of ethnomedicinal extracts used for wound healing on cellular migration and intracellular reactive oxygen species release in SC-1 fibroblasts

Ewura Seidu Yahaya,^{a,b} Werner Cordier,^a Paul Anton Steenkamp^c and Vanessa Steenkamp^a

^aDepartment of Pharmacology, School of Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

^bDepartment of Pharmacology, School of Medical Sciences, University of Cape Coast, Cape Coast, Ghana

^cDepartment of Biochemistry, University of Johannesburg, Auckland Park, South Africa

Correspondence

Vanessa Steenkamp, Department of Pharmacology, Faculty of Health Sciences, School of Medicine, University of Pretoria, Private Bag X323, Arcadia, 0007, Pretoria, South Africa. Email: vanessa.steenkamp@up.ac.za

Tel. +27(0)123192547

Highlights

- UPLC-TOF-MS revealed presence of neobavaisoflavone in *Erythrina senegalensis*.
- Extracts were not cytotoxic to SC-1 fibroblasts.
- *Aspilia africana* and *Erythrina senegalensis* extracts reduced ROS release.
- All extracts showed the tendency to hinder fibroblast migration.

Abstract

The inhibition of reactive oxygen species (ROS) and the migration of fibroblasts are key processes involved in wound healing. In this study, the ability of aqueous ethnomedicinal plant extracts prepared from *Aspilia africana* CD Adams, *Boerhavia diffusa* L. and *Erythrina senegalensis* DC. to mediate fibroblast migration and ROS release was determined. Phytochemical composition was assessed using thin-layer chromatography (TLC), whereas phytochemical markers were detected using ultra-performance liquid chromatography coupled to time of flight mass spectrometry (UPLC-TOF-MS). Sulforhodamine B staining and morphological examination via microscopy was conducted to determine cytotoxic effects on SC-1 fibroblasts. The effect on AAPH-induced oxidative stress was assessed by measuring ROS release using dichlorofluorescein diacetate activation. The scratch wound assay was used to estimate the rate of cellular migration. Alkaloids, flavonoids and phenols were detected in all three extracts using TLC, whilst UPLC-TOF-MS revealed the presence of neobavaisoflavone in *E. senegalensis*. None of the extracts was cytotoxic to the SC-1 cells at the highest in-well concentration tested (100 µg/mL). *E. senegalensis* extract reduced intracellular ROS and cellular migration by 35% and 32.5%, respectively. Although these plant extracts have the potential to minimize oxidation, they do not facilitate fibroblast migration. Further investigation into their mechanism of wound healing is required.

Keywords: Antioxidants, *Aspilia africana*, *Boerhavia diffusa*, *Erythrina senegalensis*, fibroblasts, oxidative stress, wounds

1. Introduction

Healing of wounded skin involves a cascade of complex biological processes which rely on the interaction between several mediators, including platelets, extracellular matrix molecules, fibroblasts, and several other inflammatory cells and products (Martin and Nunan, 2015). Cellular migration is generally pivotal to many physiological and pathological processes, including embryogenesis, angiogenesis, cancer growth and invasion, inflammatory response, and wound healing (Ascione et al., 2016). The proliferation and recruitment of fibroblasts, stimulated by growth factors released by platelets, monocytes and other cellular constituents during the wound healing process, is of particular importance because they are directly responsible for key events in the healing process such as deposition of extracellular matrix components, and collagen (Darby et al., 2014).

The presence of fibroblasts stimulates the production of diverse cytokines related to wound healing, such as interferons (INF)- α , β and γ for macrophage activation, and keratinocyte growth factors for keratinocyte differentiation, proliferation and migration (Broughton et al., 2006). Therefore, substances which affect fibroblast function could affect the healing process. Recently, it has been shown that apart from its antimicrobial effects, silver nanoparticles can also promote wound healing by facilitating fibroblast migration (You et al., 2017). Though much improvement has been made in conventional wound treatment across the world, many patients, particularly in poorly resourced countries have over the years relied on natural products such as medicinal plants for treatment (Pereira and Bartolo, 2016).

Medicinal plants provide a plethora of compounds which could serve as leads for drug discovery. Phytochemicals such as flavonoids and phenolic compounds, amongst a host of others, have been demonstrated to possess useful biological activities such as antioxidant and antimicrobial effects that could account for their use as healing agents (Sasidharan et al., 2011). *Aspilia africana* CD Adams (haemorrhage plant, Asteraceae), *Boerhavia diffusa* L. (spreading hogweed, Nyctaginaceae) and *Erythrina senegalensis* DC (coral flower, Papilionaceae) are examples of commonly used medicinal plants in tropical countries for management of several disease states, amongst others, wounds (Okoli et al., 2007; Kuete et al., 2014; Patil and Bhalsing,

2016). All three plants have been shown to possess a diverse array of phytochemicals with multiple pharmacological properties to attest for their ethnomedicinal use (Okoli et al., 2007; Kuete et al., 2014; Patil and Bhalsing, 2016). In a previous study, the wound healing potential of fractions sequentially extracted from *B. diffusa* was demonstrated (Yahaya et al., 2018). The current study determined the ability of aqueous ethnomedicinal plant extracts prepared from *Aspilia africana*, *Boerhavia diffusa* and *Erythrina senegalensis* to effect fibroblast migration and ROS release.

2. Methods

2.1. Plant material

Plant specimens were collected from the University of Cape Coast (UCC) botanical gardens in the Central Region of Ghana, and authenticated by a botanist at the University's School of Biological Sciences herbarium. Voucher specimens for *A. africana* (UCCH 0211215), *B. diffusa* (UCCH 0041215) and *E. senegalensis* (UCCH 0091215) are deposited at the aforementioned herbarium. Plant material was washed, shade-dried at room temperature, and powdered using an electronic grinding device. Powdered samples were stored in air-tight containers until extraction.

2.2. Extract preparation

Powdered plant material (10 g) was extracted by immersion in distilled water (100 mL), agitating on an electronic shaking device for 30 min, and incubating for 24 h at 4 °C. Samples were first filtered through a Whatman Grade-1 filter paper, and subsequently a 0.22 µm Millipore filter paper under vacuum-pressure. The resultant filtrates were freeze-dried (Labconco 31 Freezone 6), and stored at -20 °C until use.

2.3. Chemicals

Dulbecco's modified Eagle's medium (DMEM), foetal calf serum (FCS), penicillin/streptomycin, saponin, 2,2'-azobis(2-amidinopropane) dihydrochloride (AAPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS), 2',7'-dichlorofluorescein diacetate (H2-DCF-DA), sulforhodamine B (SRB), trichloroacetic acid (TCA), trolox, platelet-derived growth factor (PDGF), and potassium persulfate were purchased from Sigma-Aldrich (South Africa). Phosphate buffered saline (PBS) was acquired from Becton Dickinson (South Africa), whilst tris

(hydroxymethyl) aminomethane, and acetic acid were purchased from Merck (South Africa) and Saarchem (South Africa), respectively.

2.4. Phytochemical analysis

The presence of major phytochemical classes in the extracts was determined using thin layer chromatography (Stahl, 1962). Ultra-performance liquid chromatography, coupled to time of flight mass spectrometry (UPLC- TOF- MS), was employed to verify marker compounds in the extracts (Yahaya et al., 2018). UPLC-TOF-MS was conducted on a Waters instrument coupled in tandem to a Waters photodiode array (PDA) detector. A SYNAPT G1 HDMS mass spectrometer was used to generate accurate mass data. MassLynx 4.1 (SCN 872) software was used to control the hyphenated system as well as for data manipulation.

2.5. Cell culture and maintenance

The SC-1 mouse fibroblast cell line was purchased from the American Type Culture Collection (ATTC, CRL-1404), and cultured in DMEM supplemented with 10% FCS, and 1% antibiotic solution (penicillin-streptomycin). Cells were maintained at 37 °C in a humidified atmosphere of 5% CO₂.

2.6. Determination of the effect on cell density

The effect of extracts on cell density was determined using the SRB assay as described previously (Vichai and Kirtikara, 2006). Briefly, cells were seeded into 96-well plates at a density of 1×10^4 cells/well in 10% FCS-supplemented DMEM and incubated for 24 h at the conditions stated above to allow for adherence. Cells were then treated with extracts at final in-well concentrations of 1, 3.2, 10, 32 and 100 µg/mL (prepared in FCS-free DMEM), and incubated for 24 or 48 h. Saponin (1%) and distilled water (0.05%) were used as positive and vehicle controls, respectively. Cells were fixed by adding 50 µL of 50% cold TCA, and incubated at 4 °C overnight. Fixed cells were gently washed with water, dried in a low temperature oven, and stained with 0.057% SRB (in 1% acetic acid) solution for 30 min at room temperature. The plate was then washed three times with 200 µL of 1% acetic acid solution to remove excess dye, and dried. The bound dye was dissociated in 200 µL of a 10 mM Tris-base solution (pH 10), and the absorbance measured at 540 nm using a BioTek ELx800 microplate reader. Cell density was determined using the following equation:

$$\text{Cell density (\% of the negative control)} = \frac{A_s}{A_c} \times 100\%$$

Where A_s and A_c are the blank-corrected absorbances of sample and average negative control respectively.

2.7. Morphological studies

Cell morphology was examined using phase contrast and plasDIC microscopy. Treatments for the morphological studies were similar to that described in Section 2.6., except that cells (2.5×10^4 cells/well) were exposed to the highest concentration of extract tested (100 $\mu\text{g/mL}$) in 24-well plates. Following 24 and 48 h incubation at the conditions stated above, cells were examined microscopically and pictures taken at 10 \times (phase contrast) and 40 \times (plasDIC) magnification.

2.8. Assessment of the effect on oxidative stress

2.8.1. Antioxidant activity determination

The 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical scavenging effect of extracts was estimated using a previously described method (Re et al., 1999). The ABTS⁺ stock solution, containing 7 mM ABTS salt and 2.4 mM potassium persulfate, was prepared in distilled water and incubated in the dark for 16 h at 4 °C. The resultant ABTS⁺ solution was diluted with distilled water to an absorbance of 0.70 ± 0.02 at 734 nm (PerkinElmer Lambda 25 UV/VIS spectrometer). An aliquot of 180 μL ABTS⁺ solution was mixed with 20 μL of varying concentrations of the extracts (20 – 100 $\mu\text{g/mL}$ in-reaction), 0.6 – 5.0 $\mu\text{g/mL}$ Trolox (in-reaction, antioxidant control) or methanol (negative control) in a 96-well plate. The absorbance was read after 30 min incubation in the dark using the Synergy 2 microplate reader (BioTek Instruments, Inc.) at 734 nm. The ABTS⁺ scavenging capacity was calculated as follows:

$$\text{ABTS radical scavenging activity (\% relative to negative control)} = \frac{A_c - A_s}{A_s} \times 100$$

Where A_c and A_s represent the absorbance of the average negative control and sample, respectively.

2.8.2. Effect on AAPH-induced oxidative stress

The ability of extracts to alter intracellular ROS was assessed in an AAPH-induced oxidative stress model as described previously (Lopachev et al., 2016) with minor modifications. Cells were seeded into a 96-well plate at a density of 1×10^4 cells/well and allowed to attach overnight. The culture medium was replaced with 100 μ L fresh medium containing 10 μ M H₂-DCFH-DA and incubated for 30 min in the dark. Excess H₂-DCF-DA was removed by washing twice with 100 μ L PBS, followed by exposure to 1, 10 and 100 μ g/mL extract (in 100 μ L of 10% FCS-supplemented DMEM) for 4 h. All control wells were treated with DMEM during this period, except antioxidant controls to which 5 μ g/mL Trolox was added. Cells were washed twice with 100 μ L PBS and all wells re-suspended in 50 μ L PBS containing 100 μ M AAPH, with the exception of the negative controls to which 50 μ L PBS was added. The relative fluorescence intensity (RFI) was measured every min for 2 h at an excitation and emission wavelength of 485 nm and 530 nm respectively using a Synergy 2 microplate reader (BioTek Instruments, Inc.). Intracellular ROS was estimated as follows:

$$\text{Intracellular ROS (fold - change relative to control)} = \frac{RFIs}{RFIc}$$

Where *RFIs* and *RFIc* represents the RFI of each sample and the average negative control, respectively.

2.9. The effect on cell migration

Effect on fibroblast cell migration was assessed *in vitro* using the scratch wound assay (Liang et al., 2007). Cells were seeded into 24-well plates as described in Section 2.7. in 10% FCS-supplemented DMEM, and incubated for 24 h to allow for cellular attachment. Attached cells were gently washed with PBS, the medium replaced with 0.5% FCS-supplemented DMEM, and incubated for a further 24 h to form a monolayer. An intersecting vertical and horizontal scratch was generated in the cell monolayer of each well, with the latter serving as a reference point for image acquisition. Cellular debris was removed by washing with PBS. Cells were then exposed to extracts at concentrations of 10 and 100 μ g/mL (in-reaction) in 0.5% FCS-supplemented DMEM. PDGF (2 ng/mL in-reaction) served as positive control. The scratches were

microscopically examined and pictures taken at 0, 8, and 24 h after wound generation. The average wound area was estimated using the ImageJ software, and cell migration calculated as:

$$\text{Cell migration (\%)} = \frac{A_0 - A_t}{A_0} \times 100\%$$

Where A_0 and A_t represent average wound area at 0 h and 24 h respectively.

2.10. Statistical analysis

Data represents results of at least three independent experiments conducted in technical triplicates. Statistical analysis was performed using GraphPad Prism 5.00 data analysis software. Data was expressed as the mean \pm standard error of mean (SEM). Non-linear regression was used to determine half-maximal inhibitory concentrations (IC50). The difference between groups was determined by a Kruskal-Wallis test followed by Dunn's post-hoc test (cytotoxicity and ROS), as well as two-way ANOVA followed by Bonferroni post-hoc tests (scratch assay). P values less than 0.05 were considered significant.

3. Results and discussion

3.1. Phytochemical evaluation

Phytochemical composition of the extracts, using TLC, is represented in Table 1. All the extracts contained major phytochemical groups such as alkaloids, flavonoids and phenols. These findings are also reported in previous literature (Saidu et al., 2000; Okwu and Josiah, 2006; Sreeja and Sreeja, 2009).

Table 1. Presence of Phytochemical classes in the plants investigated in this study as determined by thin layer chromatography.

Extract	Phytochemical classes						
	Alkaloids	Flavonoids	Glycosides	Phenols	Saponins	Tannins	Terpenes
<i>Aspilia africana</i>	+	+	-	+	-	-	+
<i>Boerhavia diffusa</i>	+	+	+	+	-	-	-
<i>Erythrina senegalensis</i>	+	+	+	+	-	-	+

+ = presence of component; - = absence of component

Neobavaisoflavone was detected in the *E. senegalensis* extract using UPLC-TOF-MS (Fig. 1). The presence of three cytotoxic isoflavonoids, including neobavaisoflavone, has been reported in the plant (Kueté et al., 2014). Ascorbic acid and kaempferol were not detected, although they had previously been reported in methanol extracts of *A. africana*, and *B. diffusa*, respectively (Ferrerés et al., 2005; Okwu and Josiah, 2006). The lack of detection of these compounds could be ascribed to differing extraction solvents employed or geographical variation of harvested plants.




Fig. 1. UPLC-TOF-MS spectra of neobavaisoflavone (A) and *Erythrina senegalensis* ethnomedicinal extract (B). The arrow indicates the presence of neobavaisoflavone.

3.2. Cytotoxicity

Cellular density of fibroblasts treated with the extracts at varying concentrations (up to 100 $\mu\text{g}/\text{mL}$) resulted in a maximum of 21% death after 24 h and 48 h exposure (Fig. 2). The *E. senegalensis* extract was the most cytotoxic to the SC-1 fibroblasts, with a reduction in cellular density of 9% and 21% after 24 h (Fig. 2A) and 48 h (Fig. 2B) exposure, respectively. Furthermore, evaluation using phase contrast (Fig. 3A) and plasDIC microscopy (Fig. 3B) indicated normal morphology after treatment with the plant extracts, in contrast to the positive control, saponin, where cytotoxicity occurred.




Fig. 2. Effect of 100 µg/mL of the ethnomedicinal extracts on fibroblast cell density after (A) 24 h and (B) 48 h exposure. NC: Negative control




Fig. 3. The effect of ethnomedicinal extracts on SC-1 fibroblast morphology. Images were captured by (A) phase contrast at 10x magnification and (B) plasDIC at 40x magnification. AA: *A. africana*; BD: *B. diffusa*; ES: *E. senegalensis*; NC: Negative control; PC: positive control. Arrows indicate damaged cells. Scale bar: 100 µm.

Negligible cytotoxicity ($IC_{50} > 300 \mu\text{g/mL}$) has been reported for the hydroalcoholic extract of *B. diffusa* in drug-sensitive human T-lymphoblastoid (CCRF-CEM) and multidrug-resistant P-glycoprotein over-expressing leukaemia (CEM/ADR5000) cell lines (Tacchini et al., 2015). The isoflavones; neobavaisoflavone, sigmoidin H and isoneorautenol, have been reported to be cytotoxic (Kuate et al., 2014). These compounds could be responsible for the higher cytotoxic activity noted in this study for the *E. senegalensis* extract.

3.3. Effect on oxidation

The extract with the highest ABTS⁺ radical scavenging activity was *A. africana* (IC₅₀ = 36.18 µg/mL), compared to an IC₅₀ of 51.52 µg/mL for *E. senegalensis* and more than a 100 µg/mL for *B. diffusa*. The positive control, Trolox, had an IC₅₀ of 2.92 µg/mL. Pre-treatment of cells with 100 µg/mL *E. senegalensis* and *A. africana* extracts prior to induction of oxidative stress led to a 35% and 25% reduction in ROS release, respectively (Fig. 4). *E. senegalensis* (1 µg/mL) was better at suppressing ROS, even though *A. africana* had a higher ABTS⁺ scavenging activity. This suggests a probable diminished uptake of *A. africana* into the cells, or that a different mechanism of quenching is at play. Although the response was not dose-dependent, differences between the various concentrations tested was not statistically significant. The ethnomedicinal extract of *B. diffusa* on the other hand had no effect on the amount of ROS released, which confirms results obtained in the ABTS⁺ scavenging assay.




Fig. 4. The effect of ethnomedicinal extracts on ROS release in SC-1 fibroblasts. AA: *A. africana*; BD: *B. diffusa*; ES: *E. senegalensis*; NC: negative control; AAPH: oxidant control; Trolox: antioxidant control.

Reactive oxygen species are crucial for the maintenance of normal cellular physiology in that they regulate signalling pathways which control cell growth, metabolic rate, cell division, apoptosis, necrosis and aging (Menon and Goswami, 2007). Therefore, an optimal amount of ROS results in timeous healing of wounds. Uncontrolled ROS release or depletion of intracellular antioxidants results in oxidative stress, which delays healing by damaging cells, tissues and other components in the body (Menon

and Goswami, 2007). The ability of the extracts to suppress oxidative stress could be useful in managing diseases where oxidative stress is involved such as in delayed wound healing. Phytochemical components, such as the flavonol kaempferol, have been shown to inhibit ROS (Wang et al., 2006). Further studies are required to identify the compounds responsible for this activity.

3.4. Fibroblast migration

Cellular migration is an important process in wound repair, allowing fibroblasts to traverse local tissue environments in order to degrade, repair and remodel the extracellular matrix (Tschumperlin, 2013). Twenty four hours after artificial wound generation, none of the extracts were found to have facilitated wound closure when compared to the negative control (Figs. 5 and 6). Wound closure in the negative control cells was 64.4%, compared to 80% in the positive control (PDGF). Treatment with all three ethnomedicinal extracts decreased fibroblast migration relative to the negative control, particularly at high concentrations (100 $\mu\text{g/mL}$). Though 10 $\mu\text{g/mL}$ *A. africana* extract had no effect on cellular migration, 100 $\mu\text{g/mL}$ hindered migration by 5.2%. Treatment with *B. diffusa* extract on the other hand produced a 7.6% and 21.8% suppression of cell migration at 10 $\mu\text{g/mL}$ and 100 $\mu\text{g/mL}$, respectively compared to the negative control. The *E. senegalensis* extract had the greatest effect on fibroblast migration, with a 16.2% and 32.5% reduction in migration at 10 $\mu\text{g/mL}$ and 100 $\mu\text{g/mL}$, relative to the negative control respectively.




Fig. 5. The effect of ethnomedicinal extracts on SC-1 fibroblast migration over a 24 h period assessed using the scratch wound assay. AA: *A. africana*; BD: *B. diffusa*; ES: *E. senegalensis*; NC: negative control; PDGF: platelet-derived growth factor. Dotted line depicts maximum wound closure in NC.




Fig. 6. Micrographs of fibroblast migration after treatment with 10 and 100 µg/mL extracts. AA: *A. africana*; BD: *B. diffusa*; ES: *E. senegalensis*; NC: negative control; PDGF: platelet-derived growth factor. Scale bar = 200 µm.

The observed effect could be linked to the ROS inhibitory ability of the extracts. Whilst extracts with a higher ability to suppress oxidation such as *E. senegalensis* delayed migration to the greatest extent, those with minimal effect on oxidation recorded a much lower inhibitory effect. Cellular migration is regulated by numerous intracellular factors. Inhibition of either the ERK1/2 or p38 pathways resulted in delayed corneal epithelial wound healing (Sharma et al., 2003). Furthermore, activation of the P13K/Akt/mTOR pathway has resulted in accelerated wound closure (Squarize et al., 2010). More recently, oxidation of Akt2 kinase has been shown to facilitate cellular migration (Wani et al., 2011). These suggest a probable relationship between ROS and cellular migration. Therefore, the extracts could be inhibiting migration by indirectly inhibiting the P13K/Akt/mTOR pathway through ROS inhibition.

4. Conclusions

Results obtained in this study have shown that the ethnomedicinal extracts prepared from the three wound healing plant extracts are not cytotoxic in SC-1 fibroblasts at the

concentrations tested. Though the *E. senegalensis* extract demonstrated ability to suppress AAPH-induced oxidation, the other two extracts had no effect on ROS release. This could be due to *E. senegalensis*' better antioxidant potential, suggesting that it could be useful in the management of ROS-induced chronic wounds. All the extracts however showed tendency to inhibit fibroblast migration, particularly at higher doses. Future studies should be focused at identifying the compounds responsible for these activities, as well as their molecular mechanisms of action.

Acknowledgements

The authors are grateful to Ms Margo Nell for her technical skills during culturing of cells, and the Department of Physiology, University of Pretoria, for the use of their microscope. This work was supported by the National Research Foundation of South Africa (NRF-TWAS grant number 100000, 2016).

References

- Ascione, F., Vasaturo, A., Caserta, S., D'Esposito, V., Formisano, P., Guido, S., 2016. Comparison between fibroblast wound healing and cell random migration assays *in vitro*. *Experimental Cell Research* 347, 123-132.
- Broughton, G., Janis, J.E., Attinger, C.E., 2006. Wound healing: an overview. *Plastic and Reconstructive Surgery* 117, 1e-S-32e-S.
- Darby, I.A., Laverdet, B., Bonté, F., Desmoulière, A., 2014. Fibroblasts and myofibroblasts in wound healing. *Clinical, Cosmetic and Investigational Dermatology* 7, 301-311.
- Ferreres, F., Sousa, C., Justin, M., Valentao, P., Andrade, P.B., Llorach, R., Rodrigues, A., Seabra, R.M., Leitao, A., 2005. Characterisation of the phenolic profile of *Boerhaavia diffusa* L. by HPLC-PAD-MS/MS as a tool for quality control. *Phytochemical Analysis* 16, 451-458.
- Kuete, V., Sandjo, L.P., Kwamou, G.M., Wiench, B., Nkengfack, A.E., Efferth, T., 2014. Activity of three cytotoxic isoflavonoids from *Erythrina excelsa* and *Erythrina senegalensis* (neobavaisoflavone, sigmoidin H and isoneorautenol) toward multi-factorial drug resistant cancer cells. *Phytomedicine* 21, 682-688.

- Liang, C.-C., Park, A.Y., Guan, J.-L., 2007. *In vitro* scratch assay: a convenient and inexpensive method for analysis of cell migration *in vitro*. *Nature Protocols* 2, 329-333.
- Lopachev, A.V., Lopacheva, O.M., Abaimov, D.A., Koroleva, O.V., Vladychenskaya, E.A., Erukhimovich, A.A., Fedorova, T.N., 2016. Neuroprotective effect of carnosine on primary culture of rat cerebellar cells under oxidative stress. *Biochemistry* 81, 511-520.
- Martin, P., Nunan, R., 2015. Cellular and molecular mechanisms of repair in acute and chronic wound healing. *British Journal of Dermatology* 173, 370-378.
- Menon, S.G., Goswami, P.C., 2007. A redox cycle within the cell cycle: ring in the old with the new. *Oncogene* 26, 1101-1109.
- Okoli, C.O., Akah, P.A., Okoli, A.S., 2007. Potentials of leaves of *Aspilia africana* (Compositae) in wound care: an experimental evaluation. *BMC Complementary and Alternative Medicine* 7, 24.
- Okwu, D.E., Josiah, C., 2006. Evaluation of the chemical composition of two Nigerian medicinal plants. *African Journal of Biotechnology* 5, 357-361.
- Patil, K.S., Bhalsing, S.R., 2016. Ethnomedicinal uses, phytochemistry and pharmacological properties of the genus *Boerhavia*. *Journal of Ethnopharmacology* 182, 200-220.
- Pereira, R.F., Bartolo, P.J., 2016. Traditional therapies for skin wound healing. *Advances in Wound Care* 5, 208-229.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C., 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 26, 1231-1237.
- Saidu, K., Onah, J., Orisadipe, A., Olusola, A., Wambebe, C., Gamaniel, K., 2000. Antiplasmodial, analgesic, and anti-inflammatory activities of the aqueous extract of the stem bark of *Erythrina senegalensis*. *Journal of Ethnopharmacology* 71, 275-280.
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K.M., Latha, L.Y., 2011. Extraction, isolation and characterization of bioactive compounds from plants' extracts. *African Journal of Traditional Complementary and Alternative Medicines* 8, 1-10.
- Sharma, G.-D., He, J., Bazan, H.E., 2003. p38 and ERK1/2 coordinate cellular migration and proliferation in epithelial wound healing evidence of cross-talk

- activation between map kinase cascades. *Journal of Biological Chemistry* 278, 21989-21997.
- Squarize, C.H., Castilho, R.M., Bugge, T.H., Gutkind, J.S., 2010. Accelerated wound healing by mTOR activation in genetically defined mouse models. *PLoS One* 5, e10643.
- Sreeja, S., Sreeja, S., 2009. An *in vitro* study on antiproliferative and antiestrogenic effects of *Boerhaavia diffusa* L. extracts. *Journal of Ethnopharmacology* 126, 221-225.
- Stahl, E. 1962. Thin-layer chromatography. A laboratory handbook. Springer-Verlag, Berlin.
- Tacchini, M., Spagnoletti, A., Marieschi, M., Caligiani, A., Bruni, R., Efferth, T., Sacchetti, G., Guerrini, A., 2015. Phytochemical profile and bioactivity of traditional ayurvedic decoctions and hydro-alcoholic macerations of *Boerhaavia diffusa* L. and *Curculigo orchioides* Gaertn. *Natural Product Research* 29, 2071-2079.
- Tschumperlin, D.J., 2013. Fibroblasts and the ground they walk on. *Physiology* 28, 380-390.
- Vichai, V., Kirtikara, K., 2006. Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nature Protocols* 1, 1112-1116.
- Wang, L., Tu, Y.-C., Lian, T.-W., Hung, J.-T., Yen, J.-H., Wu, M.-J., 2006. Distinctive antioxidant and antiinflammatory effects of flavonols. *Journal of Agricultural and Food Chemistry* 54, 9798-9804.
- Wani, R., Bharathi, N.S., Field, J., Tsang, A.W., Furdui, C.M., 2011. Oxidation of Akt2 kinase promotes cell migration and regulates G1-S transition in the cell cycle. *Cell Cycle* 10, 3263-3268.
- Yahaya, E.S., Cordier, W., Steenkamp, P.A., Steenkamp, V., 2018. Attenuation of oxidative stress and artificial wound closure in C2C12 myoblasts induced by sequential extracts of *Boerhavia diffusa*. *Journal of Pharmacy and Pharmacology* 70, 111-123.
- You, C., Li, Q., Wang, X., Wu, P., Ho, J.K., Jin, R., Zhang, L., Shao, H., Han, C., 2017. Silver nanoparticle loaded collagen/chitosan scaffolds promote wound healing via regulating fibroblast migration and macrophage activation. *Scientific Reports* 7, 10489.