

Supplementary Materials for

Moving in the Anthropocene: Global Reductions in Terrestrial Mammalian Movements

Marlee A. Tucker, Katrin Böhning-Gaese, William F. Fagan, John M. Fryxell, Bram Van Moorter, Susan C. Alberts, Abdullahi H. Ali, Andrew M. Allen, Nina Attias, Tal Avgar, Hattie Bartlam-Brooks, Bayarbaatar Buuveibaatar, Jerrold L. Belant, Alessandra Bertassoni, Dean Beyer, Laura Bidner, Floris M. van Beest, Stephen Blake, Niels Blaum, Chloe Bracis, Danielle Brown, PJ Nico de Bruyn, Francesca Cagnacci, Justin M. Calabrese, Constança Camilo-Alves, Simon Chamailié-Jammes, Andre Chiaradia, Sarah C. Davidson, Todd Dennis, Stephen DeStefano, Duane Diefenbach, Iain Douglas-Hamilton, Julian Fennessy, Claudia Fichtel, Wolfgang Fiedler, Christina Fischer, Ilya Fischhoff, Christen H. Fleming, Adam T. Ford, Susanne A. Fritz, Benedikt Gehr, Jacob R. Goheen, Eliezer Gurarie, Mark Hebblewhite, Marco Heurich, A. J. Mark Hewison, Christian Hof, Edward Hurme, Lynne A. Isbell, René Janssen, Florian Jeltsch, Petra Kaczensky, Adam Kane, Peter Kappeler, Matthew Kauffman, Roland Kays, Duncan Kimuyu, Flavia Koch, Bart Kranstauber, Scott LaPoint, Peter Leimgruber, John D. C. Linnell, Pascual López-López, A. Catherine Markham, Jenny Mattisson, Emilia Patricia Medici, Ugo Mellone, Evelyn Merrill, Guilherme de Miranda Mourão, Ronaldo G. Morato, Nicolas Morellet, Thomas Morrison, Samuel L Díaz-Muñoz, Atle Mysterud, Dejid Nandintsetseg, Ran Nathan, Aidin Niamir, John Odden, Robert B. O'Hara, Luiz Gustavo R. Oliveira-Santos, Kirk A. Olson, Bruce D. Patterson, Rogerio Cunha de Paula, Luca Pedrotti, Björn Reineking, Martin Rimmler, Tracey L. Rogers, Christer Moe Rolandsen, Christopher S. Rosenberry, Daniel I. Rubenstein, Kamran Safi, Sonia Saïd, Nir Sapir, Hall Sawyer, Niels Martin Schmidt, Nuria Selva, Agnieszka Sergiel, Enkhtuvshin Shiilegdamba, João Paulo Silva, Navinder Singh, Erling J. Solberg, Orr Spiegel, Olav Strand, Siva Sundaresan, Wiebke Ullmann, Ulrich Voigt, Jake Wall, David Wattles, Martin Wikelski, Christopher C. Wilmers, John W. Wilson, George Wittemyer, Filip Zięba, Tomasz Zwijacz-Kozica, Thomas Mueller
Correspondence Author. Email: tucker.marlee@gmail.com;
thomas.mueller@senckenberg.de

This PDF file includes:

Materials and Methods
Supplementary Text
Figs. S1 to S2
Tables S1 to S5
References

Materials and Methods

Displacement Data

We compiled GPS location data for 57 mammalian species, comprising 7 339 376 locations of 803 individuals from 1998 to 2015 (Fig. 1, Supplementary Table S1). The dataset included adult male and female individuals. Datasets were obtained from the online animal tracking database *Movebank* (<https://www.movebank.org/>), the Movebank Data Repository (*Equus quagga* (1, 2) and *Loxodonta africana* (3, 4)), or were contributed by co-authors directly (Table S2). For species that are inactive at night (e.g., primates sleeping overnight in trees) and where the GPS devices had been switched off to prolong battery life, we interpolated location data during the inactive phase (i.e., using the last recorded position) with the same sampling frequency as that employed for active periods to ensure an even sampling regime.

We sub-sampled the location data with inter-location intervals at a geometric time scale from one hour to ~ ten days (i.e. 1, 2, 4, 8, 16, 32, 64, 128 and 256 hours) using the “SyncMove” R package (5). We started the sub-sampling algorithm from the first location recorded for each individual. For each of the nine time scales, we calculated the geodesic distance between the subsampled locations using the Spherical Law of Cosines using 6371 km as the mean radius of the Earth (6). This allowed a systematic investigation across time scales from within day movements to more long-term movements, and standardized the sampling regime across studies and individuals. Smaller time intervals were not available for most species and longer time intervals resulted in a significant loss in sample size. Sub-sampling precision was set to the inter-location interval $\pm 4\%$ (e.g., for the 1-hour scale resulting in inter-location intervals varying between 57 and 62 minutes). We then checked the data for outliers, specifically for maximum movement speeds that were unlikely for a terrestrial land mammal to achieve over a given time period ($> 4 \text{ m s}^{-1}$), and removed them (7). We calculated two response variables for each individual: the 0.5 quantile displacement distance and the 0.95 quantile displacement distance, the former describing the median movement behavior of that individual, and the latter describing long-distance movements (Supplementary Figure S1). All values were \log_{10} transformed prior to analyses.

Covariates

We annotated each GPS location with NDVI and human footprint index (8) (HFI; Supplementary Table S2). NDVI data was extracted from MODIS Land Terra Vegetation Indices 500-m 16-day resolution (MOD13A1 V005 (9)) using the Movebank Env-DATA system (10) (environmental-data automated track annotation; <http://www.movebank.org>). We filtered the NDVI data to remove pixels with no data (-1), snow/ice (2) and clouds (3). We also included species body mass using the PanTHERIA database (11) (where individual mass information was unknown) and diet (i.e., carnivore, herbivore or omnivore) (Table S1). Body mass values were \log_{10} transformed and the NDVI values were scaled. We then calculated the mean NDVI and human footprint value for each

inter-location interval (i.e., the average value between each sequential pair of locations) and averaged these values for each individual.

Analyses

Our final database (Supplementary Fig. 1) comprised nine median and nine 0.95 quantile movement distance values for each individual (one for each temporal scale), associated with nine mean values for body mass, NDVI, and the human footprint index. We only included individuals that had tracking data for a minimum of two months (~60 days) or 50 displacements. We ran 18 linear mixed effects models, two for each time-scale, one with the 0.5 and the other with the 0.95 quantile displacement distances as the dependent variable, and body mass, NDVI, HFI, and diet as the predictor variables. We included species identity as a nested random effect to account for taxonomy (i.e., Order/Family/Genus/Species), and a Gaussian spatial autocorrelation structure (12) including the mean longitude and latitude for each individual. For each model, we checked the residuals for normality (i.e., Q-Q plots) and removed outliers (< 2% of total data points). All correlation coefficients among the predictor variables were $|r| \leq 0.55$ and all variance inflation factors (VIFs) were ≤ 2 , well below the common cut-off values of 0.7 and 4, respectively (13, 14). All model predictions and associated standard errors were calculated using the “AICcmodavg” R package (15). All analyses were performed in R version 3.2.2 (16).

Supplementary Text

Extended List of Acknowledgements

The authors are grateful for support from the Robert Bosch Foundation, Goethe International Postdoctoral Programme, People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement no [291776], German Research Foundation (DFG, FR 3246/2-1), US National Science Foundation (NSF) ABI-1458748, NSF grant #0963022, NSF grant #1255913, Irish Research Council GOIPD/2015/81, NASA funded project: "Animals on the Move", grant NNX15AV92, Research Council of Norway (Grant number 251112), GLOBE POL-NOR/198352/85/2013, UC Berkeley Museum of Vertebrate Zoology, American Society of Mammalogists, NSF DEB-LTREB Grant # 1556248, NSF DDIG grant 0608467, NASA Earth Science Division, Ecological Forecasting Program project number NNX11AP61G, NSF Biological Infrastructure Award #1564380, German Research Foundation (DFG) AOBJ 576687, ANR FEAR, ANR SAVARID, Leverhulme Study Abroad Studentship and ERC (323401), Copenhagen Zoo, the Danish Environmental Protection Agency, 15. Juni Charity Foundation, DFG Research Training Group 2118/1 BioMove, Grant SFRH/BPD/111084/2015 from Fundação para a Ciência e Tecnologia, the “MOVEIT” ANR grant ANR-16 -CE02-0010-02, Save the Elephants, Spanish Ministry of Economy and Competitiveness (grant number IJCI-2014-19190), NSF grants (BCS 99-03949 and BCS 1266389), the L.S.B. Leakey Foundation, and the University of California, Davis, Committee on Research. Movebank is hosted by the Max Planck

Institute for Ornithology and the Movebank Data Repository is hosted by the University of Konstanz. Roe and red deer data were obtained from euroungulates, www.euroungulates.org. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. Figure 1 silhouettes by J. A. Venter, H. H. T. Prins, D. A. Balfour & R. Slotow (vectorized by T. M. Keesey) (hare and buffalo) and R. Groom (gazelle) were downloaded from www.phylopic.org and are available for re-use under the Creative Commons Attribution 3.0 Unported license. Figure 1 silhouettes by S. Traver (boar, deer, tapir, wildcat, elephant, muskox, wolverine, giraffe and khulan), O. Jones (baboon), D. Orr (coyote), T. Heath (bear and wolf) and G. Prideaux (possum) were downloaded from www.phylopic.org and are available for re-use under the Public Domain Mark 1.0 license. Puma, maned wolf and lynx silhouettes by M. Tucker.




Fig. S1.

Distributions of the median and 0.95 quantiles of the individual displacements used in the analyses. The y-axis represents the density distribution of median (0.5 quantile) and long-distance (0.95 quantile) displacements of each individual.




Fig. S2

Model coefficients (\pm CI) predicting mammalian displacements including (A) an individual-behavioral effect and (B) a species-occurrence effect of the Human footprint index (HFI). The individual-behavioral HFI was calculated as the individual HFI minus the species mean HFI, and the species-occurrence HFI was calculated as the species mean HFI. Other covariates of the model included (C) Normalized Difference Vegetation Index (NDVI), (D) body mass, and dietary guild (not shown). The models also included a nested random effect accounting for taxonomy, and a Gaussian spatial autocorrelation structure. Models were run for the median (i.e. -0.5 quantiles; blue) and long-distance (i.e. 0.95 quantiles; red) displacements of each individual calculated across different time scales. When the error bars cross the horizontal line (at 0) the effect is not significant. See Methods and Supplementary Tables S4 for additional details.

Table S1.

Data annotation summary

Variable	Unit	Temporal Resolution	Spatial Resolution	Source	Transformation
Normalised Difference Vegetation Index (NDVI)	Unitless	16 days	500 m	MODIS Land Terra Vegetation Indices 500-m 16-day (MOD13A1 V005)	Scaled
Human Footprint	Unitless	1993-2009 mean	1 km	Global terrestrial Human Footprint maps for 1993 and 2009 (8, 17)	Log ₁₀
Body Mass	Grams	Not applicable.	Not applicable.	K. E. Jones <i>et al.</i> , PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. <i>Ecology</i> . 90 , 2648 (2009).	Log ₁₀
Diet	Unitless, categorical	Not applicable.	Not applicable.	K. E. Jones <i>et al.</i> , PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. <i>Ecology</i> . 90 , 2648 (2009).	Not applicable.

Table S2.

Summary of species and number of individuals per species included in the analyses.

Species	No. Individuals	Data Source	Species	No. Individuals	Data Source
<i>Aepyceros melampus</i>	20	Co-author	<i>Madoqua guentheri</i>	15	Co-author
<i>Alces alces</i>	46	Co-author	<i>Martes pennanti</i>	13	Movebank
<i>Antilocapra americana</i>	25	Co-author	<i>Myrmecophaga tridactyla</i>	4	Co-author
<i>Beatragus hunteri</i>	4	Co-author	<i>Odocoileus hemionus</i>	25	Co-author
<i>Canis aureus</i>	1	Movebank	<i>Odocoileus hemionus columbianus</i>	14	Co-author
<i>Canis latrans</i>	19	Movebank	<i>Odocoileus virginianus</i>	30	Movebank
<i>Canis lupus</i>	12	Co-author & Movebank	<i>Ovibos moschatus</i>	14	Co-author
<i>Capreolus capreolus</i>	94	Eurodeer & co-author	<i>Panthera leo</i>	2	Movebank
<i>Cercocebus galeritus*</i>	1	Co-author	<i>Panthera onca</i>	4	Co-author
<i>Cerdocyon thous</i>	10	Co-author	<i>Panthera pardus</i>	4	Movebank
<i>Cervus elaphus</i>	47	Co-author, Eurodeer & Movebank	<i>Papio anubis</i>	4	Movebank
<i>Chlorocebus pygerythrus</i>	12	Movebank	<i>Papio cynocephalus*</i>	22	Co-author & Movebank
<i>Chrysocyon brachyurus</i>	12	Movebank	<i>Procapra gutturosa</i>	15	Co-author
<i>Connochaetes taurinus</i>	3	Co-author	<i>Procyon lotor</i>	9	Movebank
<i>Dasybus novemcinctus</i>	1	Co-author	<i>Propithecus verreauxi*</i>	28	Co-author
<i>Elephas maximus</i>	2	Movebank	<i>Puma concolor</i>	6	Co-author
<i>Equus grevyi</i>	7	Movebank	<i>Rangifer tarandus</i>	14	Co-author
<i>Equus hemionus</i>	6	Co-author	<i>Saguinus geoffroyi*</i>	3	Movebank
<i>Equus quagga</i>	27	Co-author & Movebank	<i>Saiga tatarica</i>	3	Co-author
<i>Eulemur rufifrons</i>	4	Co-author	<i>Sus scrofa</i>	26	Co-author
<i>Euphractus sexcinctus</i>	7	Co-author	<i>Syncerus caffer</i>	6	Movebank
<i>Felis silvestris</i>	5	Movebank	<i>Tamandua mexicana</i>	2	Movebank
<i>Giraffa camelopardalis</i>	5	Co-author	<i>Tapirus terrestris</i>	4	Co-author
<i>Gulo gulo</i>	5	Co-author	<i>Tolypeutes matacus</i>	5	Co-author
<i>Lepus europaeus</i>	39	Movebank	<i>Trichosurus vulpecula*</i>	29	Co-author
<i>Loxodonta africana</i>	14	Co-author & Movebank	<i>Ursus americanus</i>	21	Movebank
<i>Loxodonta africana cyclotis</i>	23	Movebank	<i>Ursus arctos</i>	13	Co-author
<i>Lynx lynx</i>	6	Co-author	<i>Vulpes vulpes</i>	5	Movebank
<i>Lynx rufus</i>	6	Movebank			

* GPS devices turned off during inactive periods to save battery (e.g., primates sleeping overnight in trees) and location data was interpolated during the stationary phases (see Methods in main text).

Table S3.

Model coefficients, r-squared and sample sizes of linear mixed effects models predicting the median and 0.95 quantiles of individual displacements from 1 to 256 hour time scales. Predictor variables included body mass, NDVI, diet and the human footprint index. The model also included a nested random effect accounting for the taxonomy, and a Gaussian spatial autocorrelation structure. We calculated the marginal r^2 (variance explained by the fixed effects) and conditional r^2 (variance explained by both fixed and random factors) values for each model using the “MuMIn” R package (18). Fixed effects included mass, NDVI, the human footprint index and diet. Random effects included taxonomy. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

	1h		2h		4h		8h		16h		32h		64h		128h		256h	
	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%
Mass	0.096	0.288***	0.138	0.268***	0.105	0.297***	0.126	0.288***	0.195*	0.301***	0.265***	0.325***	0.33***	0.321***	0.336***	0.306**	0.423***	0.403***
NDVI	0.004	-0.041*	-0.019	-0.081***	-0.04	-0.078***	-0.067***	-0.086***	-0.056**	-0.078***	-0.115***	-0.161***	-0.124***	-0.155***	-0.144***	-0.158***	-0.132***	-0.172***
HumanF	-0.001	-0.002	-0.004	-0.004*	-0.004	-0.003	-0.006***	-0.005**	-0.01***	-0.009***	-0.009***	-0.01***	-0.009***	-0.011***	-0.009***	-0.011***	-0.009***	-0.014***
Diet (H)	0.225	-0.209	0.175	-0.172	-0.018	-0.363	-0.026	-0.431	-0.342	-0.497*	-0.552*	-0.598*	-0.72**	-0.527	-0.558*	-0.342	-0.638*	-0.46
Diet (O)	0.185	-0.127	0.052	-0.066	-0.006	-0.186	0.073	-0.233	-0.123	-0.248	-0.307	-0.403	-0.494	-0.445	-0.45*	-0.346	-0.492*	-0.398
r² Marginal	0.034	0.286	0.045	0.255	0.016	0.346	0.022	0.35	0.228	0.415	0.349	0.443	0.406	0.347	0.391	0.28	0.459	0.381
r² Conditional	0.922	0.865	0.932	0.895	0.958	0.887	0.977	0.901	0.875	0.885	0.898	0.898	0.906	0.87	0.871	0.846	0.866	0.835
Species	52		53		48		45		42		41		43		46		48	
Individuals	531		606		601		544		525		526		590		598		624	

Table S4.

Model coefficients, r-squared and sample sizes of linear mixed effects models predicting the median and 0.95 quantiles of individual displacements from 1 to 256 hour time scales. Predictor variables included body mass, NDVI, diet and the human footprint index, which was split into the individual-behavioral effect (Ind_HumanF: the individual HFI minus the species mean HFI) and species-occurrence effect (Sp_HumanF: the species mean HFI). The model also included a nested random effect accounting for the taxonomy, and a Gaussian spatial autocorrelation structure. We calculated the marginal r^2 (variance explained by the fixed effects) and conditional r^2 (variance explained by both fixed and random factors) values for each model using the “MuMIn” R package (18). Fixed effects included mass, NDVI, the human footprint index and diet. Random effects included taxonomy. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

	1h		2h		4h		8h		16h		32h		64h		128h		256h	
	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%	Median	95%
Mass	0.129	0.287***	0.143	0.267***	0.127***	0.292	0.116	0.268***	0.203*	0.301***	0.254**	0.301***	0.271**	0.236*	0.279**	0.218*	0.373***	0.33***
NDVI	0.003	-0.041*	-0.019	-0.08***	-0.041	-0.077***	-0.067***	-0.085***	-0.056*	-0.078***	-0.115**	-0.16***	-0.122**	-0.152*	-0.142**	-0.154*	-0.127***	-0.166***
Ind_HumanF	-0.001	-0.002	-0.004	-0.004*	-0.004*	-0.003	-0.006**	-0.005**	-0.01***	-0.009***	-0.009***	-0.01***	-0.009***	-0.011***	-0.008***	-0.01***	-0.008***	-0.013***
Sp_HumanF	0.005	-0.002	-0.003	-0.004	0.001	-0.005	-0.008	-0.01	-0.008	-0.009	-0.011	-0.015	-0.022	-0.031	-0.025	-0.036*	-0.031*	-0.038*
Diet (H)	0.206	-0.209	0.168	-0.172	-0.023	-0.36	-0.035	-0.421	-0.352	-0.497*	-0.544*	-0.571*	-0.626*	-0.46	-0.477	-0.304	-0.66**	-0.42
Diet (O)	0.169	-0.126	0.047	-0.066	-0.018	-0.185	0.068	-0.233	-0.131	-0.249	-0.301	-0.383	-0.424	-0.384	-0.381	-0.288	-0.499*	-0.356
r^2 Marginal	0.037	0.282	0.045	0.252	0.016	0.342	0.023	0.345	0.222	0.407	0.343	0.433	0.394	0.367	0.406	0.323	0.528	0.428
r^2 Conditional	0.921	0.866	0.932	0.896	0.958	0.889	0.978	0.905	0.874	0.886	0.901	0.902	0.913	0.886	0.884	0.87	0.882	0.853
Species	52		53		48		45		42		41		43		46		48	
Individuals	531		606		601		544		525		526		590		598		624	

Table S5.

Summary of the positive (+) and negative (-) effects of barriers and anthropogenic resources on individuals, populations and ecosystems using examples from the literature.

Mechanism	Impact	Level of Impact	Effect of impact	Study Organism	References
Restricted Access to Natural Areas/Barriers	Road barriers alter genetic structure between populations.	Populations	-	Moose (<i>Alces alces</i>); desert bighorn sheep (<i>Ovis canadensis nelsoni</i>)	Wilson <i>et al.</i> (19); Epps <i>et al.</i> (20)
	Altered animal abundance.	Populations	-/+	White-tailed antelope squirrel (<i>Ammospermophilus leucurus</i>), black-tailed prairie dog (<i>Cynomys ludovicianus</i>), Merriam's kangaroo rat (<i>Dipodomys merriami</i>), kangaroo rat (<i>Dipodomys microps</i>), prairie vole (<i>Microtus ochrogaster</i>), California vole (<i>Microtus californicus</i>), house mouse (<i>Mus musculus</i>), woodrat (<i>Notoma lepida</i>), golden mouse (<i>Ochrotomys nuttalli</i>), long-tailed pocket mouse (<i>Perognathus formosus</i>), white-footed mouse (<i>Peromyscus boylii</i>), white-footed mouse (<i>Peromyscus leucopus</i>), deer mouse (<i>Peromyscus maniculatus</i>), rat (<i>Rattus rattus</i>), eastern chipmunk (<i>Tamias striatus</i>), chacoan peccary (<i>Catagonus wagneri</i>), hedgehog (<i>Erinaceus europaeus</i>), brown hare (<i>Lepus europaeus</i>), American marten (<i>Martes americana</i>), badger (<i>Meles meles</i>), koala (<i>Phascolarctos cinereus</i>), white-lipped peccary (<i>Tayassu pecari</i>), collared peccary (<i>Tayassu tajacu</i>), red fox (<i>Vulpes vulpes</i>), Impala (<i>Aepyceros</i>)	Fahrig <i>et al.</i> (21)

				<i>melampus</i>), moose (<i>Alces alces</i>), wolf (<i>Canis lupus</i>), eastern timber wolf (<i>Canis lupus lycaon</i>), black-backed jackal (<i>Canis mesomelas</i>), roe deer (<i>Capreolus capreolus</i>), elk (<i>Cervus canadensis</i>), wildebeest (<i>Connochaetes taurinus</i>), zebra (<i>Equus quagga</i>), giraffe (<i>Giraffa camelopardalis</i>), African elephant (<i>Loxodonta africana</i>), bobcat (<i>Lynx rufus</i>), Eurasian lynx (<i>Lynx lynx</i>), Iberian lynx (<i>Lynx pardinus</i>), mule deer (<i>Odocoileus hemionus</i>), Amur tiger (<i>Panthera tigris altaica</i>), warthog (<i>Phacochoerus africanus</i>), cougar (<i>Puma concolor</i>), woodland caribou (<i>Rangifer tarandus caribou</i>), bohor reedbuck (<i>Redunca redunca</i>), boar (<i>Sus scrofa</i>), eland (<i>Taurotragus oryx</i>), brown bear (<i>Ursus arctos</i>) and grizzly bear (<i>Ursus arctos horribilis</i>).	
	Decreased immigration and colonization success due to barriers.	Populations	-	Animal simulation	Fahrig (22)
	Reproduction, body mass and mobility impact susceptibility to roads.	Individual	-/+	Woodland caribou (<i>Rangifer tarandus</i>), white-footed mouse (<i>Peromyscus leucopus</i>), eastern chipmunk (<i>Tamias striatus</i>), hedgehog (<i>Erinaceus europaeus</i>), bobcat (<i>Lynx rufus</i>), grey wolf (<i>Canis lupus</i>), cougar (<i>Puma concolor</i>), black bear (<i>Ursus americanus</i>), elk (<i>Cervus elaphus</i>), moose (<i>Alces alces</i>) and grizzly bear (<i>Ursus arctos</i>).	Rytwinski <i>et al.</i> (23)
	Dirt tracks/firebreaks can increase seed dispersal.	Ecosystem	+	Wild boar (<i>Sus scrofa</i>), red deer (<i>Cervus elaphus</i>), fallow deer (<i>Dama dama</i>), red fox (<i>Vulpes vulpes</i>), Eurasian badger (<i>Meles meles</i>) and	Suarez-Esteban <i>et al.</i> (24)

				European hare (<i>Lepus europaeus</i>).	
Fragmentation and altered community composition.	Individuals and populations	-	Mammal simulations		Buchmann <i>et al.</i> (25)
Tortuosity increases near roads and trails.	Individuals	-	Wolf (<i>Canis lupus</i>)		Whittington <i>et al.</i> (26)
Small home range and increased overlap near hard boundaries (e.g., roads) and altered genetic composition.	Individuals and populations	-	Coyote (<i>Canis latrans</i>) and bobcats (<i>Lynx rufus</i>).		Riley <i>et al.</i> (27)
Reduced population densities near infrastructure.	Populations	-	Moose (<i>Alces alces</i>), coyote (<i>Canis latrans</i>), red fox (<i>Vulpes vulpes</i>), duiker (<i>Cephalophus</i> sp), elk (<i>Cervus canadensis</i>), blue wildebeest (<i>Connochaetes taurinus</i>), Emin's pouched rat (<i>Cricetomys emini</i>), link rat (<i>Deomys ferrugineus</i>), desert kangaroo rat (<i>Dipodomys deserti</i>), plains zebra (<i>Equus quagga</i>), red-cheeked rope squirrel (<i>Funisciurus leucogenys</i>), shining thicket rat (<i>Grammomys rutilans</i>), African dormice (<i>Graphiurus</i> sp), African smoky mouse (<i>Heimyscus fumosus</i>), Peters' striped mouse (<i>Hybomys univittatus</i>), beaded wood mouse (<i>Hylomyscus aeta</i>), Allen's wood mouse (<i>Hylomyscus alleni</i>), European hare (<i>Lepus europaeus</i>), fire-bellied brush-furred rat (<i>Lophuromys nudicaudus</i>), African elephant (<i>Loxodonta africana</i>), forest elephant (<i>Loxodonta africana cyclotis</i>), bobcat (<i>Lynx rufus</i>), fawn-footed mosaic-tailed rat (<i>Melomys cervinipes</i>), mule deer (<i>Odocoileus hemionus</i>), white-tailed deer (<i>Odocoileus virginianus</i>), Tullberg's soft-furred mouse (<i>Praomys</i>		Benitez-Lopez <i>et al.</i> (28)

				<i>tullbergi</i>), reindeer (<i>Rangifer tarandus</i>), rat (<i>Rattus</i> spp), round-tailed ground squirrel (<i>Spermophilus tereticaudus</i>), target rat (<i>Stochomys longicaudatus</i>), eland (<i>Taurotragus</i> spp), bohor reedbuck (<i>Redunca redunca</i>), giant white-tailed rat (<i>Uromys caudimaculatus</i>), brown bear (<i>Ursus arctos</i>) and black-backed jackal (<i>Canis mesomelas</i>).	
	Reduced population densities near infrastructure and restricted movements caused by infrastructure.	Populations	-	Forest elephants (<i>Loxodonta africana cyclotis</i>).	Blake <i>et al.</i> (29)
	Reduced movements due to human settlements/roads and reduced flow of females between populations.	Individuals and populations	-	Grizzly bears (<i>Ursus arctos</i>).	Proctor <i>et al.</i> (30)
Restricted Access AND Increased Resources	Movements tied to artificial water sources and increased recursive movements due to fences, resulting in increased pressure on local resources.	Individuals, populations and ecosystems	-	African elephant (<i>Loxodonta africana</i>).	Loarie <i>et al.</i> (31)
	Smaller home ranges due to supplemental feeding and road barriers.	Individuals and populations	-	Red deer (<i>Cervus elaphus</i>)	Jerina <i>et al.</i> (32)
	Urban resources as an ecological trap: urban sink populations and urban islands impact population genetic structure/flow and increase in conflict with humans due to expanding population numbers.	Individuals and populations	-	Wild boar (<i>Sus scrofa</i>)	Stillfried <i>et al.</i> (33)
	Increased productivity/reproduction, altered migration timing and increased grazing pressure at winter sites due to supplemental feeding, and population declines due to habitat loss.	Individual, population and ecosystem	-/+	Mule deer (<i>Odocoileus hemionus</i>)	DeVos <i>et al.</i> (34); Sandoval <i>et al.</i> (35); Peterson <i>et al.</i> (36) ; Bishop <i>et al.</i> (37).

	Landscape elements (e.g., fruit trees) act as food supplements, allowing populations to persist in fragmented landscapes.	Individuals and populations.	+	Howler monkeys (<i>Alouatta palliata mexicana</i>)	Asensio <i>et al.</i> (38)
Increased Resources (Anthropogenic)	Crop damage leading to human-wildlife conflict.	Individuals and populations	-	Wild boars (<i>Sus scrofa</i>); Red deer (<i>Cervus elaphus</i>).	Honda <i>et al.</i> (39); Barrios-Garcia <i>et al.</i> (40); Bleier <i>et al.</i> (41)
	Increase in parasite load and diseases.	Individual and population	-	Elk (<i>Cervus canadensis</i>); white-tailed deer (<i>Odocoileus virginianus</i>).	Hines <i>et al.</i> (42); Miller <i>et al.</i> (43); Sorensen <i>et al.</i> (44)
	Increase group size.	Population	+	Arctic fox (<i>Vulpes lagopus</i>).	Elmhagen <i>et al.</i> (45)
	Increased survival rate, increased reproductive rate, improved winter condition, increased hunting, increased population growth rate and reduced density dependence, changed spatial genetic structure, reduced natural selection, increased aggression, increased stress, increased local browsing or grazing, changed plant species composition, invasion of non-native weed species, increased parasitism due to spatial aggregation and increased contact rates and reduced parasitism due to improved body condition.	Individual, population and ecosystem	-/+	European bison (<i>Bison bonasus</i>), wild boar (<i>Sus scrofa</i>), white-tailed deer (<i>Odocoileus virginianus</i>), elk (<i>Cervus canadensis</i>) and moose (<i>Alces alces</i>).	Milner <i>et al.</i> (46)
	Disruption of movement patterns, circadian rhythm, denning behavior, increased individual interactions, increase population size, culling, increase in diseases, human-animal conflict, alter natural foraging and trophic cascades.	Individual, population and ecosystem	-/+	Brown bears (<i>Ursus arctos</i>).	Penteriani <i>et al.</i> (47)
	Consumption of valuable tree species, altered social structure, space	Individual, population and	-/+	European bison (<i>Bison bonasus</i>); moose (<i>Alces alces</i>).	Kowalczyk <i>et al.</i> (48); Mathisen <i>et</i>

use and parasites.	ecosystem			<i>al. (49)</i>
Sustain populations in resource poor areas and trophic cascades.	Population and ecosystem	-/+	Dingo (<i>Canis lupus dingo</i>).	Newsome <i>et al.</i> (50, 51)
Trophic cascades.	Ecosystem	-	African wild dog (<i>Lycaon pictus</i>), yellow baboon (<i>Papio cynocephalus</i>), black-backed jackal (<i>Canis mesomelas</i>), bobcat (<i>Lynx rufus</i>), chilla fox (<i>Pseudalopex griseus</i>), coyote (<i>Canis latrans</i>), culpeo fox (<i>Pseudalopex culpaeus</i>), dhole (<i>Cuon alpinus</i>), common genet (<i>Genetta genetta</i>), Geoffroy's cat (<i>Oncifelis geoffroyii</i>), golden jackal (<i>Canis aureus</i>), Indian fox (<i>Vulpes bengalensis</i>), pampas fox (<i>Pseudalopex gymnocercus</i>), red fox (<i>Vulpes vulpes</i>) and San Joaquin kit fox (<i>Vulpes macrotis mutica</i>), Arabian wolf (<i>Canis lupus arabs</i>), black bear (<i>Ursus americanus</i>), brown bear (<i>Ursus arctos</i>), cheetah (<i>Acinonyx jubatus</i>), dingo (<i>Canis dingo</i>), Ethiopian wolf (<i>Canis simensis</i>), Eurasian lynx (<i>Lynx lynx</i>), grey wolf (<i>Canis lupus</i>), Mexican grey wolf (<i>Canis lupus baileyi</i>), Iberian lynx (<i>Lynx pardinus</i>), Iberian wolf (<i>Canis lupus signatus</i>), jaguar (<i>Panthera onca</i>), leopard (<i>Panthera pardus</i>), lion (<i>Panthera leo</i>), polar bear (<i>Ursus maritimus</i>), puma (<i>Puma concolor</i>), snow leopard (<i>Panthera uncia</i>), spotted hyena (<i>Crocuta crocuta</i>), tiger (<i>Panthera tigris</i>); white-tailed deer (<i>Odocoileus virginianus</i>); moose (<i>Alces alces</i>).	Newsome <i>et al.</i> (52); Cooper <i>et al.</i> (53); Gundersen <i>et al.</i> (54)

Increase in stress hormones.	Individual	-	Asiatic black bears (<i>Ursus thibetanus</i>).	Malcolm <i>et al.</i> (55)
Animal-human conflict: death and monetary costs.	Population	-	Brown bear (<i>Ursus arctos</i>).	Kavčič <i>et al.</i> (56)
Reduced natural selection effects on juveniles.	Individual and population	+	Red deer (<i>Cervus elaphus</i>).	Schmidt <i>et al.</i> (57)
Reduced and stable home range size due to resources.	Individual	+	Racoon (<i>Procyon lotor</i>) ; Roe deer (<i>Capreolus capreolus</i>) ; Red deer (<i>Cervus elaphus</i>); Iberian lynx (<i>Lynx pardinus</i>).	Prange <i>et al.</i> (58); Ossi <i>et al.</i> (59); Lopez-Bao <i>et al.</i> (60)
Reduce migration distance and time spent at summer grounds (less quality forage).	Individual	-	Elk (<i>Cervus canadensis</i>).	Jones <i>et al.</i> (61)
Smaller home range size, covered more distance, nocturnal activity and increase movement speeds.	Individual	+	Wild boar (<i>Sus scrofa</i>).	Podgorski <i>et al.</i> (62)
Anthropogenic food resources reduce home range size and increases home range overlap, with implications for rabies transmission between individuals.	Individual and populations	-	Indian mongoose (<i>Herpestes javanicus</i>).	Quinn <i>et al.</i> (63)
Food provisions impact movement behaviors, amplify pathogen invasion due to increased host aggregation and tolerance, but also reduces transmission if provisioned food decreases dietary exposure to parasites.	Individuals and populations	-/+	Elk (<i>Cervus canadensis</i>), long-tail macaque (<i>Macaca fascicularis</i>) , red fox (<i>Vulpe vulpes</i>), white-tailed deer (<i>Odocoileus virginianus</i>), common vampire bat (<i>Desmodus rotundus</i>) and flying fox (<i>Pteropus giganteus</i>).	Becker <i>et al.</i> (64)
Anthropogenic resources reduce home range size and increases livestock kills by wildlife.	Individuals	-	Spotted hyena (<i>Crocuta crocuta</i>).	Kolowski <i>et al.</i> (65)
Anthropogenic food reduced core home range size and increases population size.	Individuals and populations	+	Banded mongoose (<i>Mungos mungo</i>).	Gilchrist <i>et al.</i> (66)

Reference and Notes

1. H. L. A. Bartlam-Brooks, P. S. A. Beck, G. Bohrer, S. Harris, Data from: In search of greener pastures: using satellite images to predict the effects of environmental change on zebra migration. *Movebank data Repos.* (2013).
2. H. L. A. Bartlam-Brooks, P. S. A. Beck, G. Bohrer, S. Harris, In search of greener pastures: Using satellite images to predict the effects of environmental change on zebra migration. *J. Geophys. Res. Biogeosciences.* **118**, 1427–1437 (2013).
3. J. Wall, G. Wittemyer, V. LeMay, I. Douglas-Hamilton, B. Klinkenberg, Data from: Elliptical Time-Density model to estimate wildlife utilization distributions. *Movebank data Repos.* (2014).
4. J. Wall, G. Wittemyer, V. LeMay, I. Douglas-Hamilton, B. Klinkenberg, Elliptical Time-Density model to estimate wildlife utilization distributions. *Methods Ecol. Evol.* **5**, 780–790 (2014).
5. M. Rimmler, T. Mueller, SyncMove: Subsample Temporal Data to Synchronal Events and Compute the MCI. R package version 0.1-0 (2015), (available at <http://cran.r-project.org/package=SyncMove>).
6. F. Chambat, B. Valette, Mean radius, mass, and inertia for reference Earth models. *Phys. Earth Planet. Inter.* **124**, 237–253 (2001).
7. K. Bjørneraas, B. Van Moorter, C. M. Rolandsen, I. Herfindal, Screening Global Positioning System Location Data for Errors Using Animal Movement Characteristics. *J. Wildl. Manage.* **74**, 1361–1366 (2010).
8. O. Venter *et al.*, Data from: Global terrestrial Human Footprint maps for 1993 and 2009. *Sci. Data* (2016), , doi:doi:10.5061/dryad.052q5.
9. K. Didan, MOD13A1 MODIS/Terra Vegetation Indices 16-Day L3 Global 500m SIN Grid V005. NASA EOSDIS Land Processes DAAC. <https://doi.org/10.5067/MODIS/MOD13A1.006> (2015).
10. S. Dodge *et al.*, The environmental-data automated track annotation (Env-DATA) system: linking animal tracks with environmental data. *Mov. Ecol.* **1**, 1 (2013).
11. K. E. Jones *et al.*, PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology.* **90**, 2648 (2009).
12. C. F. Dormann *et al.*, Methods to account for spatial autocorrelation in the analysis of species distributional data: a review. *Ecography (Cop.)*. **30**, 609–628 (2007).
13. C. F. Dormann *et al.*, Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography (Cop.)*. **36**, 27–46 (2013).
14. A. F. Zuur, E. N. Ieno, C. S. Elphick, A protocol for data exploration to avoid common statistical problems. *Methods Ecol. Evol.* **1**, 3–14 (2010).
15. M. J. Mazerolle, AICcmodavg: Model selection and multimodel inference based on (Q)AIC(c). R package version 2.1-0.itle (2016), (available at <https://cran.r-project.org/package=AICcmodavg>).
16. R. D. C. Team, R: A Language and Environment for Statistical Computing. Vienna, Austria, ISBN 3-900051-07-0. <http://www> (2012).
17. O. Venter *et al.*, Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nat. Commun.* **7** (2016).
18. K. Barton, MuMIn: Multi-Model Inference. R package version 1.15.6e (2016),

- (available at <http://cran.r-project.org/package=MuMIn>).
19. R. E. Wilson, S. D. Farley, T. J. McDonough, S. L. Talbot, P. S. Barboza, A genetic discontinuity in moose (*Alces alces*). *Conserv. Genet.* **16**, 791–800 (2015).
 20. C. W. Epps *et al.*, Highways block gene flow and cause a rapid decline in genetic diversity of desert bighorn sheep. *Ecol. Lett.* **8**, 1029–1038 (2005).
 21. L. Fahrig, T. Rytwinski, Effects of roads on animal abundance: an empirical review and synthesis. *Ecol. Soc.* **14** (2009).
 22. L. Fahrig, Non-optimal animal movement in human-altered landscapes. *Funct. Ecol.* **21**, 1003–1015 (2007).
 23. T. Rytwinski, L. Fahrig, Do species life history traits explain population responses to roads? A meta-analysis. *Biol. Conserv.* **147**, 87–98 (2012).
 24. A. Suárez-Esteban, M. Delibes, J. M. Fedriani, Barriers or corridors? The overlooked role of unpaved roads in endozoochorous seed dispersal. *J. Appl. Ecol.* **50**, 767–774 (2013).
 25. C. M. Buchmann, F. M. Schurr, R. Nathan, F. Jeltsch, Habitat loss and fragmentation affecting mammal and bird communities—The role of interspecific competition and individual space use. *Ecol. Inform.* **14**, 90–98 (2013).
 26. J. Whittington, C. C. St Clair, G. Mercer, Path tortuosity and the permeability of roads and trails to wolf movement. *Ecol. Soc.* **9**, 4 (2004).
 27. S. P. D. Riley *et al.*, FAST-TRACK: A southern California freeway is a physical and social barrier to gene flow in carnivores. *Mol. Ecol.* **15**, 1733–1741 (2006).
 28. A. Benítez-López, R. Alkemade, P. A. Verweij, The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biol. Conserv.* **143**, 1307–1316 (2010).
 29. S. Blake *et al.*, Roadless Wilderness Area Determines Forest Elephant Movements in the Congo Basin. *PLoS One.* **3**, e3546 (2008).
 30. M. F. Proctor *et al.*, Population fragmentation and inter-ecosystem movements of grizzly bears in western Canada and the northern United States. *Wildl. Monogr.* **180**, 1–46 (2012).
 31. S. R. Loarie, R. J. Van Aarde, S. L. Pimm, Fences and artificial water affect African savannah elephant movement patterns. *Biol. Conserv.* **142**, 3086–3098 (2009).
 32. K. Jerina, Roads and supplemental feeding affect home-range size of Slovenian red deer more than natural factors. *J. Mammal.* **93**, 1139–1148 (2012).
 33. M. Stillfried *et al.*, Do cities represent sources, sinks or isolated islands for urban wild boar population structure? *J. Appl. Ecol.* (2016).
 34. J. C. DeVos, M. R. Conover, N. E. Headrick, *Mule deer conservation: issues and management strategies* (Jack H. Berryman Institute Press, Utah State University, 2003).
 35. L. Sandoval, J. Holechek, J. Biggs, R. Valdez, D. VanLeeuwen, Elk and Mule Deer Diets in North-Central New Mexico. *Rangel. Ecol. Manag.* **58**, 366–372 (2005).
 36. C. Peterson, T. A. Messmer, Effects of Winter-Feeding on Mule Deer in Northern Utah. *J. Wildl. Manage.* **71**, 1440–1445 (2007).
 37. C. J. Bishop, G. C. White, D. J. Freddy, B. E. Watkins, T. R. Stephenson, Effect of Enhanced Nutrition on Mule Deer Population Rate of Change. *Wildl. Monogr.*, 1–

- 28 (2009).
38. N. Asensio, V. Arroyo-Rodríguez, J. C. Dunn, J. Cristóbal-Azkarate, Conservation value of landscape supplementation for howler monkeys living in forest patches. *Biotropica*. **41**, 768–773 (2009).
 39. T. Honda, M. Sugita, Environmental factors affecting damage by wild boars (*Sus scrofa*) to rice fields in Yamanashi Prefecture, central Japan. *Mammal Study*. **32**, 173–176 (2007).
 40. M. N. Barrios-Garcia, S. A. Ballari, Impact of wild boar (*Sus scrofa*) in its introduced and native range: a review. *Biol. Invasions*. **14**, 2283–2300 (2012).
 41. N. Bleier, R. Lehoczki, D. Újváry, L. Szemethy, S. Csányi, Relationships between wild ungulates density and crop damage in Hungary. *Acta Theriol. (Warsz)*. **57**, 351–359 (2012).
 42. A. M. Hines, V. O. Ezenwa, P. Cross, J. D. Rogerson, Effects of supplemental feeding on gastrointestinal parasite infection in elk (*Cervus elaphus*): Preliminary observations. *Vet. Parasitol.* **148**, 350–355 (2007).
 43. R. Miller, J. B. Kaneene, S. D. Fitzgerald, S. M. Schmitt, Evaluation of the influence of supplemental feeding of white-tailed deer (*Odocoileus virginianus*) on the prevalence of bovine tuberculosis in the Michigan wild deer population. *J. Wildl. Dis.* **39**, 84–95 (2003).
 44. A. Sorensen, F. M. van Beest, R. K. Brook, Impacts of wildlife baiting and supplemental feeding on infectious disease transmission risk: a synthesis of knowledge. *Prev. Vet. Med.* **113**, 356–363 (2014).
 45. B. Elmhagen, P. Hersteinsson, K. Norén, E. R. Unnsteinsdottir, A. Angerbjörn, From breeding pairs to fox towns: the social organisation of arctic fox populations with stable and fluctuating availability of food. *Polar Biol.* **37**, 111–122 (2014).
 46. J. M. Milner, F. M. Van Beest, K. T. Schmidt, R. K. Brook, T. Storaas, To feed or not to feed? Evidence of the intended and unintended effects of feeding wild ungulates. *J. Wildl. Manage.* **78**, 1322–1334 (2014).
 47. V. Penteriani *et al.*, Consequences of brown bear viewing tourism: A review. *Biol. Conserv.* **206**, 169–180 (2017).
 48. R. Kowalczyk *et al.*, Influence of management practices on large herbivore diet—Case of European bison in Białowieża Primeval Forest (Poland). *For. Ecol. Manage.* **261**, 821–828 (2011).
 49. K. M. Mathisen, J. M. Milner, F. M. van Beest, C. Skarpe, Long-term effects of supplementary feeding of moose on browsing impact at a landscape scale. *For. Ecol. Manage.* **314**, 104–111 (2014).
 50. T. M. Newsome, G.-A. Ballard, C. R. Dickman, P. J. S. Fleming, C. Howden, Anthropogenic Resource Subsidies Determine Space Use by Australian Arid Zone Dingoes: An Improved Resource Selection Modelling Approach. *PLoS One*. **8**, e63931 (2013).
 51. T. M. Newsome *et al.*, Human-resource subsidies alter the dietary preferences of a mammalian top predator. *Oecologia*. **175**, 139–150 (2014).
 52. T. M. Newsome *et al.*, The ecological effects of providing resource subsidies to predators. *Glob. Ecol. Biogeogr.* **24**, 1–11 (2015).
 53. S. M. Cooper, M. K. Owens, R. M. Cooper, T. F. Ginnett, Effect of supplemental feeding on spatial distribution and browse utilization by white-tailed deer in semi-

- arid rangeland. *J. Arid Environ.* **66**, 716–726 (2006).
54. H. Gundersen, H. P. Andreassen, T. Storaas, Supplemental feeding of migratory moose *Alces alces*: forest damage at two spatial scales. *Wildlife Biol.* **10**, 213–223 (2004).
 55. K. D. Malcolm *et al.*, Increased stress in Asiatic black bears relates to food limitation, crop raiding, and foraging beyond nature reserve boundaries in China. *Glob. Ecol. Conserv.* **2**, 267–276 (2014).
 56. I. Kavčič *et al.*, Fast food bears: brown bear diet in a human-dominated landscape with intensive supplemental feeding. *Wildlife Biol.* **21**, 1–8 (2015).
 57. K. T. Schmidt, H. Hoi, Supplemental feeding reduces natural selection in juvenile red deer. *Ecography (Cop.)*. **25**, 265–272 (2002).
 58. S. Prange, S. D. Gehrt, E. P. Wiggers, Influences of anthropogenic resources on raccoon (*Procyon lotor*) movements and spatial distribution. *J. Mammal.* **85**, 483–490 (2004).
 59. F. Ossi *et al.*, Plastic response by a small cervid to supplemental feeding in winter across a wide environmental gradient. *Ecosphere*. **8**, e01629–n/a (2017).
 60. J. V López-Bao, F. Palomares, A. Rodríguez, M. Delibes, Effects of food supplementation on home-range size, reproductive success, productivity and recruitment in a small population of Iberian lynx. *Anim. Conserv.* **13**, 35–42 (2010).
 61. J. D. Jones *et al.*, Supplemental feeding alters migration of a temperate ungulate. *Ecol. Appl.* **24**, 1769–1779 (2014).
 62. T. Podgórski *et al.*, Spatiotemporal behavioral plasticity of wild boar (*Sus scrofa*) under contrasting conditions of human pressure: primeval forest and metropolitan area. *J. Mammal.* **94**, 109–119 (2013).
 63. J. H. Quinn, D. A. Whisson, The effects of anthropogenic food on the spatial behaviour of small Indian mongooses (*Herpestes javanicus*) in a subtropical rainforest. *J. Zool.* **267**, 339–350 (2005).
 64. D. J. Becker, D. G. Streicker, S. Altizer, Linking anthropogenic resources to wildlife–pathogen dynamics: a review and meta-analysis. *Ecol. Lett.* **18**, 483–495 (2015).
 65. J. M. Kolowski, K. E. Holekamp, Effects of an open refuse pit on space use patterns of spotted hyenas. *Afr. J. Ecol.* **46**, 341–349 (2008).
 66. J. S. Gilchrist, E. Otali, The effects of refuse-feeding on home-range use, group size, and intergroup encounters in the banded mongoose. *Can. J. Zool.* **80**, 1795–1802 (2002).