

Animal welfare considerations for using large carnivores and guardian dogs as vertebrate biocontrol tools against other animals

Benjamin L. Allen^{1,2*}, Lee R. Allen³, Guy Ballard^{4,5}, Marine Drouilly⁶, Peter J.S. Fleming^{4,7}, Jordan O. Hampton⁸, Matthew W. Hayward^{2,9,10}, Graham I.H. Kerley¹⁰, Paul D. Meek^{4,11}, Liaan Minnie¹², M. Justin O'Riain⁶, Daniel M. Parker^{12,13}, Michael J. Somers¹⁴.

Author affiliations

¹University of Southern Queensland, Institute for Life Sciences and the Environment, Toowoomba, Queensland 4350, Australia. Email: benjamin.allen@usq.edu.au

²Centre for Invasion Biology, Mammal Research Institute, University of Pretoria, Pretoria 0002, South Africa.

³Robert Wicks Pest Animal Research Centre, Biosecurity Queensland, Toowoomba, Queensland 4350, Australia.

⁴Ecosystem Management, School of Environmental and Rural Science, University of New England, Armidale, New South Wales 2351, Australia.

⁵Vertebrate Pest Research Unit, New South Wales Department of Primary Industries, Armidale, New South Wales 2350, Australia.

⁶Institute for Communities and Wildlife in Africa, Department of Biological Sciences, University of Cape Town, Upper Campus, Rondebosch, 7700, South Africa.

⁷Vertebrate Pest Research Unit, New South Wales Department of Primary Industries, Orange, New South Wales 2800, Australia.

⁸Murdoch University, Murdoch, Western Australia 6150, Australia.

⁹Conservation Biology Lab, School of Environmental and Life Sciences, University of Newcastle, Callaghan New South Wales 2308, Australia.

¹⁰Centre for African Conservation Ecology, Nelson Mandela University, Port Elizabeth 6034, South Africa.

¹¹Vertebrate Pest Research Unit, New South Wales Department of Primary Industries, National Marine Science Centre, Coffs Harbour, New South Wales 2450, Australia.

¹²School of Biology and Environmental Sciences, University of Mpumalanga, Nelspruit 1200, South Africa.

¹³Wildlife and Reserve Management Research Group, Department of Zoology and Entomology, Rhodes University, Grahamstown 6140, South Africa.

¹⁴Centre for Invasion Biology, Eugène Marais Chair of Wildlife Management, Mammal Research Institute, University of Pretoria, Pretoria 0002, South Africa.

*Corresponding author.

Acknowledgements

We thank Glen Saunders and Trudy Sharp for ongoing discussions about the five domains assessment model for invasive animal control tools. Gail Potgeiter assisted with the assessment and contributed substantial input to the manuscript, including provision of many helpful comments on earlier drafts. BA and MJS thank the DST-NRF Centre of Excellence for Invasion Biology in Stellenbosch and Pretoria, South Africa, for providing funding that enabled the development of this study.

Author contributions

BA initiated the concept of using the five domains approach to assess the animal welfare impacts of dingoes as a biocontrol tool against other species. GB and PF expanded the concept to include guardian dogs, and LM and DP expanded it further to include leopards. BA led the writing, all authors participated in the assessment and contributed to the results, and all authors contributed to the writing and editing of the manuscript.

Abstract

Introducing consumptive and non-consumptive effects into food webs can have profound effects on individuals, populations and communities. This knowledge has led to the deliberate use of predation and/or fear of predation as an emerging technique for controlling wildlife. Many now advocate for the intentional use of large carnivores and livestock guardian dogs as more desirable alternatives to traditional wildlife control approaches like fencing, shooting, trapping, or poisoning. However, there has been very little consideration of the animal welfare implications of deliberately using predation as a wildlife management tool. We assess the animal welfare impacts of using dingoes, leopards and guardian dogs as biocontrol tools against wildlife in Australia and South Africa following the 'Five Domains' model commonly used to assess other wildlife management tools. Application of this model indicates that large carnivores and guardian dogs cause considerable lethal and non-lethal animal welfare impacts to the individual animals they are intended to control. These impacts are likely similar across different predator-prey systems, but are dependent on specific predator-prey combinations; combinations that result in short chases and quick kills will be rated as less harmful than those that result in long chases and protracted kills. Moreover, these impacts are typically rated greater than those caused by traditional wildlife control techniques. The intentional lethal and non-lethal harms caused by large carnivores and guardian dogs should not be ignored or dismissively assumed to be negligible. A greater understanding of the impacts they impose would benefit from empirical studies of the animal welfare outcomes arising from their use in different contexts.

Keywords: animal ethics; animal welfare; biocontrol; decision matrix; dingo; guardian dog; fear effects; humaneness; landscape of fear; leopard; predator-prey relationships.

Introduction

Large carnivores influence ecosystems through consumptive and non-consumptive effects. They frighten, displace, harass, chase, attack and kill other animals (e.g. [van Bommel 2010](#); [Thorn et al. 2012](#); [Potgieter et al. 2013](#); [Fleming et al. 2014](#)). Adding predators to multi-predator multi-prey systems can produce a variety of outcomes ([Hairston et al. 1960](#); [Holt and Lawton 1994](#)), including profound welfare effects on other species ([Fleming et al. 2012](#)). Both the non-lethal and lethal impacts of predators cause distress and/or death to individual prey ([Fox 1969](#); [Power and Compion 2009](#); [Behrendorff et al. 2018](#)), which can lead to population declines and local extinction of some species ([Kruuk 1971](#); [Sinclair et al. 1998](#); [Woinarski et al. 2015](#)) and population increases and recoveries of others ([Terborgh and Estes 2010](#); [Estes et al. 2013](#); [Ripple et al. 2014](#)). Besides killing prey, predators also wound prey and act as vectors for pathogens debilitating to prey, such as *Echinococcus granulosus* or *Neospora caninum* (e.g. [Barnes et al. 2008](#); [King et al. 2011](#)). Predators can also alter space use and foraging patterns of individual prey and prey populations ([Ripple and Beschta 2004](#); [Atwood et al. 2009](#); [Moll et al. 2017](#); [Palmer et al. 2017](#)), causing local extinctions in places where these effects are strong. The direct effects of predation can be a driver of ecosystem structure ([Hairston et al. 1960](#); [Mech 1966](#); [Barbosa and Castellanos 2005](#); [Eisenberg 2011](#); [Linnell 2011](#); [Peterson et al. 2014](#)), as can indirect effects ([Brown et al. 1999](#); [Creel and Christianson 2008](#); [Clinchy et al. 2013](#); [Bleicher 2017](#)). Thus, consumptive and non-consumptive mechanisms are expected to be present and strongly influence species abundance, distribution and behaviour in multi-predator multi-prey systems. These direct and indirect effects make predators attractive as potential biocontrol tools for use against susceptible wildlife species that humans wish to control.

Human conflict with wildlife is a global issue, and a wide variety of wildlife control tools and techniques are used to reduce the distribution, abundance and impacts of the animals involved (e.g. [Fleming et al. 2014](#); [du Plessis et al. 2018](#)). Lethal techniques include shooting, trapping and poisoning (e.g. [Bothma 1971](#); [Saunders and McLeod 2007](#)), and aim to maximise mortality. Non-lethal techniques include aversive conditioning or deterrents (e.g. [Breck et al. 2017](#); [Smith and Appleby 2018](#)), and aim to scare or displace animals. Other techniques, such as exclusion fencing ([de Tores and Marlow 2011](#); [Allen and West 2013](#)) and guardian animals ([Potgieter et al. 2013](#); [Linnell and Lescureux 2015](#); [Allen et al. 2016](#)), operate in both lethal and non-lethal ways. All wildlife control tools are typically applied against a select number of target species within multi-predator multi-prey systems, sometimes producing variable and uncertain outcomes ([Treves et al. 2016](#); [Lennox et al. 2018](#); [van Eeden et al. 2018](#); [Campbell et al. 2019](#)).

Encouraged by the positive ecological changes reported following the reintroduction of grey wolves (*Canis lupus*) to Yellowstone National Park in North America (e.g. [Ripple and Beschta 2012](#)) and many success stories of using livestock guardian dogs (*Canis familiaris*, such as Maremmas) to protect livestock (e.g. [van Bommel 2010](#); [Potgieter et al. 2013](#); [Linnell and Lescureux 2015](#)), many now advocate for the intentional use of large carnivores and/or guardian dogs as more desirable 'natural' alternatives to traditional approaches to wildlife control like fencing, shooting, trapping, or poisoning (e.g. [Ritchie et al. 2012](#); [van Bommel and Johnson 2012](#); [Letnic 2014](#); [Wallach 2014](#); [Minnie et al. 2015](#); [Newsome et al. 2015](#); [Atkins et al. 2017](#)). But while such traditional tools have often been subject to intense and repeated formal assessment of their efficacy (e.g. [Eldridge et al. 2002](#); [Allen et al. 2014](#); [Campbell et al. 2019](#)) and welfare impacts (e.g. [Fleming et al. 1998](#); [Marks et al. 2004](#); [Meek et al. 2019](#); [Allen et al. In press](#)), to date there has been very little consideration of

the animal welfare implications of deliberately using predation and/or fear of predation as a wildlife management tool ([Allen et al. 2017](#)). This is symptomatic of the disproportionate animal welfare scrutiny that is applied to different conservation practices, with human-caused predation representing an under-addressed issue ([Hampton and Hyndman In press](#)).

Predation is not usually considered an animal welfare issue given that the behaviour is not usually considered to be anthropogenic, but rather natural. Nearly all ethical viewpoints applied to animal welfare require consideration of only those processes that humans impose on animals ([Palmer 2010](#); [Fraser and MacRae 2011](#)), though this rule is complicated and is not universal ([Torres 2015](#); [Horta 2017](#)). When predation occurs between wild animals living independent of humans, few (if any) ethicists would consider humans to have a moral responsibility to consider the welfare of prey ([Palmer 2010](#)). However, when predators are deliberately introduced (or reintroduced) into an ecosystem by humans, a process sometimes referred to as rewilding ([Soulé and Noss 1998](#)), predator effects on prey could be considered to be anthropogenic and therefore warrant ethical scrutiny. Some authors have considered the implications of knowingly exposing prey to predation when prey are introduced or reintroduced (e.g. [Swaigood 2010](#); [Harrington et al. 2013](#)), but few authors have considered the implications of intentionally causing predation when predators are introduced or reintroduced ([Allen et al. 2017](#)). Here we refer to the deliberate release, reintroduction or deployment of predators or use of predation or fear of predation as anthropogenic predation. Alternative viewpoints may not consider such predation events to be anthropogenic or may view the level of human responsibility for them to attenuate over time. This question is complex, but important, because it determines whether humans are responsible for the animal welfare impacts arising from introducing predators or not. We assume that humans are responsible for the purposes of this assessment, but we revisit this issue later.

Most (if not all) methods for managing wildlife are controversial (e.g. [Fitzgerald 2009](#); [Suryawanshi et al. 2014](#); [Linnell et al. 2017](#); [Mormile and Hill 2017](#); [Slagle et al. 2017](#)). A prominent source of contention is the animal welfare impacts imposed on target and non-target species, and the ethical issues these raise ([Dubois et al. 2017](#)). For example, poisoning is commonly used to control dingoes (*Canis familiaris*), coyotes (*Canis latrans*) and black-backed jackals (*Canis mesomelas*; hereafter jackals) in attempts to reduce their impacts on sheep (*Ovis aries*) and goats (*Capra hircus*; e.g. [Fleming et al. 2014](#); [Minnie et al. 2016a](#); [du Plessis et al. 2018](#)). However, poisoning (like all other approaches) is often undertaken without first demonstrating that the target individuals or species are actually impacting livestock, without measuring the efficacy of poisoning at reducing these livestock impacts, and without assessing potential impacts of poisoning on non-target species. These 'shots in the dark' (*sensu* [Treves et al. 2016](#)) have raised ethical concerns about the justification for implementation of predator control (e.g. [Marks et al. 2000](#); [Berger 2006](#); [Reddiex et al. 2006](#); [Allen et al. 2014](#); [Doherty and Ritchie 2017](#)). Related to, but distinct from ethical concerns, critics have also highlighted the many animal welfare impacts associated with common wildlife control tools ([Sherley 2007](#); [Twigg and Parker 2010](#); [Littin et al. 2014](#); [Mallick et al. 2016](#); [RSPCA 2016](#)), calling for their replacement with techniques claimed to be less-harmful or for the cessation of wildlife control altogether (e.g. [Letnic 2014](#); [McManus et al. 2014](#); [AJP 2015](#); [Smith and Appleby 2018](#)). Recognition of these concerns and the ongoing desire to improve wildlife control methods have contributed to the refinement of wildlife management tools and techniques. This includes, for example, the development and adoption of less-harmful traps and similar devices, less-harmful poisons, and more-effective strategies for their use (e.g. [Marks et al. 2004](#); [Anon 2014](#); [Eason et al. 2014](#); [Meek et](#)

[al. 2018](#); [Meek et al. 2019](#); [Allen et al. In press](#)). Great progress has also been made in assessing the relative welfare impacts of wildlife management tools and techniques in some regions of the world (e.g. [Sharp and Saunders 2011](#); [Littin et al. 2014](#); [Baker et al. 2016](#); [Hampton et al. 2016](#)). These assessments have included examination of both direct and intentional impacts (e.g. those that arise from poisoning), and also indirect and unintentional impacts (e.g. those that arise from fencing), which are collectively referred to as ‘harms’ (*sensu* [Fraser and MacRae 2011](#)).

Here, we assess the animal welfare impacts of deploying large carnivores and guardian dogs as biocontrol tools against a variety of wildlife known or suspected to be in conflict with humans. Our goal is to provide an assessment of the animal welfare impacts associated with this control technique to assist wildlife managers and decision makers in identifying the most appropriate wildlife control technique for a given situation. We hope to highlight knowledge gaps and stimulate discussion on the advantages and disadvantages of introducing large carnivores and guardian dogs to multi-predator multi-prey systems, and whether or not their use truly can be considered to represent an animal welfare improvement over the traditional tools commonly used by wildlife managers worldwide.

Methods

Study systems

We assessed the animal welfare impacts of large carnivores and guardian dogs as biocontrol tools in two different ecological systems: Australia and South Africa.

In Australia, dingoes have been proposed as biocontrol tools for use against European red foxes (*Vulpes vulpes*), feral cats (*Felis catus*), European rabbits (*Oryctolagus cuniculus*), feral pigs (*Sus scrofa*), feral goats, exotic small mammals such as house mice (*Mus musculus*) and black rats (*Rattus rattus*), and overabundant native macropods such as red kangaroos (*Osphranter rufus*; e.g. [Letnic et al. 2012](#); [Ritchie et al. 2012](#); [Wallach 2014](#)). In some parts of Australia (such as the sheep farming zone in the south-east), all of these species co-occur, interact with each other, and cause considerable social, economic and environmental impacts ([West 2008](#); [McLeod 2016](#)). Dingoes are absent from or occur in low densities across much of this zone given their historical extirpation to enable sheep and goat production ([Yelland 2001](#); [Allen and West 2013, 2015](#)). Dingoes are Australia’s largest non-human terrestrial predator and kill, consume and elicit fear in these prey species ([Glen and Dickman 2005](#); [Letnic et al. 2012](#); [Allen and Leung 2014](#); [Allen et al. 2018](#)). As such, some have proposed the intentional reintroduction and active restoration of the extant dingo populations within this zone as a means of suppressing target wildlife species (e.g. [Wallach 2014](#); [Newsome et al. 2015](#)). Opposition has arisen from livestock producers and other stakeholders about the potential impacts of dingoes on livestock and threatened native fauna (e.g. [Allen and Fleming 2012](#)). In response, proponents further advocate for the broad-scale deployment of livestock guardian dogs to protect the tens of millions of livestock from the recovering dingo population ([van Bommel and Johnson 2012](#); [Smith and Appleby 2018](#)). The proponents envisage a scenario where dingoes will control target wildlife, guardian dogs will protect livestock from dingoes, and consequently humans will not need to engage in lethal wildlife control. This approach has been described by proponents as a “humane”, “virtuous”, “predator-friendly”, “efficient”, “compassionate, cost-effective, sustainable and ethical approach” to problematic wildlife control

(e.g. [Ritchie et al. 2012](#); [van Bommel and Johnson 2012](#); [Wallach 2014](#); [Wallach et al. 2015](#); [Johnson and Wallach 2016](#)).

In South Africa, leopards (*Panthera pardus*) have been proposed as biocontrol tools for use against black-backed jackals and caracals (*Caracal caracal*) on sheep and goat farms ([Minnie et al. 2015](#); [du Plessis et al. 2018](#)), and also for managing chacma baboons (*Papio ursinus*), warthogs (*Phacochoerus africanus*), Himalayan tahr (*Hemitragus jemlahicus*) and other species in other areas. In some parts of South Africa, jackals and caracals occupy the trophic position of apex predator given that larger predators, such as lions (*Panthera leo*) and leopards, have been locally extirpated to enable sheep and goat production ([Van Sittert 1998](#); [Skead 2007, 2011](#)). Both jackals and caracals cause substantial impacts to small livestock species and are subject to widespread poisoning, trapping and shooting ([Bergman et al. 2013](#); [Minnie et al. 2016b](#); [Drouilly et al. 2018a](#); [du Plessis et al. 2018](#); [Minnie et al. 2018](#)). Warthogs damage fencing infrastructure (facilitating the dispersal of jackals; [Minnie et al. 2018](#)), exhibit other nuisance behaviours ([Mason 1982](#)), and have been introduced to many areas outside their historical range ([Somers 1992](#); [Skead 2007, 2011](#)). Baboons are frequently implicated in livestock predation, crop-raiding activities, and also perform a variety of other nuisance behaviours ([Minnie 2009](#); [Hoffman and O'Riain 2012](#); [Drouilly et al. 2018b](#); [Somers et al. 2018](#)). Big cats, such as leopards, are known to kill, consume and illicit fear in these wildlife species ([Hayward et al. 2012](#); [Clements et al. 2014](#)). Leopards present on livestock farms might therefore be expected to act as biocontrol tools to reduce predation losses experienced by livestock producers ([Minnie et al. 2015](#)). Permitting large predators to reside on livestock farms is opposed by many livestock producers who seek to prevent or mitigate all sources of livestock predation. Some have proposed that livestock guardian dogs (such as Anatolian shepherd or Kangal dogs) be deployed to protect livestock from predators ([Potgieter et al. 2013](#); [du Plessis et al. 2018](#)); they envisage a similar scenario where large carnivores can control smaller predators, guardian dogs protect livestock from large carnivores, and humans do not need to engage in lethal wildlife control. This approach has been described by proponents as an “effective”, “non-lethal”, “cost-effective”, “selective”, “useful, practical, and economically feasible” way to achieve “improved animal welfare and reduced non-target casualties” ([Marker et al. 2005](#); [Potgieter et al. 2013](#); [Rust et al. 2013](#); [McManus et al. 2014](#)).

The Australian and South African systems we describe are similar, in that they both reflect multi-predator and multi-prey systems where predators are expected to negatively influence prey through consumptive (direct, kill) and non-consumptive (indirect, fear) mechanisms (e.g. [Glen and Dickman 2005](#); [Letnic et al. 2012](#); [Valeix et al. 2012](#); [Riginos 2015](#)). The two systems we describe are also different, in that the Australian predator-prey guild is largely comprised of a depauperate assemblage of relatively small-sized, invasive and exotic species introduced since the European settlement of Australia in the late 1700s, whereas, the South African predator-prey guild is largely comprised of co-evolved native species of all sizes. We assess these two systems to explore the welfare impacts that arise for different potential applications of vertebrate biocontrols in different multi-predator multi-prey systems. Our assessment provides insights into the welfare outcomes of using large carnivores as biocontrol tools; we do not assess their efficacy, cost-effectiveness, sustainability or utility.

Animal welfare assessment

We assessed the animal welfare impacts of using dingoes, leopards and guardian dogs as biocontrol tools using the ‘Five Domains’ approach developed by Mellor and Reid (1994). This approach has since been widely utilised to assess animal welfare impacts imposed by different wildlife control tools in a variety of contexts, including European rabbit control in the United Kingdom (Baker et al. 2016) and brushtail possum (*Trichosurus vulpecula*) poisons in New Zealand (Beausoleil et al. 2016). It has also been adopted in Australia for assessing the control tools used against multiple invasive carnivore and herbivore species (RSPCA 2010; Sharp and Saunders 2011), where codes of practice and standard operating procedures have been further developed for a wide variety of particular applications (available at www.pestsmart.org.au). Using this approach, both lethal and non-lethal animal control tools are given an overall ‘humaneness score’ that enables their animal welfare impacts to be compared to other potential tools. As a result, for example, ground-shooting a feral goat in the head (rating = 3A) is considered to produce superior welfare outcomes to aerial-shooting (rating = 4C), which are both superior to mustering, holding, drafting, transporting, and eventually slaughtering the goat in an abattoir (e.g. Fig. 1; Sharp and Saunders 2011). This welfare assessment approach does not intend to prescribe which tool should be used by wildlife managers or declare a given tool ‘humane’ or ‘inhumane’, but rather makes the animal welfare impacts associated with each tool quantifiable and explicit. This allows comparison of different tools and techniques so that inferred animal welfare impacts can be ranked and considered along with efficacy, efficiency and other practical criteria when determining the most appropriate animal control tool for a given situation.




Fig. 1 – The relative humaneness of a variety of control tools used against (A) European red foxes and (B) feral goats in Australia. Adapted from Appendix 12 in Sharp and Saunders 2011, and used here to illustrate the outcomes of the five domains animal welfare impact assessment process.

The full assessment approach has been described in detail previously (e.g. [Mellor and Reid 1994](#); [Sharp and Saunders 2011](#); [Beausoleil and Mellor 2015](#); [Mellor and Beausoleil 2015](#); [Baker et al. 2016](#)). In summary, the assessment consists of two parts (A and B) that are collectively rated on a scale of 1–8, with another two welfare-related factors considered for each part (duration of suffering, and level of suffering) and collectively rated A–H (Fig. 2). Part A assesses the welfare of the controlled, individual animal before death. Part B assesses the mode of death. For each part, impacts are assessed within five domains used to determine an overall score: domain 1 is food or water deprivation, domain 2 is environmental challenge, domain 3 is disease/injury/functional impairment, domain 4 is behavioural restriction, and domain 5 is anxiety/fear/distress. The duration of suffering is assessed in categories such as immediate to seconds, minutes, hours, days, or longer. The level or intensity of suffering (which considers pain or discomfort, fear and distress) is assessed in categories such as mild, moderate, severe, or extreme. In other words, Part A assesses pre-death welfare and is scored 1–8 (with 1 indicating least and 8 indicating most suffering), Part B assesses the mode of death and is scored A–H (with A indicating least and H indicating most suffering), and judgments of suffering account for both the duration (in units of time) and the level of suffering (along a gradient of severity). This assessment approach produces overall scores such as 2B, 4D, 6F, and so on (Fig. 1), with a score of 2A reflecting a tool that produces superior welfare outcomes to a tool with a score of 3C, for example.

Part A	Duration of the chase				
	<60 second s	1–60 minute s	1–24 hours	1–7 days	>7 days
Extreme	5	6	7	8	8
Severe	4	5	6	7	8
Moderate	3	4	5	6	7
Mild	2	3	4	5	6
No impact	1	1	1	1	1

Part B	Duration of the kill				
	<60 second s	1–60 minute s	1–24 hours	1–7 days	>7 days
Extreme	E	F	G	H	H
Severe	D	E	F	G	H
Moderate	C	D	E	F	G
Mild	B	C	D	E	F
No impact	A	A	A	A	A

Fig. 2 – The overall humaneness scoring matrices for (A) Part A, the welfare impact associated with ‘the chase’ or the period preceding death, and (B) Part B, the welfare impact associated with ‘the kill’ or mode of death (adapted from [Sharp and Saunders 2011](#), where detailed definitions can be found explaining the mild, moderate, severe and extreme categories).

Table 1 – Overview of the subject-matter experience of the expert panel (author group) performing the humaneness assessment.

Author / assessor	Years of experience	Dingo ecology	Leopard ecology	Guardian dog ecology	Use of traditional wildlife control tools	Vertebrate biocontrol / predator introductions	Welfare assessments of wildlife control tools*	Policy planning, review & implementation	Conservation research & management	Animal welfare research & management	Pest animal research & management	Publications in Scopus	Link to Scopus author profile
Author 1	14	x		x	x	x	x	x	x	x	x	48	< INSERT LINK >
Author 2	38	x		x	x	x	x	x	x	x	x	30	< INSERT LINK >
Author 3	12	x		x	x	x	x	x	x	x	x	38	< INSERT LINK >
Author 4	8		x	x	x			x	x		x	6	< INSERT LINK >
Author 5	35	x		x	x	x	x	x	x	x	x	75	< INSERT LINK >
Author 6	16				x		x	x	x	x	x	34	< INSERT LINK >
Author 7	15	x	x		x	x			x		x	94	< INSERT LINK >
Author 8	38		x	x	x	x		x	x		x	198	< INSERT LINK >
Author 9	30	x			x	x	x	x	x	x	x	47	< INSERT LINK >
Author 10	10		x		x				x		x	7	< INSERT LINK >
Author 11	15		x	x	x		x	x	x	x	x	63	< INSERT LINK >
Author 12	11		x	x	x	x	x	x	x			52	< INSERT LINK >
Author 13	21		x			x		x	x			92	< INSERT LINK >

*Not including the current assessment.

The scores should ideally be derived by an experienced and diverse panel of participants considered to be expert in their subject-matter ([Sharp and Saunders 2011](#)). Baker et al. ([2016](#)) used only one author to perform the assessment and Beausoleil et al. ([2016](#)) used a panel of six scientists with various levels of expertise in animal welfare science, pest control and veterinary toxicology. In our assessment, we used a large, relevantly experienced and diverse author group who have collectively contributed to 784 published articles during our combined 263 years of experience (Table 1) at the time of writing. When performing our assessment, all 13 authors thoroughly discussed all overall scores before arriving at a unanimous consensus, which we report below. Despite our efforts, we acknowledge that because the five domains approach uses expert panels to make the assessment, it suffers from the problems of subjectivity inherent to the use of expert opinion, including how panel members are chosen. We attempted to minimise this issue by engaging participants with diverse backgrounds, but the conclusions we reach through use of this model are ultimately qualitative in nature ([Mellor and Beausoleil 2015](#)).

Dingoes, leopards and guardian dogs have not previously been assessed as control tools of wildlife ([Allen et al. 2017](#)). Hence, we assessed dingoes and guardian dogs as control tools of European red foxes, feral cats, feral goats, small mammals (such as house mice and black rats), kangaroos, feral pigs, and rabbits in Australia. We also assessed leopards and guardian dogs as control tools of jackals, caracals, cheetahs (*Acinonyx jubatus*), baboons, warthogs, fallow deer (*Dama dama*), Himalayan tahr, feral pigs and Cape foxes (*Vulpes chama*) in South Africa. The choice of prey species assessed for each predator was intended to reflect those that some people have argued need to be controlled (because of conflict with humans) and are potentially controllable by these predators. For convenience, in the text we collectively refer to all animals attacked or killed as 'prey', regardless of their functional role (carnivore or herbivore etc.) or whether or not they are consumed. We also refer to all those animals that do the attacking or killing as 'predators', regardless of their trophic position (top-predator, mesopredator, etc.) and whether or not their attacks successfully deter or kill prey.

Results

Performing the assessment

Though prey might technically be deprived of food or have their behaviour restricted during a predation event, we considered there to be 'no impact' in Domains 1, 2 and 4, and we do not discuss these further. We considered Domains 3 (e.g. injury, functional impairment) and 5 (e.g. anxiety, fear, pain and distress) to be most applicable to the mode of action (predation or fear of predation) in our case. We discussed and scored each predator against each prey over a period of several weeks using the assessment approach described above. During this process, the general paucity of empirical studies measuring the animal welfare impacts experienced during predation led to some personal variation in initial scores, with different values being attributed to the level of suffering authors perceived for individual prey species. In other words, some authors initially considered a given prey to be experiencing only mild suffering, whereas others felt that same prey might be experiencing extreme suffering. After considering these initial individual views, group discussion led to unanimous agreement that once a prey animal becomes aware that a predator is about to kill it, that prey animal can reasonably be assumed to be experiencing extreme suffering in those moments preceding capture; and that once the predator captures and proceeds to kill the prey, that prey

animal can reasonably be assumed to be experiencing extreme suffering in those moments prior to insensibility or death. Thus, scores for Domain 3 and Domain 5 were considered to be extreme in all cases.

Dingoes in Australia

The minimum time taken by dingoes to chase prey may be <60 seconds, and the minimum time taken to kill prey may also be <60 seconds. Assuming that prey experience extreme harm during both the chase and kill components of the process, the minimum score that any prey species can receive is 5E. This was the case only for small mammals, such as mice and rats (Fig. 3). Given that chases may last minutes (but unlikely hours) and time-to-death may also last minutes (but unlikely hours; e.g. [Behrendorff 2018](#)), scores for all other animals produce ranges, from 5E – 6E for feral cats and European rabbits, from 5E – 6F for red foxes and kangaroos, and 5E – 6H for feral goats and feral pigs (Fig. 3). The higher maximum scores for these prey reflect the longer chasing and handling times that might be expected for larger individuals of these prey species.

Guardian dogs in Australia

The minimum time taken by guardian dogs to chase Australian prey may also be <60 seconds, and the minimum time taken to kill prey may also be <60 seconds. Assuming that prey experience extreme harm during both the chase and kill components of the assessment, the minimum score that any prey species can receive is also 5E. Given the way guardian dogs function in relation to prey species ([Allen et al. 2016](#)), scores for all animals produce ranges, from 5E – 6F for feral cats and red foxes, from 5E – 6H for kangaroos, from 5F – 6H for feral goats and feral pigs, and 5F – 7H for dingoes (Fig. 3). The higher maximum scores for dingoes reflect the longer time it would normally take for guardian dogs to chase, subdue and kill dingoes.

Leopards in South Africa

The minimum time taken by leopards to chase prey may be <60 seconds, and the minimum time taken to kill prey may also be <60 seconds. Assuming that prey experience extreme harm during both the chase and kill components of the process, the minimum score that any prey species can receive is still 5E. Given the way leopards ambush and kill their prey ([Hubel et al. 2018](#)), scores for all animals produce ranges, from 5E – 5F for Cape foxes, jackals, warthogs, fallow deer, Himalayan tahr, and feral pigs, and from 5E – 6F for baboons and caracals (Fig. 3). The higher maximum scores for baboons and caracals reflect the longer time it might take for leopards to chase, subdue and kill these prey given their similar morphology (caracals) or use of group defence strategies (baboons).

Guardian dogs in South Africa

Like guardian dogs in Australia, the minimum time taken by guardian dogs to chase South African prey may also be <60 seconds, and the minimum time taken to kill prey may also be <60 seconds. Assuming that prey experience extreme harm during both the chase and kill components of the assessment, the minimum score that any prey species can receive is also 5E. Guardian dogs in South Africa produce ranges of scores from 5E – 6F for Cape foxes, jackals, warthogs and caracals, and from 5F – 6H for cheetahs and baboons (Fig. 3). The higher maximum scores for these two species reflect the longer expected time it might take for guardian dogs to chase, subdue and kill these dangerous prey. As an aside, deployment of guardian dogs is not without risk to the guardian dogs,

which are also killed by dingoes and other wild predators (e.g. [Allen et al. 2016](#)), producing similar harms to the guardian dog.




Fig. 3 – The relative humaneness of using dingoes in Australia (top left), guardian dogs in Australia (top right; lines used instead of circles for clarity), leopards in South Africa (bottom left), and guardian dogs in South Africa (bottom right) as vertebrate biocontrol tools against a variety of wildlife species that come into conflict with humans. Humaneness scores for applications shown in the bottom-left of each panel are considered less harmful than scores in the top-right of each panel.

Discussion

Large carnivores and guardian dogs are increasingly being recognised for the ecosystem services they can provide (e.g. [Linnell and Lescureux 2015](#); [van Bommel and Johnson 2016](#); [Gilbert et al. 2017](#); [O'Bryan et al. 2018](#)), including their ability to act as vertebrate biocontrol tools through consumptive and non-consumptive mechanisms (e.g. [Allen 2015](#); [Potgieter et al. 2016](#); [Williams et al. 2017](#); [Thinley et al. 2018](#)). Accordingly, predators are now being deliberately used or recommended as management tools for reducing the distribution, abundance and impacts of a variety of prey species, including both carnivores and herbivores (e.g. [Allen et al. 1998](#); [Minnie et al. 2015](#); [Atkins et al. 2017](#)). While much attention has been given to the animal welfare impacts associated with the use of traditional tools and techniques like poisoning, trapping or shooting ([Sharp and Saunders 2011](#); [Littin et al. 2014](#)), almost no attention has been given to the animal welfare impacts associated with 'natural tools', such as infectious biocontrols ([Hampton and Hyndman In press](#)) or the intentional use of large carnivores and guardian dogs as vertebrate biocontrol tools ([Allen et al. 2017](#)). Stakeholders have polarized views regarding such use of large carnivores. Some claim that deployment of large carnivores represents an "ethical", "humane" and "virtuous" approach (e.g. [Wallach 2014](#); [Johnson and Wallach 2016](#)), while others claim that it represents a "cruel", "wrong", "immoral" and non-compassionate approach (e.g. [Schwartz 2016](#); [Wallach et al. 2018](#)). This becomes particularly confusing when individuals draw opposite ethical conclusions depending on the predator-prey combination being discussed (e.g. [Wallach 2014](#) advocates using dingoes to control foxes and feral cats, but later denigrates others' use of dingoes to control feral goats; [Wallach et al. 2018](#)). With few empirical studies to draw from, the results of our formal assessment of using large carnivores and guardian dogs as biocontrol tools indicates that their animal welfare impacts vary depending on the large carnivore involved, the prey species they are intended to control, and the way in which the carnivores interact with those animals (Fig. 3). In general, the consumptive and non-consumptive mechanisms used by large carnivores and guardian dogs cause considerable animal welfare impacts to the individual animals they are intended to control, and this harm is typically rated greater than those harms caused by most traditional techniques (see [Sharp and Saunders 2011](#) for details).

Using dingoes to control red foxes, for example, rates as the most harmful tool of all those that are currently in use or proposed for use against red foxes. Trapping red foxes with padded leg-hold traps is rated at 6B, or the most harmful of all current red fox control tools (Fig. 1; [Sharp and Saunders 2011](#)). This score conservatively accounts for the worst instances in which foxes remain in traps for up to 24 hours and chew their own toes or feet ([Meek et al. 1995](#); [Fleming et al. 1998](#)). Dingoes are cursorial predators that 'bite and shake' their prey to death ([Corbett 2001](#); [Behrendorff et al. 2018](#)). Using dingoes to chase, attack and kill red foxes was conservatively scored as 5E – 6F depending on how long red foxes are chased and how long it takes dingoes to kill them (Fig. 3). Even at best – 5E, which represents extreme harm being experienced in a chase that lasts <60 seconds, and extreme harm being experienced in a kill that lasts <60 seconds – using dingoes against red foxes is rated as being more harmful to the red fox than using padded leg-hold traps (compare Fig. 1 with Fig. 3). At worst – 6F, which represents extreme harm being experienced in a chase that lasts >60 seconds, and extreme harm being experienced in a kill that lasts >60 seconds – using dingoes against red foxes substantially exceeds the suffering experienced by using firearms, traps, poisons (both sodium fluoroacetate or 'Compound 1080' and para-aminopropiophenone or 'PAPP'), or any other red fox control tool. This is similarly the case for using dingoes against feral cats (Fig. 3).

Using leopards to control various wildlife in South Africa is rated as 5E – 6F depending on the target species and the time taken by leopards to catch and kill the prey (Fig. 3). Leopards are ambush predators that stalk and approach prey unseen prior to initiating a short chase or a single pounce ([Hubel et al. 2018](#)), and as such, leopard prey are likely to experience extreme harm in a chase that lasts <60 seconds. Furthermore, leopards employ a skull bite to kill smaller prey and a throat or nape bite for larger prey ([Brain 1981](#)), which will result in prey experiencing extreme harm in a kill that lasts <60 seconds. This type of ‘stalk and pounce’ predatory behaviour produces lower humaneness scores (such as 5E; Fig. 3) than ‘chase and catch’ types of predatory behaviour. However, baboons and caracals are formidable prey with teeth and claws which can be used in defence against leopard attack ([Brain 1981](#); [Jooste et al. 2012](#)). This is likely to increase capture and handling time and thus a time-to-death of >60 seconds, resulting in a rating closer to 6F (Fig. 3). No comparative data exists to evaluate our results against other commonly used wildlife control tools in South Africa, such as ground shooting, cage trapping, leg-hold traps (or gin traps) and poisoning ([Minnie 2009](#); [du Plessis et al. 2018](#)). However, one may assume that the animal welfare impacts of similar management tools for similar species (i.e. jackals and red foxes, and caracal and feral cats) would be comparable between South Africa and Australia. Thus, we can assume that the animal welfare impacts of a jackal caught in a leg-hold trap in South Africa would be similar to that of a red fox caught in a leg-hold trap in Australia, which was rated at 6B (Fig. 1). Similarly, we can assume that the animal welfare impacts of shooting a caracal in a cage in South Africa would be similar to that of a feral cat shot in a cage in Australia, which was rated at 4B – 5B ([Sharp and Saunders 2011](#)). Thus, even in a best-case scenario (rating = 5E; Fig. 3), using leopards against jackals, caracals, and other prey species would be rated as being more harmful to jackals and caracals than using leg-hold traps, ground shooting, or poisoning.

Guardian dogs have been used as biocontrol tools for centuries, and can be very effective at reducing livestock predation in some contexts ([Coppinger and Coppinger 1993](#); [van Bommel 2010](#); [Potgieter et al. 2013](#); [Linnell and Lescureux 2015](#)). They typically work by defensively guarding livestock, frightening potential predators away and seldom seeking-out or hunting-down predators and other wildlife (e.g. [Allen et al. 2016](#)). They are usually thought of as a non-lethal tool, intended to non-consumptively instil fear and repel predators ([van Bommel and Johnson 2014](#)). However, guardian dogs are known to consumptively kill target and non-target animals ([van Bommel 2010](#); [Potgieter et al. 2016](#)), and should thus be classified as both a non-lethal (intentional harm) and lethal (unintentional harm) tool. When such lethal effects occur, the associated harms to the prey are relatively straightforward to assess and are rated similarly to other large carnivores (Fig. 3). An exception to this occurs when a guardian dog is involved in a fight that is not immediately fatal to the target animal, but where death is delayed and the animal dies as a consequence of its injuries (e.g. by secondary infection or starvation) sometime later. This scenario produced the higher limits of the range of some humaneness scores for dingoes and guardian dogs (Fig. 3), where extreme harm can be experienced over a time-to-death that can last longer than 24 hours. Though extreme, at their worst these outcomes are still similar to other natural predation behaviours exhibited by other wild-living large carnivores. For example, the restraining and killing of elephants (*Loxodonta africana*) by lions (*Panthera leo*), which can last longer than 24 hours ([Power and Compion 2009](#)), would be scored at a maximum rating of 6H were it to be considered anthropogenic predation worthy of scrutiny. Likewise, the intentional biting, envenomation and eventual (days later) death of water buffalo (*Bubalus bubalis*) by Komodo dragons (*Varanus komodoensis*; [Auffenberg 1981](#); [Fry et al. 2009](#); [Bull et al. 2010](#)) would be rated at 8F – 6H, depending on when the chase is considered to

stop and the kill is considered to start. Deploying large carnivores as biocontrol tools essentially represents a management decision to introduce considerable consumptive and non-consumptive effects into food webs as a means of mitigating the undesirable impacts of target wildlife. In almost all cases, however, these ‘natural’ biocontrols will impose greater animal welfare impacts to target animals than any other anthropogenic tool used by managers, which typically have no chase period (e.g. poisoning), an immediate or very short kill period (e.g. aerial shooting), or both (e.g. ground shooting, cyanide poisoning; [Sharp and Saunders 2011](#)).

Essential for assessing the animal welfare impacts of using large carnivores and guardian dogs as biocontrol tools using the five domains approach is an understanding of how carnivores hunt and catch their prey, or their fine-scale interactions with other predator and prey species. Despite a rich body of such literature (e.g. [Barbosa and Castellanos 2005](#); [Schmitz 2008](#); [Thaker et al. 2011](#); [Belgrad and Griffen 2016](#); [Moll et al. 2016](#); [Allen et al. 2018](#)), patterns of predation behaviour are rarely simple ([MacNulty et al. 2007](#)) and relatively few studies quantify the animal welfare outcomes experienced by prey during predation (but see [Creel and Christianson 2008](#); [Kluever et al. 2008](#); [Creel et al. 2009](#); [Behrendorff et al. 2018](#)). Thus, in making assessments such as ours (Fig. 3), we are reliant on expert opinion and will almost certainly be working with few relevant empirical data. Moreover, all qualitative index approaches for assessing animal welfare impacts, such as the one we have used, have limitations in predicting the actual welfare outcomes for animals ([Beausoleil and Mellor 2015](#)). “Because of the dearth of objective data relating to welfare in this particular field, some judgements will have to be made subjectively” ([Sharp and Saunders 2011](#); pg. 40). Making assumptions is unavoidable. Quantifying welfare impacts requires physiological measurements of stress and injury responses over time for both the chase (Part A) and the kill (Part B) phases. These data must then be compared with similar measurements for the other methods (e.g. traditional tools) that form the basis of the comparisons of animal welfare impacts. Given this, our assessment (1) represents testable predictions based on current (lack of) literature and expert opinion, (2) represents a valuable starting point which stimulates discussion on the animal welfare implications of using large carnivores and guardian dogs to control other species, and (3) highlights the type of empirical information required to repeat our assessment and advance our understanding of the harms associated with these biocontrol tools.

In our assessment we assumed that in most (if not all) individual predator-prey altercations the prey experiences an extreme level of suffering during both the chase and kill components considered in the assessment process. We also assumed that prey are aware of their impending predation. For example, when a jackal becomes aware that a leopard is chasing it with intent to kill it, the jackal is probably experiencing some of the following criteria from Domain 5, described by Sharp and Saunders ([2011](#); pg. 48) viz., “extreme inescapable or unrelieved anxiety, fear, pain, sickness, breathlessness, nausea, lethargy/weakness, dizziness, unsatisfied thirst and/or hunger or [some] other negative affective experience causing distress which is judged to be at or beyond the limits of reasonable endurance and results in the death of the animal”. Any re-assessment of our results (Fig. 3) – by any other assessment panel, either with or without empirical data – cannot yield lower animal welfare scores than what we report unless (A) it is assumed or can be shown that prey are unaware they are being chased or killed or (B) prey experience less-than-extreme welfare harms when they are knowingly being chased and killed. It might reasonably be argued that in some individual predation events (particularly of leopards against smaller prey), prey may be unaware of their impending predation and/or may experience a relatively instantaneous death. If, when or

where this is true, then the humaneness scores we report will be overestimated. We considered assuming prey unawareness or less-than-extreme suffering as a minimum in Domains 3 and 5, but felt more confident in assuming that such events would be the exception rather than the rule. If prey are aware and the intensity of suffering is conservatively assumed to be extreme (as we have done), then the outcomes of assessing large carnivores as biocontrol tools largely depend on the duration of harm, which will vary in different predator-prey combinations. For example, the animal welfare impacts of guardian dogs on jackals (5E – 6F) are relatively more harmful than those of leopards on jackals (5E – 5F) given that guardian dogs would typically rely on a prolonged chase to subdue prey while leopards are ambush predators which either pounce or engage in a very brief chase. Furthermore, canids typically kill their prey with a bite-and-shake movement (e.g. [Behrendorff et al. 2018](#)) rather than the swift and crushing skull or neck bite used by leopards (e.g. [Brain 1981](#)). Regardless of the particular predator-prey combination, short chases and quick kills will be rated less harmful than long chases and protracted kills.

In contrast to the lethal, consumptive outcomes of interactions between large carnivore biocontrol tools and prey (discussed above), their non-lethal, non-consumptive harms are much more difficult to assess reliably. If large carnivore and guardian dog biocontrol tools are successful in establishing a landscape of fear (*sensu* [Brown et al. 1999](#)) and animals suffer as a result of that fear, then harm is caused. Large carnivores and guardian dogs are obviously intended to work in this way (e.g. [Linnell and Lescureux 2015](#); [van Bommel and Johnson 2015](#)); the challenge lies in determining just how harmful these effects are for the target and non-target wildlife they are deployed against, and how these harms compare to other wildlife management tools.

The duration of harm component of the assessment process is relatively easy to complete given knowledge of large carnivore contact times with other species or the time they spend in close proximity to each other. This will typically fall in the range of seconds to minutes (e.g. [Behrendorff 2018](#)), but may sometimes extend to hours ([Barbosa and Castellanos 2005](#); [Allen et al. 2016](#)). However, these interactions are repeated over time and may represent ‘cumulative effects’ or ‘compounded welfare impacts’ ([Sharp and Saunders 2011](#)), the magnitude of which is determined in large part by their frequency. What is difficult to assess is how harmful these cumulative welfare effects are to prey and whether these are more or less harmful than those of other tools. Several difficult questions arise. For example, is dying from poison over several hours a better welfare outcome than being repeatedly chased by a predator over many months or years, or being displaced and forced into starvation or conflict with conspecifics? Or, should the harm be assumed less for co-evolved predators and prey (e.g. leopards and jackals) than those species that have only come into contact through anthropogenic activities (e.g. leopards and guardian dogs)? Prey species frequently come in close proximity to their predators in multi-predator multi-prey systems. Some species (e.g. kleptoparasitic scavengers) even follow their dominant predators around, hoping to scavenge a meal from them ([Iyengar 2008](#); [Cusack et al. 2017](#)). Species that coexist in this way likely evolved behavioural and physiological adaptations that allow them to tolerate – to some degree – these non-consumptive, cumulative harms without debilitating physiological effects. On the other hand, prey can suffer greatly from non-lethal interactions, with the landscape of fear affecting their movement patterns, sociality, foraging, reproduction, fitness and survival ([Kluever et al. 2008](#); [Creel et al. 2009](#); [Clinchy et al. 2013](#)). Proactive anti-predator behaviours typically carry food-mediated costs, reactive anti-predator behaviours typically carry stress-mediated costs, and both of these costs can be manifest as reduced fitness, fecundity and survival ([Creel 2018](#)). Thus, there is likely to be a strong

positive relationship between the strength of a landscape of fear and harmful animal welfare impacts experienced by the prey (Fig. 4), and one cannot argue that a strong landscape of fear exists while at the same time arguing that the prey are not experiencing any harm from that fear. The cumulative welfare impacts that arise from repeated non-consumptive predator-prey interactions will always last longer than the effects of any lethal anthropogenic wildlife control tool. Thus, 'non-lethal' does not necessarily equate to 'least harmful'. Though challenging to quantify and rank, the potential for large carnivores and guardian dogs to cause substantial cumulative harms to both target and non-target animals cannot be disregarded and requires further inquiry.




Fig. 4 – The predicted relationship between the strength of the landscape of fear and the welfare harms experienced by individual animals.

Conclusion and recommendations

Large carnivore and guardian dog biocontrol tools act through both lethal (consumptive) and non-lethal (non-consumptive) mechanisms. When intentionally deployed by humans for wildlife management purposes, the harms they impose on other animals could be considered anthropogenic in origin and require ethical scrutiny. The passive or active use of such anthropogenic predation may offer important benefits to wildlife management and conservation, but, like all other tools, they cause harm to target and non-target animals. The animal welfare impacts of large carnivores and guardian dogs as biocontrol tools appear similar across different predator-prey systems, but are highly dependent on specific predator-prey combinations. In most cases where they can be compared to other tools, our results indicate that predation and/or fear of predation produces more harm to target animals than most other alternative 'human tools' assessed. The intentional lethal and non-lethal harms caused by introducing somewhat uncontrollable consumptive and non-consumptive mechanisms into food webs should not be ignored or dismissively assumed to be negligible. These findings complement knowledge of the harm caused by traditional wildlife management tools and have important implications for those considering rewilding programs or the use of large carnivores or guardian dogs as biocontrols of wildlife within multi-predator multi-prey systems.

Some have argued that common control techniques, such as the poisoning of red foxes with Compound 1080 or PAPP in Australian sheep production zones, should be prohibited or abandoned ostensibly on animal welfare grounds, and instead be replaced with the deployment of dingoes as biocontrol tools against red foxes and deployment of guardian dogs to protect sheep from dingoes (e.g. [Johnson and Wallach 2016](#); [Smith and Appleby 2018](#)). However, deliberately deploying carnivores for this purpose is undeniably anthropogenic and assessment of their animal welfare impacts suggests that this course of action represents abandonment of less-harmful tools in favour of more-harmful ones. Some have also argued that traditional forms of jackal control (such as ground shooting at night) should be discouraged because they impose greater animal welfare impacts than the deployment of guardian dogs ([du Plessis 2013](#); [McManus et al. 2014](#)). However, guardian dogs kill jackals directly and may also create a landscape of fear that displaces jackals, which together have been rated here as causing greater animal welfare impacts than ground shooting. How these harms compare to other tools is presently unclear given that formal animal welfare assessments have not been completed for other wildlife control tools used in South Africa. Once completed, such assessments may enable more informed discussion about the animal welfare impacts associated with different wildlife control tools used there (see [du Plessis et al. 2018](#)), but it should be noted that the five domains assessment approach does not generally consider indirect or unintentional harms ([Beausoleil and Mellor 2015](#)), which are also important ([Fraser and MacRae 2011](#)).

We predict that large carnivores and guardian dogs can be effective vertebrate biocontrol tools in some contexts (e.g. [Allen et al. 1998](#)), both despite and because of the consumptive and non-consumptive harms they cause to prey animals. A greater understanding of their utility as wildlife management tools would benefit from further demonstrations of their efficacy, advantages and disadvantages in different contexts, including explicit empirical assessment of their animal welfare impacts on prey. This information will play an increasingly important role in the future social license of using guardian animals specifically and rewilding with large carnivores more generally ([Hampton and Teh-White 2019](#)).

Framing our assessment was the view that when predators are introduced to an environment by humans, the impacts predators exert in that environment constitute anthropogenic effects in perpetuity. This logic is an extension of the ethic applied to the ecological impacts incurred by invasive species and the generally accepted moral responsibility of humans to mitigate these impacts in the field of conservation ([Russell et al. 2016](#)). However, we acknowledge that an alternative view may be that when humans reintroduce predation (less so for novel introductions) it restores certain previously occurring processes such that an ecological system can henceforth function autonomously (although, determining an acceptable historical benchmark creates its own problems; [Hayward 2012](#)). Under this premise, subsequent predation events would not be considered anthropogenic and would not be morally equivalent to other wildlife management tools used by humans. This complex question is further complicated by variation in what might be regarded as the naturalness of the predator and the environment ([Torres 2015](#)). For example, reintroducing wolves to Yellowstone may be argued to be highly natural given that wolves previously occurred there, were extirpated by anthropogenic processes, and were absent for a period of only decades. On the other end of the spectrum, the introduction, training and maintenance of domestic guardian dogs in an environment that never supported natural wild populations of dogs could only tenuously be considered natural.

The ethical consequences of this issue are profound for conservation and invasive species disciplines. If it is asserted that the harm imposed by all introduced and reintroduced predators deserve equal animal welfare consideration as other wildlife management tools, then predation and fear of predation are likely to be seen as an unappealing option for managers. Under this ethical interpretation, whereby anthropogenic predation constitutes an 'immoral' approach, the social license of rewilding programs and all predator reintroductions would be extremely fragile ([Allen et al. 2017](#)). Even the widely celebrated reintroduction of grey wolves to Yellowstone National Park could be argued to constitute an unacceptable anthropogenic harm imposed on the naïve prey animals of that ecosystem. In contrast, should anthropogenic predation not be considered anthropogenic at all (i.e. if humans are deemed to not be responsible for the predation, fear and distress arising from introducing predators), then concerns about the welfare effects of using large carnivores as vertebrate biocontrols are void, and managers may begin to deploy vertebrate biocontrols more frequently. In any event, we assert that the animal welfare impacts arising from deliberate use of large carnivores and guardian dogs requires more critical examination, and we invite discussion.

References

AJP. (2015). 'Animal Justice Party: Dingo policy.' (Animal Justice Party: Strawberry Hills, New South Wales, Australia)

Allen, B.L. (2015). More buck for less bang: reconciling competing wildlife management interests in agricultural food webs. *Food Webs* 2, 1-9.

Allen, B.L., Allen, L.R., Engeman, R.M, Leung, L.K.-P. (2014). Sympatric prey responses to lethal top-predator control: predator manipulation experiments. *Frontiers in Zoology* 11, 56.

Allen, B.L., Ballard, G., Fleming, P.J.S. (2017). *The humaneness of dingoes and guardian dogs as vertebrate biocontrol tools*. In 'Proceedings of the Australasian Wildlife Management Society Conference.' (Australasian Wildlife Management Society: Katoomba)

Allen, B.L., Fawcett, A., Anker, A., Engeman, R.M., Lisle, A., Leung, L.K.-P. (2018). Environmental effects are stronger than human effects on mammalian predator-prey relationships in arid Australian ecosystems. *Science of the Total Environment* 610-611C, 451-461.

Allen, B.L., Fleming, P.J.S. (2012). Reintroducing the dingo: the risk of dingo predation to threatened vertebrates of western New South Wales. *Wildlife Research* 39, 35-50.

Allen, B.L., Leung, K.-P. (2014). The (non)effects of lethal population control on the diet of Australian dingoes. *PLoS ONE* 9, e108251.

Allen, B.L., West, P. (2013). The influence of dingoes on sheep distribution in Australia. *Australian Veterinary Journal* 91, 261-267.

Allen, B.L., West, P. (2015). Dingoes are a major causal factor for the decline and distribution of sheep in Australia. *Australian Veterinary Journal* 93, 90-92.

Allen, B.L., Wishart, J., Staples, L. (In press). Para-aminopropiophenone (PAPP) in Canid Pest Ejectors (CPEs) kills dingoes and European red foxes quickly and humanely. *Environmental Science and Pollution Research* xx, xx-xx.

Allen, L., Lee, J., Gonzalez, A. (1998). 'The management and eradication of feral goats from Townshend Island. Final report to Department of Defence.' (Department of Natural Resources: Toowoomba)

Allen, L.R., Stewart-Moore, N., Byrne, D., Allen, B.L. (2016). Guardian dogs protect sheep by guarding sheep, not by establishing territories and excluding predators. *Animal Production Science* 57, 1118-1127.

Anon. (2014). 'National wild dog action plan: promoting and supporting community-driven action for landscape-scale wild dog management.' (WoolProducers Australia: Barton, Australian Capital Territory)

Atkins, A., Redpath, S.M., Little, R.M., Amar, A. (2017). Experimentally manipulating the landscape of fear to manage problem animals. *The Journal of Wildlife Management* 81, 610-616.

Atwood, T.C., Gese, E.M., Kunkel, K.E. (2009). Spatial partitioning of predation risk in a multiple predator–multiple prey system. *Journal of Wildlife Management* 73, 876-884.

Auffenberg, W. (1981). 'The behavioral ecology of the Komodo monitor.' (University Presses of Florida: Gainesville)

Baker, S.E., Sharp, T.M., Macdonald, D.W. (2016). Assessing animal welfare impacts in the management of European rabbits (*Oryctolagus cuniculus*), European moles (*Talpa europaea*) and carrion crows (*Corvus corone*). *PLOS ONE* 11, e0146298.

Barbosa, P., Castellanos, I. (2005). 'Ecology of predator-prey interactions.' (Oxford University Press: New York)

Barnes, T.S., Goldizen, A.W., Morton, J.M., Coleman, G.T. (2008). Cystic echinococcosis in a wild population of the brush-tailed rock-wallaby (*Petrogale penicillata*), a threatened macropodid. *Parasitology* 135, 715-723.

Beausoleil, N.J., Fisher, P., Littin, K.E., Warburton, B., Mellor, D.J., Dalefield, R.R., Cowan, P. (2016). A systematic approach to evaluating and ranking the relative animal welfare impacts of wildlife control methods: poisons used for lethal control of brushtail possums (*Trichosurus vulpecula*) in New Zealand. *Wildlife Research* 43, 553-565.

Beausoleil, N.J., Mellor, D.J. (2015). Advantages and limitations of the Five Domains model for assessing welfare impacts associated with vertebrate pest control. *New Zealand Veterinary Journal* 63, 37-43.

Behrendorff, L. (2018). Clever girl? An observation of innovative prey handling by a dingo (*Canis dingo*). *Pacific Conservation Biology* 24, 194-197.

Behrendorff, L., Belonje, G., Allen, B.L. (2018). Intraspecific killing behaviour of canids: how dingoes kill dingoes. *Ethology, Ecology and Evolution* 30, 88-98.

Belgrad, B.A., Griffen, B.D. (2016). Predator–prey interactions mediated by prey personality and predator hunting mode. *Proceedings of the Royal Society of London B: Biological Sciences* 283, 2016.0408.

Berger, K.M. (2006). Carnivore-livestock conflicts: effects of subsidized predator control and economic correlates on the sheep industry. *Conservation Biology* 20, 751-761.

Bergman, D.L., De Waal, H., Avenant, N.L., Bodenchuk, M., Marlow, M.C., Nolte, D.L. JB Armstrong, GR Gallagher (Eds) (2013) '*The need to address black-backed jackal and caracal predation in South Africa*, Proceedings of the 15th Wildlife Damage Management Conference.' Clemson, South Carolina.

Bleicher, S.S. (2017). The landscape of fear conceptual framework: definition and review of current applications and misuses. *PeerJ* 5, e3772.

Bothma, J.D.P. (1971). Control and ecology of the black-backed jackal *Canis mesomelas* in the Transvaal. *Zoologica Africana* 6, 187-193.

Brain, C.K. (1981). 'The hunters or the hunted? An introduction to African cave taphonomy.' (University of Chicago Press: Chicago)

Breck, S.W., Poessel, S.A., Bonnell, M.A. (2017). Evaluating lethal and nonlethal management options for urban coyotes. *Human-Wildlife Interactions* 11, 133-145.

Brown, J.S., Laundre, J.W., Gurung, M. (1999). The ecology of fear: optimal foraging, game theory, and trophic interactions. *Journal of Mammalogy* 80, 385-399.

Bull, J.J., Jessop, T.S., Whiteley, M. (2010). Deathly drool: evolutionary and ecological basis of septic bacteria in Komodo Dragon mouths. *PLOS ONE* 5, e11097.

Campbell, G., Coffey, A., Miller, H., Read, J.L., Brook, A., Fleming, P.J.S., Bird, P., Eldridge, S., Allen, B.L. (2019). Dingo baiting did not reduce foetal/calf losses in beef cattle in northern South Australia. *Animal Production Science* 59, 319-330.

Clements, H.S., Tambling, C.J., Hayward, M.W., Kerley, G.I.H. (2014). An objective approach to determining the weight ranges of prey preferred by and accessible to the five large African carnivores. *PLoS ONE* 9, e101054.

Clinchy, M., Sheriff, M.J., Zanette, L.Y. (2013). The ecology of stress: predator-induced stress and the ecology of fear. *Functional Ecology* 27, 56-65.

Coppinger, L., Coppinger, R. (1993). *Dogs for herding and guarding livestock*. In 'Livestock handling and transport.' Ed. T Grandin. pp. 179-196. (Inkata Press: Melbourne, Australia)

Corbett, L.K. (2001). 'The dingo in Australia and Asia.' (J.B. Books, South Australia: Marleston)

Creel, S. (2018). The control of risk hypothesis: reactive vs. proactive antipredator responses and stress-mediated vs. food-mediated costs of response. *Ecology Letters* 21, 947-956.

Creel, S., Christianson, D. (2008). Relationships between direct predation and risk effects. *Trends in Ecology and Evolution* 23, 194-201.

Creel, S., Winnie, Jr., J.A., Christianson, D. (2009). Glucocorticoid stress hormones and the effect of predation risk on elk reproduction. *Proceedings of the National Academy of Sciences of the USA* 106, 12,388-12,393.

Cusack, J.J., Dickman, A.J., Kalyahe, M., Rowcliffe, J.M., Carbone, C., Macdonald, D.W., Coulson, T. (2017). Revealing kleptoparasitic and predatory tendencies in an African mammal community using camera traps: a comparison of spatiotemporal approaches. *Oikos* 126, 812-822.

de Tores, P.J., Marlow, N.J. (2011). *A review of the relative merits of predator exclusion fencing and repeated 1080 fox baiting for protection of native fauna: five case studies from Western Australia*. In 'Fencing for conservation.' Eds MJ Somers, MW Hayward. pp. 21-42. (Springer: New York)

Doherty, T.S., Ritchie, E.G. (2017). Stop jumping the gun: a call for evidence-based invasive predator management. *Conservation Letters* 10, 15-22.

Drouilly, M., Nattrass, N., O'Riain, M.J. (2018a). Dietary niche relationships among predators on farmland and a protected area. *The Journal of Wildlife Management* 82, 507-518.

Drouilly, M., Tafani, M., Nattrass, N., O’Riain, J. (2018b). Spatial, temporal and attitudinal dimensions of conflict between predators and small-livestock farmers in the Central Karoo. *African Journal of Range & Forage Science* 35, 245-255.

du Plessis, J., Avenant, N., Botha, A., Mkhize, N., Muller, L., Mzileni, N., O’Riain, J., Parker, D., Potgieter, G., Richardson, P., Rode, S., Viljoen, N., Tafani, M. (2018). *Past and current management of predation on livestock*. In 'Livestock predation and its management in South Africa: a scientific assessment.' Eds GIH Kerley, SL Wilson, D Balfour. pp. 125-177. (Centre for African Conservation Ecology, Nelson Mandela University: Port Elizabeth, South Africa)

du Plessis, J.J. (2013). *Towards a sustainable management strategy for Canis mesomelas and Caracal caracal on rangeland*. PhD thesis, University of the Free State.

Dubois, S., Fenwick, N., Ryan, E.A., Baker, L., Baker, S.E., Beausoleil, N.J., Carter, S., Cartwright, B., Costa, F., Draper, C., Griffin, J., Grogan, A., Howald, G., Jones, B., Littin, K.E., Lombard, A.T., Mellor, D.J., Ramp, D., Schuppli, C.A., Fraser, D. (2017). International consensus principles for ethical wildlife control. *Conservation Biology* 31, 753-760.

Eason, C.T., Miller, A., MacMorran, D.B., Murphy, E.C. (2014). Toxicology and ecotoxicology of para-aminopropiophenone (PAPP) – a new predator control tool for stoats and feral cats in New Zealand. *New Zealand Journal of Ecology* 38, 177-188.

Eisenberg, C. (2011). 'The wolf's tooth: keystone predators, top predators, trophic cascades, and biodiversity.' (Island Press: Washington D.C.)

Eldridge, S.R., Shakeshaft, B.J., Nano, T.J. (2002). 'The impact of wild dog control on cattle, native and introduced herbivores and introduced predators in central Australia. Final report to the Bureau of Rural Sciences.' (Parks and Wildlife Commission of the Northern Territory: Alice Springs)

Estes, J., Crooks, K., Holt, R.D. (2013). *Ecological role of predators*. In 'Encyclopedia of biodiversity (Second edition).' Ed. SA Levin. Vol. 6 pp. 229-249. (Academic Press: Waltham)

Fitzgerald, G. (2009). 'Public attitudes to current and proposed forms of pest animal control.' (Invasive Animals Cooperative Research Centre: Canberra)

Fleming, P.J., Allen, L.R., Berghout, M.J., Meek, P.D., Pavlov, P.M., Stevens, P.L., Strong, K., Thompson, J.A., Thomson, P.C. (1998). The performance of wild-canid traps in Australia: efficiency, selectivity, and trap-related injuries. *Wildlife Research* 25, 327-338.

Fleming, P.J.S., Allen, B.L., Allen, L.R., Ballard, G., Bengsen, A.J., Gentle, M.N., McLeod, L.J., Meek, P.D., Saunders, G.R. (2014). *Management of wild canids in Australia: free-ranging dogs and red foxes*. In 'Carnivores of Australia: past, present and future.' Eds AS Glen, CR Dickman. pp. 105-149. (CSIRO Publishing: Collingwood)

Fleming, P.J.S., Allen, B.L., Ballard, G. (2012). Seven considerations about dingoes as biodiversity engineers: the socioecological niches of dogs in Australia. *Australian Mammalogy* 34, 119-131.

Fox, M.W. (1969). Ontogeny of prey-killing behavior in Canidae. *Behaviour* 35, 259-272.

Fraser, D., MacRae, A.M. (2011). Four types of activities that affect animals: implications for animal welfare science and animal ethics philosophy. *Animal Welfare* 20, 581-590.

Fry, B.G., Wroe, S., Teeuwisse, W., van Osch, M.J.P., Moreno, K., Ingle, J., McHenry, C., Ferrara, T., Clausen, P., Scheib, H., Winter, K.L., Greisman, L., Roelants, K., van der Weerd, L., Clemente, C.J., Giannakis, E., Hodgson, W.C., Luz, S., Martelli, P., Krishnasamy, K., Kochva, E., Kwok, H.F., Scanlon, D., Karas, J., Citron, D.M., Goldstein, E.J.C., Mcnaughtan, J.E., Norman, J.A. (2009). A central role for venom in predation by *Varanus komodoensis* (Komodo Dragon) and the extinct giant *Varanus (Megalania) priscus*. *Proceedings of the National Academy of Sciences* 106, 8969-8974.

Gilbert, S.L., Sivy, K.J., Pozzanghera, C.B., DuBour, A., Overduijn, K., Smith, M.M., Zhou, J., Little, J.M., Prugh, L.R. (2017). Socioeconomic benefits of large carnivore recolonization through reduced wildlife-vehicle collisions. *Conservation Letters* 10, 431-439.

Glen, A.S., Dickman, C.R. (2005). Complex interactions among mammalian carnivores in Australia, and their implications for wildlife management. *Biological Reviews* 80, 387-401.

Hairston, N., Smith, F., Slobodkin, L. (1960). Community structure, population control and competition. *American Naturalist* 94, 421-425.

Hampton, J.O., Hyndman, T.H. (In press). Underaddressed animal-welfare issues in conservation. *Conservation Biology* xx, xx-xx.

Hampton, J.O., Hyndman, T.H., Laurence, M., Perry, A.L., Adams, P., Collins, T. (2016). Animal welfare and the use of procedural documents: limitations and refinement. *Wildlife Research* 43, 599-603.

Hampton, J.O., Teh-White, K. (2019). Animal welfare, social license, and wildlife use industries. *The Journal of Wildlife Management* 83, 12-21.

Harrington, L.A., Moehrensclager, A., Gelling, M., Atkinson, R.P.D., Hughes, J., Macdonald, D.W. (2013). Conflicting and complementary ethics of animal welfare considerations in reintroductions. *Conservation Biology* 27, 486-500.

Hayward, M. (2012). Time to agree on a conservation benchmark for Australia. *Pacific Conservation Biology* 18, 69-76.

Hayward, M.W., Jędrzejewski, W., Jędrzejewska, B. (2012). Prey preferences of the tiger *Panthera tigris*. *Journal of Zoology* 286, 221-231.

Hoffman, T.S., O'Riain, M.J. (2012). Monkey management: using spatial ecology to understand the extent and severity of human-baboon conflict in the Cape Peninsula, South Africa. *Ecology and Society* 17, 13-29.

Holt, R.D., Lawton, J.H. (1994). The ecological consequences of shared natural enemies. *Annual Review of Ecology and Systematics* 25, 495-520.

Horta, O. (2017). Animal suffering in nature: the case for intervention. *Environmental Ethics* 39, 261-279.

Hubel, T.Y., Golabek, K.A., Rafiq, K., McNutt, J.W., Wilson, A.M. (2018). Movement patterns and athletic performance of leopards in the Okavango Delta. *Proceedings of the Royal Society B* 285, 20172622.

Iyengar, E.V. (2008). Kleptoparasitic interactions throughout the animal kingdom and a re-evaluation, based on participant mobility, of the conditions promoting the evolution of kleptoparasitism. *Biological Journal of the Linnean Society* 93, 745-762.

Johnson, C.J., Wallach, A.D. (2016). The virtuous circle: predator-friendly farming and ecological restoration in Australia. *Restoration Ecology* 24, 821-826.

Jooste, E., Pitman, R.T., van Hoven, W., Swanepoel, L.H. (2012). Unusually high predation on chacma baboons (*Papio ursinus*) by female leopards (*Panthera pardus*) in the Waterberg Mountains, South Africa. *Folia primatologica: International Journal of Primatology* 83, 353-360.

King, J.S., Jenkins, D.J., Ellis, J.T., Fleming, P.J.S., Windsor, P.A., Slapeta, J. (2011). Implications of wild dog ecology on the sylvatic and domestic life cycle of *Neospora caninum* in Australia. *The Veterinary Journal* 188, 24-33.

Kluever, B.M., Breck, S.W., Howery, L.D., Krausman, P.R., Bergman, D.L. (2008). Vigilance in cattle: the influence of predation, social interactions, and environmental factors. *Rangeland Ecology & Management* 61, 321-328.

Kruuk, H. (1971). Surplus killing by carnivores. *Journal of Zoology, London* 166, 233-244.

Lennox, R.J., Gallagher, A.J., Ritchie, E.G., Cooke, S.J. (2018). Evaluating the efficacy of predator removal in a conflict-prone world. *Biological Conservation* 224, 277-289.

Letnic, M. (2014). 'Stop poisoning dingoes to protect native animals.' (UNSW Newsroom: University of New South Wales, Sydney, available at <http://newsroom.unsw.edu.au/news/science/stop-poisoning-dingoes-protect-native-mammals>, accessed 1 April 2014)

Letnic, M., Ritchie, E.G., Dickman, C.R. (2012). Top predators as biodiversity regulators: the dingo *Canis lupus dingo* as a case study. *Biological Reviews* 87, 390-413.

Linnell, J.D.C. (2011). The relative importance of predators and people in structuring and conserving ecosystems. *Conservation Biology* 25, 646-647.

Linnell, J.D.C., Lescureux, N. (2015). 'Livestock guarding dogs: cultural heritage icons with a new relevance for mitigating conservation conflicts.' (Norwegian Institute for Nature Research: Trondheim)

Linnell, J.D.C., Trouwborst, A., Fleurke, F.M. (2017). When is it acceptable to kill a strictly protected carnivore? Exploring the legal constraints on wildlife management within Europe's Bern Convention. *Nature Conservation* 21, 129-157.

Littin, K., Fisher, P., Beausoleil, N.J., Sharp, T. (2014). Welfare aspects of vertebrate pest control and culling: ranking control techniques for humaneness. *Scientific and Technical Review of the Office International des Epizooties* 33, 281-289.

MacNulty, D.R., Mech, L.D., Smith, D.W. (2007). A proposed ethogram of large-carnivore predatory behavior, exemplified by the wolf. *Journal of Mammalogy* 88, 595-605.

Mallick, S., Pauza, M., Eason, C., Mooney, N., Gaffney, R., Harris, S. (2016). Assessment of non-target risks from sodium fluoroacetate (1080), para-aminopropiophenone (PAPP) and sodium cyanide (NaCN) for fox-incursion response in Tasmania. *Wildlife Research* 43, 140-152.

Marker, L.L., Dickman, A.J., Macdonald, D.W. (2005). Perceived effectiveness of livestock-guarding dogs placed on Namibian farms. *Rangeland Ecology & Management* 58, 329-336.

Marks, C.A., Allen, L., Gigliotti, F., Busana, F., Gonzalez, T., Lindeman, M., Fisher, P.M. (2004). Evaluation of the trap tranquiliser device (TTD) for improving the humaneness of dingo trapping. *Animal Welfare* 13, 393-399.

Marks, CA, Hackman, C, Busana, F, Gigliotti, F. (2000). Assuring that 1080 toxicosis in the red fox (*Vulpes vulpes*) is humane: fluoroacetic acid (1080) and drug combinations. *Wildlife Research* 27, 483-494.

Mason, D.R. (1982). *Studies on the biology and ecology of the warthog Phacochoerus aethiopicus sundevalli (Lönnberg, 1908) in Zululand*. PhD thesis, University of Pretoria.

McLeod, R. (2016). 'Cost of pest animals in NSW and Australia, 2013-14. Report prepared for the NSW Natural Resources Commission.' (eSYS Development Pty Ltd: Sydney)

McManus, J.S., Dickman, A.J., Gaynor, D., Smuts, B.H., Macdonald, D.W. (2014). Dead or alive? Comparing costs and benefits of lethal and non-lethal human-wildlife conflict mitigation on livestock farms. *Oryx* 49, 687-695.

Mech, L.D. (1966). 'The wolves of Isle Royale.' (National Parks Fauna Series No. 7. U.S. Government Printing Office: Washington)

Meek, P.D., Brown, S.C., Wishart, J., Milne, H., Aylett, P., Humphrys, S., Ballard, G., Fleming, P.J.S. (2019). Efficacy of lethal-trap devices to improve the welfare of trapped wild dogs. *Wildlife Research* xx, xx-xx.

Meek, P.D., Jenkins, D.J., Morris, B., Ardler, A.J., Hawksby, R.J. (1995). Use of two humane leg-hold traps for catching pest species. *Wildlife Research* 22, 733-739.

Meek, P.D., Shorter, K., Falzon, G. (2018). Do lethal trap devices threaten foot-hold trap capture efficacy? *International Journal of Pest Management* 2018.1462538.

Mellor, D.J., Beausoleil, N.J. (2015). Extending the 'Five Domains' model for animal welfare assessment to incorporate positive welfare states. *Animal Welfare* 24, 241-253.

Mellor, D.J., Reid, C.S.W. (1994). *Concepts of animal well-being and predicting the impact of procedures on experimental animals*. In 'Improving the well-being of animals in the research environment.' Eds R Baker, G Jenkin, D Mellor. pp. 3-18. (Australian and New Zealand Council for the Care of Animals in Research and Teaching: Sydney, Australia)

Minnie, L. (2009). *Socio-economic and ecological correlates of leopard-stock farmer interactions in the Baviaanskloof Mega-Reserve, Eastern Cape*. Masters thesis, Nelson Mandela Metropolitan University.

Minnie, L., Avenant, N.L., Kamler, J.F., Butler, H., Parker, D.M., Drouilly, M., Du Plessis, J.J., Do Linh San, E. (2016a). *A conservation assessment of Canis mesomelas. The Red List of Mammals of South Africa, Swaziland and Lesotho*. Eds MF Child, L Roxburgh, E Do Linh San, D Raimondo, HT Davies-Mostert. (South African National Biodiversity Institute, Endangered Wildlife Trust: Pretoria)

Minnie, L., Boshoff, A.F., Kerley, G.I.H. (2015). Vegetation type influences livestock predation by leopards: implications for conservation in agro-ecosystems. *African Journal of Wildlife Research* 45, 204-214.

- Minnie, L., Gaylard, A., Kerley, G.I.H. (2016b). Compensatory life history responses of a mesopredator may undermine carnivore management efforts. *Journal of Applied Ecology* 53, 379-387.
- Minnie, L., Zalewski, A., Zalewska, H., Kerley, G.I.H. (2018). Spatial variation in anthropogenic mortality induces a source–sink system in a hunted mesopredator. *Oecologia* 186, 939-951.
- Moll, R.J., Killion, A.K., Montgomery, R.A., Tambling, C.J., Hayward, M.W. (2016). Spatial patterns of African ungulate aggregation reveal complex but limited risk effects from reintroduced carnivores. *Ecology* 97, 1123-1134.
- Moll, R.J., Redilla, K.M., Mudumba, T., Muneza, A.B., Gray, S.M., Abade, L., Hayward, M.W., Millspaugh, J.J., Montgomery, R.A. (2017). The many faces of fear: a synthesis of the methodological variation in characterizing predation risk. *Journal of Animal Ecology* 86, 749-765.
- Mormile, J.E., Hill, C.M. (2017). Living with urban baboons: exploring attitudes and their implications for local baboon conservation and management in Knysna, South Africa. *Human Dimensions of Wildlife* 22, 99-109.
- Newsome, T.M., Ballard, G., Crowther, M.S., Dellinger, J.A., Fleming, P.J.S., Glen, A.S., Greenville, A.C., Johnson, C.N., Letnic, M., Moseby, K.E., Nimmo, D.G., Nelson, M.P., Read, J.L., Ripple, W.J., Ritchie, E.G., Shores, C.R., Wallach, A.D., Wirsing, A.J., Dickman, C.R. (2015). Resolving the value of the dingo in ecological restoration. *Restoration Ecology* 23, 201-208.
- O’Bryan, C.J., Braczkowski, A.R., Beyer, H.L., Carter, N.H., Watson, J.E.M., McDonald-Madden, E. (2018). The contribution of predators and scavengers to human well-being. *Nature Ecology & Evolution* 2, 229-236.
- Palmer, C. (2010). 'Animal ethics in context.' (Columbia University Press: New York)
- Palmer, M.S., Fieberg, J., Swanson, A., Kosmala, M., Packer, C. (2017). A ‘dynamic’ landscape of fear: prey responses to spatiotemporal variations in predation risk across the lunar cycle. *Ecology Letters* 20, 1364-1373.
- Peterson, R.O., Vucetich, J.A., Bump, J.M., Smith, D.W. (2014). Trophic cascades in a multicausal world: Isle Royale and Yellowstone. *Annual Review of Ecology, Evolution, and Systematics* 45, 325-345.
- Potgieter, G., Kerley, G.I.H., Marker, L.L. (2016). More bark than bite? The role of livestock guarding dogs in predator control on Namibian farmlands. *Oryx* 50, 514- 522.
- Potgieter, G.C., Marker, L.L., Avenant, L.L., Kerley, G.I.H. (2013). Why Namibian farmers are satisfied with the performance of their livestock guardian dogs. *Human Dimensions of Wildlife* 18, 403-415.

Power, R.J., Compion, R.X.S. (2009). Lion predation on elephants in the Savuti, Chobe National Park, Botswana. *African Zoology* 44, 36-44.

Reddiex, B., Forsyth, D.M., McDonald-Madden, E., Einoder, L.D., Griffioen, P.A., Chick, R.R., Robley, A.J. (2006). Control of pest mammals for biodiversity protection in Australia. I. Patterns of control and monitoring. *Wildlife Research* 33, 691-709.

Riginos, C. (2015). Climate and the landscape of fear in an African savanna. *Journal of Animal Ecology* 84, 124-133.

Ripple, W.J., Beschta, R.L. (2004). Wolves and the ecology of fear: can predation risk structure ecosystems? *Bioscience* 54, 755-766.

Ripple, W.J., Beschta, R.L. (2012). Trophic cascades in Yellowstone: the first 15 years after wolf reintroduction. *Biological Conservation* 145, 205-213.

Ripple, W.J., Estes, J.A., Beschta, R.L., Wilmers, C.C., Ritchie, E.G., Hebblewhite, M., Berger, J., Elmhagen, B., Letnic, M., Nelson, M.P., Schmitz, O.J., Smith, D.W., Wallach, A.D., Wirsing, A.J. (2014). Status and ecological effects of the world's largest carnivores. *Science* 343, 151-163.

Ritchie, E.G., Elmhagen, B., Glen, A.S., Letnic, M., Ludwig, G., McDonald, R.A. (2012). Ecosystem restoration with teeth: what role for predators? *Trends in Ecology and Evolution* 27, 265-271.

RSPCA. (2010). 'RSPCA Policy E02 Management of wild animals (Article ID: 422, Revision 1, Updated 7 December 2010).' (Royal Society for the Prevention of Cruelty to Animals (RSPCA): Canberra)

RSPCA. (2016). 'Position statement: How can pest control programs be made more humane? (Article ID: 152; Revision 2; Updated 16 March 2016).' (Royal Society for the Prevention of Cruelty to Animals (RSPCA): Canberra)

Russell, J.C., Jones, H.P., Armstrong, D.P., Courchamp, F., Kappes, P.J., Seddon, P.J., Opper, S., Rauzon, M.J., Cowan, P.E., Rocamora, G., Genovesi, P., Bonnaud, E., Keitt, B.S., Holmes, N.D., Tershy, B.R. (2016). Importance of lethal control of invasive predators for island conservation. *Conservation Biology* 30, 670-672.

Rust, N.A., Whitehouse-Tedd, K.M., MacMillan, D.C. (2013). Perceived efficacy of livestock-guarding dogs in South Africa: implications for cheetah conservation. *Wildlife Society Bulletin* 37, 690-697.

Saunders, G., McLeod, L. (2007). 'Improving fox management strategies in Australia.' (Bureau of Rural Sciences: Canberra)

Schmitz, O.J. (2008). Effects of predator hunting mode on grassland ecosystem function. *Science* 319, 952-954.

Schwartz, D. (2016). RSPCA wants to stop 'cruel' dingo cull of feral goats on Great Barrier Reef island. *ABC News* 28 July 2016, <http://www.abc.net.au/news/2016-07-28/rspca-seeks-to-stop-dingo-cull-of-feral-goats-on-barrier-reef/7668268>.

Sharp, T., Saunders, G. (2011). 'A model for assessing the relative humaneness of pest animal control methods.' (Australian Government Department of Agriculture, Fisheries and Forestry: Canberra, ACT)

Sherley, M. (2007). Is sodium fluoroacetate (1080) a humane poison? *Animal Welfare* 16, 449-458.

Sinclair, A.R.E., Pech, R.P., Dickman, C.R., Hik, D., Mahon, P., Newsome, A.E. (1998). Predicting the effects of predation on conservation of endangered prey. *Conservation Biology* 12, 564-575.

Skead, C.J. (Eds AF Boshoff, GIH Kerley, PH Lloyd (2007). 'Historical incidence of the larger land mammals in the broader Eastern Cape, Second Edition.' (Centre for African Conservation Ecology, Nelson Mandela Metropolitan University: Port Elizabeth)

Skead, C.J. (Eds AF Boshoff, GIH Kerley, PH Lloyd (2011). 'Historical incidence of the larger land mammals in the broader Western and Northern Cape, Second edition ' (Centre for African Conservation Ecology, Nelson Mandela Metropolitan University: Port Elizabeth)

Slagle, K., Bruskotter, J.T., Singh, A.S., Schmidt, R.H. (2017). Attitudes toward predator control in the United States: 1995 and 2014. *Journal of Mammalogy* 98, 7-16.

Smith, B., Appleby, R. (2018). Promoting human-dingo co-existence in Australia: moving toward more innovative methods of protecting livestock rather than killing dingoes (*Canis dingo*). *Wildlife Research* 45, 1-15.

Somers, M.J. (1992). *The implications of social structure for the conservation and control of a warthog Phacochoerus aethiopicus population in the Andries Vosloo Kudu Reserve, Eastern Cape province*. MSc thesis, University of Pretoria.

Somers, M.J., Davies-Mostert, J., Mzileni, N., Swanepoel, L.H., Do Linh San, E., Botha, A.J., Tjelele, J., Dumalisile, L., Marnewick, K., Tafani, M., Hunnicutt, A., Tambling, C., Minnie, L., Hawkins, H.-J. (2018). *Biology, ecology and interaction of other predators with livestock*. In 'Livestock predation and its management in South Africa: a scientific assessment.' Eds GIH Kerley, SL Wilson, D Balfour. pp. 228-254. (Centre for African Conservation Ecology, Nelson Mandela University: Port Elizabeth, South Africa)

Soulé, M.E., Noss, R. (1998). Rewilding and biodiversity: complementary goals for continental conservation. *Wild Earth* 8, 1-11.

Suryawanshi, K.R., Bhatia, S., Bhatnagar, Y.V., Redpath, S., Mishra, C. (2014). Multiscale factors affecting human attitudes toward snow leopards and wolves. *Conservation Biology* 28, 1657-1666.

Swaigood, R.R. (2010). The conservation-welfare nexus in reintroduction programmes: a role for sensory ecology. *Animal Welfare* 19, 125-137.

Terborgh, J., Estes, J.A. (2010). 'Trophic cascades: predator, prey, and the changing dynamics of nature.' (Island Press: Washington D.C.)

Thaker, M., Vanak, A.T., Owen, C.R., Ogden, M.B., Niemann, S.M., Slotow, R. (2011). Minimizing predation risk in a landscape of multiple predators: effects on the spatial distribution of African ungulates. *Ecology* 92, 398-407.

Thinley, P., Rajaratnam, R., Lassoie, J.P., Morreale, S.J., Curtis, P.D., Vernes, K., Leki, L., Phuntsho, S., Dorji, T., Dorji, P. (2018). The ecological benefit of tigers (*Panthera tigris*) to farmers in reducing crop and livestock losses in the eastern Himalayas: implications for conservation of large apex predators. *Biological Conservation* 219, 119-125.

Thorn, M., Green, M., Dalerum, F., Bateman, P.W., Scott, D.M. (2012). What drives human-carnivore conflict in the north west province of South Africa? *Biological Conservation* 150, 23-32.

Torres, M. (2015). The case for intervention in nature on behalf of animals: a critical review of the main arguments against intervention. *Relations: Beyond Anthropocentrism* 3, 33-49.

Treves, A., Krofel, M., McManus, J. (2016). Predator control should not be a shot in the dark. *Frontiers in Ecology and the Environment* 14, 380-388.

Twigg, L.E., Parker, R.W. (2010). Is sodium fluoroacetate (1080) a humane poison? The influence of mode of action, physiological effects, and target specificity. *Animal Welfare* 19, 249-263.

Valeix, M., Hemson, G., Loverage, A.J., Mills, G., Macdonald, D.W. (2012). Behavioural adjustments of a large carnivore to access secondary prey in a human-dominated landscape. *Journal of Applied Ecology* 49, 73-81.

van Bommel, L. (2010). 'Guardian dogs: best practice manual for the use of livestock guardian dogs.' (Invasive Animal Cooperative Research Centre: Canberra)

van Bommel, L., Johnson, C.N. (2012). Good dog! Using livestock guardian dogs to protect livestock from predators in Australia's extensive grazing systems. *Wildlife Research* 39, 220-229.

van Bommel, L., Johnson, C.N. (2014). *Protecting livestock while conserving ecosystem function: non-lethal management of wild predators*. In 'Carnivores of Australia: past, present and future.' Eds AS Glen, CR Dickman. pp. 323-353. (CSIRO Publishing: Collingwood, Australia)

van Bommel, L., Johnson, C.N. (2015). How guardian dogs protect livestock from predators: territorial enforcement by Maremma sheepdogs. *Wildlife Research* 41, 662-672.

van Bommel, L., Johnson, C.N. (2016). Livestock guardian dogs as surrogate top predators? How Maremma sheepdogs affect a wildlife community. *Ecology and Evolution* 6, 6702-6711.

van Eeden, L.M., Crowther, M.S., Dickman, C.R., Macdonald, D.W., Ripple, W.J., Ritchie, E.G., Newsome, T.M. (2018). Managing conflict between large carnivores and livestock. *Conservation Biology* 32, 26-34.

Van Sittert, L. (1998). Keeping the enemy at bay: the extermination of wild Carnivora in the Cape Colony, 1889-1910. *Environmental History* 3, 333-356.

Wallach, A.D. (2014). Australia should enlist dingoes to control invasive species. *The Conversation* (<https://theconversation.com/australia-should-enlist-dingoes-to-control-invasive-species-24807>) 26 March, 2014.

Wallach, A.D., Bekoff, M., Batavia, C., Nelson, M.P., Ramp, D. (2018). Summoning compassion to address the challenges of conservation. *Conservation Biology* 32, 1255-1265.

Wallach, A.D., Bekoff, M., Nelson, M.P., Ramp, D. (2015). Promoting predators and compassionate conservation. *Conservation Biology* 29, 1481-1484.

West, P. (2008) *Assessing invasive animals in Australia 2008*. National Land and Water Resources Audit, The Invasive Animals Cooperative Research Centre, Canberra.

Williams, S.T., Maree, N., Taylor, P., Belmain, S.R., Keith, M., Swanepoel, L.H. (2017). Predation by small mammalian carnivores in rural agro-ecosystems: an undervalued ecosystem service? *Ecosystem Services* xx, xx-xx.

Woinarski, J.C.Z., Burbidge, A.A., Harrison, P.L. (2015). Ongoing unraveling of a continental fauna: decline and extinction of Australian mammals since European settlement. *Proceedings of the National Academy of Sciences* 112, 4531-4540.

Yelland, L. (2001). 'Holding the line: a history of the South Australian Dog Fence Board, 1947 to 2000.' (Primary Industries and Resources South Australia: Adelaide)