

SUPPLEMENTAL TABLE S2: Staging investigations

Total patients	n = 390	
Chest X-ray (CXR)	Number of studies	Percentage
Total	n = 163	23.1%
Normal study	82	50.3%
Pulmonary effusions	17	10.4%
Mediastinal mass	26	16%
Lung metastasis	12	8%
Infections	7	4.2%
Paravertebral	3	1.8%
Skull X-ray	Number of studies	Percentage
Total	n = 41	10.5%
Normal	20	48.7%
Coin lesions	18	43.9%
Orbital lesions	2	4.9%
Other	1	2.4%
Skeletal screening	Number of studies	Percentage
Total	n = 45	11.5%
No bone lesions	20	44.4%
Bone lesions reported	25	55.6%
Abdominal ultrasound (U/S)	Number of studies	Percentage
Total	n = 205	52.6%
Adrenal	97	47.3%
Abdominal	51	24.9%
Paraspinal	8	39%
Liver lesions	12	58.5%
Lymphadenopathy	28	13.6%
CT/MRI scans soft tissue	Number of studies	Percentage
Total	n = 299	76.7%
Adrenal primary	126	42.1%
Nonadrenal primary	133	44.5%
Liver metastasis	13	43.5%
Paraspinal	15	50.2%
Lymphadenopathy	54	18.1%
Lung metastasis	15	5%
CT/MRI scans bone lesions	Number of studies	Percentage
Total	n = 136	34.9%
No bone lesions	46	33.8%
Bone lesions	90	66.2%
¹²³I-MIBG scans	Number of studies	Percentage
Total	n = 163	41.8%
Avid	141	86.5%
Nonavid	22	13.5%
Bone scans	Number of studies	Percentage
Total	n = 157	40.3%

Positive	97	61.8%
Negative	60	38.2%
Bone marrow aspirates	Number of studies	Percentage
Total	n = 321	82.3%
Involved	204	63.6%
Not involved	117	36.4%