





























- McIntosh, B. 1971. The aedine subgenus *Neomelanicion* Newstead (Culicidae, Diptera) in southern Africa with descriptions of two new species. *J. Entom. Soc. South Africa* 34: 319–333.
- McIntosh, B. 1978. Olifantsvlei (OLI). *Am. J. Trop. Med. Hyg.* 27: 404–405.
- Moncayo, A.C., J.D. Edman, and J.T. Finn. 2000. Application of geographic information technology in determining risk of eastern equine encephalomyelitis virus transmission. *J. Am. Mosq. Contr. Assoc.* 8: 28–35.
- Munhenga, G., B.D. Brooke, B. Spillings, L. Essop, R.H. Hunt, S. Midzi, D. Govender, L. Braack, and L.L. Koekemoer. 2014. Field study site selection, species abundance and monthly distribution of anopheline mosquitoes in the northern Kruger National Park, South Africa. *Malar. J.* 13: 27.
- Muspratt, J. 1955. Research on South African Culicini (Diptera, Culicidae). III. A check-list of the species and their distribution, with notes on taxonomy, bionomics and identification. *J. Entom. Soc. South Africa* 18: 149–207.
- Newhouse, V.F., R. Chamberlain, J. Johnston, and W.D. Sudia. 1966. Use of dry ice to increase mosquito catches of the CDC miniature light trap. *Mosq. News* 26: 30–35.
- Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrol. Earth Syst. Sci.* 4: 439–473.
- Reisen, W.K. 2010. Landscape epidemiology of vector-borne diseases. *Annu. Rev. Entomol.* 55: 461–483.
- Reiskind, M., R. Griffin, M. Janairo, and Hopperstad. 2017. Mosquitoes of field and forest: the scale of habitat segregation in a diverse mosquito assemblage. *Med. Vet. Entomol.* 31: 44–54.
- Roche, B., P. Rohani, A.P. Dobson, and J.-F. Guégan. 2012. The impact of community organization on vector-borne pathogens. *Am. Nat.* 181: 1–11.
- Rueda, L., K. Patel, R. Axtell, and R. Stinner. 1990. Temperature-dependent development and survival rates of *Culex quinquefasciatus* and *Aedes aegypti* (Diptera: Culicidae). *J. Med. Entomol.* 27: 892–898.
- Rutherford, M.C., L. Mucina, and W. Powrie. 2006. Biomes and bioregions of southern Africa. The vegetation of South Africa, Lesotho and Swaziland. South African National Biodiversity Institute, Pretoria. South Africa. pp. 30–51.
- Ruwanza, S. and D. Mulaudzi. 2018. Soil physico-chemical properties in Lapalala Wilderness old agricultural fields, Limpopo Province of South Africa. *Appl. Ecol. Environ. Res.* 16: 2475–2486.
- Schulz, K.H., J. Steyn, and R. Rose-Innes. 1958. A Culicine mosquito survey of the Kruger National Park. *Koedoe* 1: 189–200.
- Smith, D.L., J. Dushoff, and F.E. McKenzie. 2004. The risk of a mosquito-borne infection in a heterogeneous environment. *PLoS Biol.* 2: e368.
- Steyn, J.J., K. H. Schultz, and R. Innes. 1955. A Culicine mosquito survey of the upper Limpopo River Valley. *J. Entom. Soc. South Africa* 18: 238–246.
- Swanepoel, R., 2003. Classification, epidemiology and control of arthropod-borne viruses. In: J.A.W. Coetzer and R.C. Tustin (eds.), *Infectious Diseases Of Livestock*. Oxford University Press, Cape Town. South Africa.
- Swanepoel, R. and J. Coetzer. 2004. Rift valley fever. In: J.A.W. Coetzer and R.C. Tustin (eds.), *Infectious Diseases of Livestock*. Oxford University Press, Cape Town. South Africa.
- van Der Linde, A. 1982. Species richness and relative abundance of female mosquitoes at a site in the western Orange Free State. *J. Entom. Soc. South. Africa* 45: 57–67.
- van Eeden, C., J.H. Williams, T.G.H. Gerdes, E. van Wilpe, A. Viljoen, R. Swanepoel, and M. Venter. 2012. Shuni virus as cause of neurologic disease in horses. *Emerg. Infect. Dis.* 18: 318–321.
- van Niekerk, S., S. Human, J. Williams, E. van Wilpe, M. Pretorius, R. Swanepoel, and M. Venter. 2015. Sindbis and Middelburg old world alphaviruses associated with neurologic disease in horses, South Africa. *Emerg. Infect. Dis.* 21: 2225–2229.
- van Staden, P. and G. Bredenkamp. 2005. Major plant communities of the Marakele National Park. *Koedoe* 48: 59–70.
- Venter, M. 2018. Assessing the zoonotic potential of arboviruses of African origin. *Curr. Opin. Virol.* 28: 74–84.
- Venter, M., M. Pretorius, J.A. Fuller, E. Botha, M. Rakgotho, V. Stivaktas, C. Weyer, M. Romito, and J. Williams. 2017. West Nile Virus lineage 2 in horses and other animals with neurologic disease, South Africa, 2008–2015. *Emerg. Infect. Dis.* 23: 2060–2064.
- Verberk, W. 2012. Explaining general patterns in species abundance and distributions. *Nature Education Knowledge* 3: 38.
- Worth, C. and B. Meillon. 1960. Culicine mosquitoes (Diptera: Culicidae) recorded from the province of Mocambique (Portuguese East Africa) and their relationship to arthropod-borne viruses. *An. Inst. Med. Trop.* 17: 231–256.
- Worth, C.B. and H. Paterson. 1961. Culicine mosquitoes in southern Africa. *Rev. Ent. Moq.* 4: 65–80.
- Worth, C.B., H. Paterson, and B. De Meillon. 1961. The incidence of arthropod-borne viruses in a population of culicine mosquitoes in Tongaland, Union of South Africa (January, 1956, through April, 1960). *Am. J. Trop. Med. Hyg.* 10: 583–592.
- Yoccoz, N.G., J.D. Nichols, and T. Boulinier. 2001. Monitoring of biological diversity in space and time. *Trends Ecol. Evol.* 16: 446–453.
- Zhou, G., N. Minakawa, A. Githeko, and G. Yan. 2004. Spatial distribution patterns of malaria vectors and sample size determination in spatially heterogeneous environments: a case study in the west Kenyan highland. *J. Med. Entomol.* 41: 1001–1009.