

Do black-backed jackals exhibit spatial partitioning with African wild dogs and lions?

Jan F. Kamler¹ | Samantha Nicholson² | Ute Stenkewitz³ |
Tazarve Gharajehdaghpour⁴ | Harriet Davies-Mostert^{2,5}

¹Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Abingdon, UK

²Endangered Wildlife Trust, Modderfontein, South Africa

³University of Iceland's Research Centre at Snæfellsnes, Stykkishólmur, Iceland

⁴Canadian Wildlife Services, Environment and Climate Change Canada, Edmonton, AB, Canada

⁵Department of Zoology and Entomology, Eugène Marais Chair of Wildlife Management, Mammal Research Institute, University of Pretoria, Pretoria, South Africa

Correspondence: Jan F. Kamler, Wildlife Conservation Research Unit, Department of Zoology, Oxford University, Tubney House, Abingdon Road, Tubney, Abingdon OX13 5QL, UK.

Email: jan.f.kamler@gmail.com

Funding information

Wildlife Conservation Society

Keywords: *Canis mesomelas*, home-range overlap, *Lycaon pictus*, *Panthera leo*, South Africa, spatial avoidance

1 | INTRODUCTION

Interspecific competition among mammalian carnivore species is common and results in a range of different behavioural mechanisms to partition resources and coexist. For example, sympatric carnivore species can coexist via spatial, dietary, habitat or temporal partitioning (Dröge, Creel, Becker, & M'soka, 2017; Kamler, Stenkewitz, Klare, Jacobsen, & Macdonald, 2012; Vanak et al., 2013). Spatial partitioning (i.e. spatial avoidance) is a common mechanism for some, but not all, smaller carnivores to coexist with large carnivores, especially when larger carnivores kill smaller carnivores (Kamler, Ballard, Gilliland, & Mote, 2003; Marneweck et al., 2019; Vanak et al., 2013). For example, black-backed jackals (*Canis mesomelas*), hereafter referred to as jackals, killed cape foxes (*Vulpes chama*), and thus cape foxes spatially avoided jackal core areas when foraging (Kamler et al., 2012). In contrast, although jackals also killed bat-eared foxes (*Otocyon megalotis*) and black-footed cats (*Felis nigripes*), these species did not spatially avoid jackal core areas when they foraged, although bat-eared foxes avoided jackal core areas when denning (Kamler et al., 2012, 2015). Among large carnivores, lions (*Panthera leo*) kill African wild dogs (*Lycaon pictus*), therefore the latter spatially avoided areas of high lion activity (Creel & Creel, 2002; Darnell, Graf, Somers, Slotow,

& Gunther, 2014; Marneweck et al., 2019; Vanak et al., 2013). In contrast, although lions kill cheetahs (*Acinonyx jubatus*) and spotted hyaenas (*Crocuta crocuta*; Périquet et al., 2016), spatial avoidance was not found to occur among these species (Broekhuis, Cozzi, Valeix, McNutt, & Macdonald, 2013; Dröge et al., 2017; Vanak et al., 2013). Reasons for the differences in spatial avoidance of larger carnivores by smaller carnivores is not known, but could be due to the use of other behavioural mechanisms to facilitate coexistence (Broekhuis et al., 2013; Dröge et al., 2017; Kamler et al., 2012; Périquet et al., 2016; Vanak et al., 2013).

Compared to research on the relationships among large carnivores in Africa, the relationships between large carnivores and mesocarnivores have received little attention. This is surprising, given that large carnivores in Africa are sympatric with numerous mesocarnivores, such as jackals, throughout their distributions. Consequently, competitive interactions between large carnivores and mesocarnivores might be greater than is generally acknowledged. For example, jackals have reportedly been killed or consumed by lions (Schaller, 1972; Stander, 1992), spotted hyaenas (van Lawick & van Lawick-Goodall, 1970), brown hyaenas (*Hyaena brunnea*; Mills, 1982), leopards (*Panthera pardus*; Kamler, Loveridge, O'Donnell, & Macdonald, 2020), cheetahs (Hayward, Hofmeyr,

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *African Journal of Ecology* published by John Wiley & Sons Ltd.

O'Brien, & Kerley, 2006) and African wild dogs (Kamler, Davies-Mostert, Hunter, & Macdonald, 2007), yet the spatial relationships between jackals and larger carnivores have never been studied.

This study determined if spatial partitioning occurred at different scales between jackals and both lions and African wild dogs. Research was carried out on De Beers Venetia Limpopo Nature Reserve (VLNR), where jackals were confirmed to be killed by African wild dogs (Kamler et al., 2007). To our knowledge, this is the first study to investigate the spatial relationships between jackals and large carnivores. Our research will help determine whether jackals use spatial partitioning at different scales to coexist with large carnivores.

2 | METHODS

2.1 | Study site

We conducted research from on VLNR (316 km²) in Limpopo Province, northeastern South Africa (22°15'–22°30'S and 29°12'–29°18'E). The VLNR has the full array of African large carnivores, including lions, leopards, cheetahs, spotted hyaenas and brown hyaenas, and is enclosed by an electrified predator-proof fence. African wild dogs were the only large carnivore previously extirpated from VLNR, and a pack of 16 individuals was reintroduced there in January 2002, and thereafter their numbers fluctuated annually between 11 and 25 individuals (Davies-Mostert, Mills, & Macdonald, 2013). Lion numbers fluctuated between 12 and 30 across years, whereas population sizes of cheetah, leopard and both hyaena species were low but stable (Davies-Mostert et al., 2013). See Davies-Mostert et al., (2013) for a more detailed description of the study site.

2.2 | Data collection and analysis

We captured jackals using padded foothold traps (see Kamler, Jacobsen, & Macdonald, 2008 for more details) and fitted radio collars (Advanced Telemetry Systems; 190 g) on three captured jackals (1 male, 2 females), which represented three different family groups. No nontarget species were captured during the study. During the first year of monitoring, the male jackal changed groups, and therefore, a total of four different family groups were monitored during the study. The total number of jackal groups in VLNR was unknown. In 2008, three individual African wild dogs were collared from a pack that consisted of 4 adults and up to 11 pups. For lions, at least one male and two females from each pride were collared. Three prides occurred on VLNR during 2018, and each consisted of 3–5 adults and 2–3 cubs. Two few locations were obtained from two lone male lions that were killed in early 2008, so they were excluded from the analysis. Our capture, handling and radio-collaring protocols were approved under permit (AG10610) from the Limpopo Province Department of Agriculture and Environment.

To obtain locations of study animals, Yagi antennas were used to home in on study animals from a vehicle until a visual observation was made, and then, a UTM coordinate was recorded, although if animals were in thick vegetation then sometimes triangulating from short distances was necessary. Study animals were radio-tracked continuously on nearly a daily basis throughout the study on a rotational basis, which typically consisted of two tracking sessions per day (one in morning, one in afternoon), although we used only the first location per tracking session to ensure independence of the location data. When several individuals were collared within the same group, we only used locations of one individual per group. We determined the annual home-range size of study animals using 90% fixed-kernel density estimates (KDE; Worton, 1989) and 50% KDEs to represent their core areas (Kamler et al., 2012; Kamler, Stenkewitz, & Macdonald, 2013). See Kamler, Rostro-García, and Macdonald (2017) for more details about kernel analyses.

We compared data at the home-range level (hereafter, broad-scale) and within home ranges (hereafter, fine scale). At the broad-scale, overlap of home ranges and core areas were compared between species. At the fine scale, we used utilisation distributions (UD) which provide a useful summary of an animal's use of space within its home range, by transforming the spatial data into a probability distribution (Fieberg & Kochanny, 2005). To measure overlap of intensity of use within home ranges, we used the utilisation distribution overlap index (UDOI; Fieberg & Kochanny, 2005). For two non-uniformly distributed UDs, UDOI values > 1 indicate a high degree of overlap, whereas values < 1 indicate less overlap than expected (Fieberg & Kochanny, 2005). The UDOI was calculated at both 90% and 50% UD isopleth levels based on all locations collected in 2018, to represent intensity of use within home ranges and core areas, respectively. See Kamler et al. (2017) for more details about UDOI analyses. The UDOIs (90% and 50%) were compared between jackals and both African wild dogs and lions using Mann–Whitney *U* tests, to determine whether jackals spatially avoided one species more than the other. Although some individuals were monitored over several years, only in 2008 were members of the four jackal groups and all three lion prides radio collared, and therefore, we only used data from 2008 in our analyses.

3 | RESULTS

The annual home-range sizes (90% KDE) for the four jackal groups ranged from 2.7 to 9.0 km² (\bar{X} = 6.4 km²). The annual home range of the pack of African wild dogs was 339.5 km² in 2008. The annual home ranges of the three lion prides ranged from 112.8 to 208.5 km² (\bar{X} = 167.4 km²).

At the broadscale, all home ranges (90% KDE) of jackals were completely overlapped by those of African wild dogs and lions (Figure 1). Similarly, 3 jackal core areas (50% KDE) were completely overlapped by those of African wild dogs and lions, whereas 1 jackal core area did not overlap either species (Figure 1). At the fine scale, overlap of the UDs within home ranges was low between jackals and




FIGURE 1 The annual (a) home ranges (90% kernel density estimate [KDE]) and (b) core areas (50% KDE) representing four groups of black-backed jackals, one pack of African wild dogs, and three prides of lions that were monitored simultaneously during 2008 in De Beers Venetia Limpopo Nature Reserve, South Africa

African wild dogs ($\bar{X} \pm SE = 0.011 \pm 0.008$; range = 0.001–0.036) and between jackals and lions (0.034 ± 0.021 ; range = 0.000–0.255), and there was no significant difference in overlap between jackals and both African wild dogs and lions ($Z = 0.672$, $n = 16$, $p = .501$). We excluded the jackal core area that did not overlap either species, and the overlap of the UD within the remaining core areas was low between jackals and African wild dogs ($\bar{X} \pm SE = 0.006 \pm 0.006$; range = 0.000–0.018) and between jackals and lions (0.011 ± 0.008 ; range = 0.000–0.063), and there was no significant difference in overlap between jackals and both African wild dogs and lions ($Z = 0.000$, $n = 12$, $p = 1.000$).

4 | DISCUSSION

The extent of spatial partitioning between jackals and both African wild dogs and lions depended on scale. At the broadscale, jackals did not exhibit spatial partitioning with either African wild dogs or lions. In contrast, at the fine scale, jackals appeared to exhibit spatial partitioning with both African wild dogs and lions because the UDOI overlap was relatively low, which indicated that the likelihood of animals occurring in the same area within their home ranges was relatively small. The fine-scale spatial partitioning between jackals and African wild dogs and lions was not surprising, given that both large carnivores kill jackals. Although our sample size of collared jackals was relatively small, we assumed our study animals exhibited typical behaviour and movements of other jackals on VLNR, and thus we feel that our results adequately represent the mechanisms of spatial partitioning between jackals and both large carnivores. That being said, our results should be viewed with caution because our study occurred within a relatively small fenced reserve, so our results might not be applicable to large or unfenced reserves where movements of carnivores are less confined. Also, our study occurred during 1 year, therefore it does not represent the dynamic spatial relationships that might occur between carnivores over the long term. Furthermore, spatial relationships between carnivores likely differ among areas based on differences in large carnivore densities, prey abundance and diversity, and habitat (Gigliotti et al., 2020; du Preez, Hart, Loveridge, & Macdonald, 2015; Rostro-García, Kamler, & Hunter, 2015). Nonetheless, our results provide the first evidence of spatial partitioning between jackals and large carnivores, and we hope future researchers examine spatial partitioning between these species under a variety of environmental conditions across Africa, to help build a more complete understanding of their interactions.

The lack of spatial partitioning between jackals and both African wild dogs and lions at the broadscale was likely due to the excessively large home ranges of the latter two species. The annual home range of the African wild dog pack covered the entire area of VLNR, whereas the home ranges of the lion prides each covered most of VLNR, and thus there was not adequate space in VLNR for jackals to avoid the large carnivores at the home-range level. In comparison, the low overlap of UD was likely facilitated by the wide-ranging movements of African wild dogs and lions, which allowed jackals to have small areas of concentrated use within the larger home ranges and core area of both large carnivores. The movements and annual home ranges in our study were typical of that previously reported for jackals (Kamler, Stenkewitz, Gharajehdaghypour, & Macdonald, 2019), African wild dogs (Davies-Mostert et al., 2012; Pomila, McNutt, & Jordan, 2015) and lion prides (Loveridge et al., 2009), therefore similar spatial relationships probably occur in other areas where these species are sympatric. In addition to lions and African wild dogs, other large carnivores also kill jackals, so future research should investigate whether spatial partitioning occurs between jackals and leopards, cheetahs and hyaenas.

In addition to spatial partitioning at the fine scale, jackals may have avoided African wild dogs and lions via other mechanisms not investigated during this study. For example, jackals may have used habitat, dietary or temporal partitioning to facilitate coexistence with African wild dogs and lions, similar to that previously reported for other African carnivores (Broekhuis et al., 2013; Dröge et al., 2017; Kamler et al., 2012; Marneweck et al., 2019; Périquet, Fritz, & Revilla, 2015; Périquet et al., 2016; Vanak et al., 2013; Vogel, Somers, & Venter, 2019). Future research should investigate whether jackals use these additional mechanisms to coexist with African wild dogs, lions and other large carnivores.

ACKNOWLEDGEMENTS

We thank Global Vision International, J. Power, W. Collinson and K. Potgieter-Forsman for help with tracking the radio-collared animals. This project was an initiative of the Carnivore Conservation Programme of the Endangered Wildlife Trust, South Africa, and the Wildlife Conservation Research Unit, Oxford University, U.K. Funding for J.F.K. was provided by a Research Fellowship from the Wildlife Conservation Society, New York, and a Marie Curie Fellowship from the European Commission, Brussels, Belgium. Sponsorship was provided by Jaguar Land Rover South Africa and De Beers Consolidated Mines.

DATA AVAILABILITY STATEMENT

Data are available on request.

ORCID

Jan F. Kamler  <https://orcid.org/0000-0003-4148-2088>

REFERENCES

- Broekhuis, F., Cozzi, G., Valeix, M., McNutt, J. W., & Macdonald, D. W. (2013). Risk avoidance in sympatric large carnivores: Reactive or predictive? *Journal of Animal Ecology*, *82*, 1098–1105. <https://doi.org/10.1111/1365-2656.12077>
- Creel, S. C., & Creel, N. M. (2002). *The African Wild Dog: Behavior, ecology, and conservation*. Princeton, NJ: Princeton University Press.
- Darnell, A. M., Graf, J. A., Somers, M. J., Slotow, R., & Gunther, M. S. (2014). Space use of African wild dogs in relation to other large carnivores. *PLoS ONE*, *9*, e98846. <https://doi.org/10.1371/journal.pone.0098846>
- Davies-Mostert, H. T., Kamler, J. F., Mills, M. G. L., Jackson, C. R., Rasmussen, G. S. A., Groom, R. J., & Macdonald, D. W. (2012). Long-distance transboundary dispersal of African wild dogs among protected areas in southern Africa. *African Journal of Ecology*, *50*, 500–506.
- Davies-Mostert, H. T., Mills, M. G. L., & Macdonald, D. W. (2013). Hard boundaries influence African wild dogs' diet and prey selection. *Journal of Applied Ecology*, *50*, 1358–1366.
- Dröge, E., Creel, S., Becker, M. S., & M'soka, J. (2017). Spatial and temporal avoidance of risk within a large carnivore guild. *Ecology and Evolution*, *7*, 189–199.
- du Preez, B., Hart, T., Loveridge, A. J., & Macdonald, D. W. (2015). Impact of risk on animal behaviour and habitat transition probabilities. *Animal Behaviour*, *100*, 22–37. <https://doi.org/10.1016/j.anbehav.2014.10.025>
- Fieberg, J., & Kochanny, C. O. (2005). Quantifying home-range overlap: The importance of the utilization distribution. *Journal of Wildlife Management*, *69*, 1346–1359. [https://doi.org/10.2193/0022-541X\(2005\)69\[1346:QHOTIO\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2005)69[1346:QHOTIO]2.0.CO;2)
- Gigliotti, L. C., Slotow, R., Hunter, L. T. B., Fattebert, J., Sholto-Douglas, C., & Jachowski, D. S. (2020). Context dependency of top-down, bottom-up and density-dependent influences on cheetah demography. *Journal of Animal Ecology*, *89*, 449–459.
- Hayward, M. W., Hofmeyr, M., O'Brien, J., & Kerley, G. I. H. (2006). Prey preferences of the cheetah (*Acinonyx jubatus*) (Felidae: Carnivora): Morphological limitations or the need to capture rapidly consumable prey before kleptoparasites arrive? *Journal of Zoology*, *270*, 615–627. <https://doi.org/10.1111/j.1469-7998.2006.00184.x>
- Kamler, J. F., Ballard, W. B., Gilliland, R. L., & Mote, K. (2003). Spatial relationships between swift foxes and coyotes in northwestern Texas. *Canadian Journal of Zoology*, *81*, 168–172.
- Kamler, J. F., Davies-Mostert, H. T., Hunter, L., & Macdonald, D. W. (2007). Predation on black-backed jackals (*Canis mesomelas*) by African wild dogs (*Lycaon pictus*). *African Journal of Ecology*, *45*, 667–668.
- Kamler, J. F., Jacobsen, N. F., & Macdonald, D. W. (2008). Efficiency and safety of Soft Catch traps for capturing black-backed jackals and excluding non-target species. *South African Journal of Wildlife Research*, *38*, 113–116.
- Kamler, J. F., Loveridge, A. J., O'Donnell, H., & Macdonald, D. W. (2020). Effects of humans and large carnivores on the survival of black-backed jackals. *African Journal of Ecology*, *58*, In press.
- Kamler, J. F., Rostro-García, S., & Macdonald, D. W. (2017). Seasonal changes in social behavior and movements of bat-eared foxes in South Africa: Disease implications. *Journal of Mammalogy*, *98*, 1426–1433. <https://doi.org/10.1093/jmammal/gyx092>
- Kamler, J. F., Stenkewitz, U., Gharajehdaghpour, T., & Macdonald, D. W. (2019). Social organization, home ranges, and extraterritorial forays of black-backed jackals. *Journal of Wildlife Management*, *83*, 1800–1808.
- Kamler, J. F., Stenkewitz, U., Klare, U., Jacobsen, N. F., & Macdonald, D. W. (2012). Resource partitioning among cape foxes, bat-eared foxes, and black-backed jackals in South Africa. *Journal of Wildlife Management*, *76*, 1241–1253.
- Kamler, J. F., Stenkewitz, U., & Macdonald, D. W. (2013). Lethal and sublethal effects of black-backed jackals on cape foxes and bat-eared foxes. *Journal of Mammalogy*, *94*, 295–306. <https://doi.org/10.1644/12-MAMM-A-122.1>
- Kamler, J. F., Stenkewitz, U., Wilson, B., Herrick, J., Lamberski, N., Sliwa, A., & Macdonald, D. W. (2015). Ecological relationships of black-footed cats (*Felis nigripes*) and sympatric canids in South Africa. *Mammalian Biology*, *80*, 122–127.
- Loveridge, A. J., Valeix, M., Davidson, Z., Murindagomo, F., Fritz, H., & Macdonald, D. W. (2009). Changes in home range size of African lions in relation to pride size and prey biomass in a semi-arid savanna. *Ecography*, *32*, 953–962. <https://doi.org/10.1111/j.1600-0587.2009.05745.x>
- Marneweck, C., Marneweck, D. G., van Schalkwyk, O. L., Beverley, G., Davies-Mostert, H. T., & Parker, D. M. (2019). Spatial partitioning by a subordinate carnivore is mediated by conspecific overlap. *Oecologia*, *191*, 531–540. <https://doi.org/10.1007/s00442-019-04512-y>
- Mills, M. G. L. (1982). *Hyaena brunnea*. *Mammalian Species*, *194*, 1–5.
- Périquet, S., Fritz, H., & Revilla, E. (2015). The Lion King and Hyaena Queen: Large carnivore interactions and coexistence. *Biological Reviews*, *90*, 1197–1214.
- Périquet, S., Mapendere, C., Revilla, E., Banda, J., Macdonald, D. W., Loveridge, A. J., & Fritz, H. (2016). A potential role for interference competition with lions in den selection and attendance by spotted hyaenas. *Mammalian Biology*, *81*, 227–234.
- Pomila, M. A., McNutt, J. W., & Jordan, N. R. (2015). Ecological predictors of African wild dog ranging patterns in northern Botswana. *Journal of Mammalogy*, *96*, 1214–1223. <https://doi.org/10.1093/jmammal/gyv130>

- Rostro-García, S., Kamler, J. F., & Hunter, L. T. B. (2015). To kill, stay or flee: The effects of lions and landscape factors on habitat and kill site selection of cheetahs in South Africa. *PLoS ONE*, *10*, e0117743. <https://doi.org/10.1371/journal.pone.0117743>
- Schaller, G. B. (1972). *The Serengeti Lion: A study of predator-prey relations*. Chicago, IL: University of Chicago Press.
- Stander, P. E. (1992). Foraging dynamics of lions in a semi-arid environment. *Canadian Journal of Zoology*, *70*, 8–21.
- van Lawick, H., & van Lawick-Goodall, J. (1970). *Innocent Killers*. Boston, MA: Houghton Mifflin.
- Vanak, A. T., Fortin, D., Thaker, M., Ogden, M., Owen, C., Greatwood, S., & Slotow, R. (2013). Moving to stay in place: Behavioral mechanisms for coexistence of African large carnivores. *Ecology*, *94*, 2619–2631. <https://doi.org/10.1890/13-0217.1>
- Vogel, J. T., Somers, M. J., & Venter, J. A. (2019). Niche overlap and dietary resource partitioning in an African carnivore guild. *Journal of Zoology*, *309*, 212–223.
- Worton, B. J. (1989). Kernel methods for estimating the utilization distribution in home-range studies. *Ecology*, *70*, 164–168. <https://doi.org/10.2307/1938423>

How to cite this article: Kamler JF, Nicholson S, Stenkewitz U, Gharajehdaghpour T, Davies-Mostert H. Do black-backed jackals exhibit spatial partitioning with African wild dogs and lions?. *Afr J Ecol*. 2020;58:552–556. <https://doi.org/10.1111/aje.12740>