


# A rapid and simple single-step method for the purification of *Toxoplasma gondii* tachyzoites and bradyzoites

Saeed El-Ashram<sup>1,2</sup>  | Yu Zhang<sup>1</sup> | Yongsheng Ji<sup>3</sup> | Dina Salama<sup>4,5</sup> | Kun Mei<sup>1</sup> | Li Zhili<sup>1</sup> | Huang Shujian<sup>1</sup> | Haoji Zhang<sup>1</sup> | Shawky M. Aboelhadid<sup>5</sup> | Reem A. Alajmi<sup>6</sup> | Dina M. Metwally<sup>6</sup> | Manal F. El-Khadragy<sup>7</sup> | Billy M. Hargis<sup>8</sup> | Guillermo Tellez-Isaias<sup>8</sup> | Beniamino T. Cenci-Goga<sup>9</sup> | Musafiri Karama<sup>10</sup> | Munyaradzi C. Marufu<sup>11</sup> | Fathi Abouhajer<sup>12</sup> | Gamal Ali Abdelhafez Hamady<sup>13</sup> | Abeer El Wakil<sup>14</sup> | Ibrahim Al Nasr<sup>15,16</sup> | Xun Suo<sup>17</sup>

<sup>1</sup>College of Life Science and Engineering, Foshan University, Foshan, Guangdong Province, China

<sup>2</sup>Faculty of Science, Kafrelsheikh University, Kafr el-Sheikh, Egypt

<sup>3</sup>School of Basic Medical Sciences, Division of Life Science and Medicine, University of Science and Technology of China, Hefei, China

<sup>4</sup>Department of Parasitology and Animal Disease, National Research Centre, Dokki, Giza, Egypt

<sup>5</sup>Parasitology Department, Faculty of Veterinary Medicine, Beni Suef University, Beni-Suef, Egypt

<sup>6</sup>Department of Zoology, Faculty of Science, King Saud University, Riyadh, Saudi Arabia

<sup>7</sup>Department of Zoology and Entomology, Faculty of Science, Helwan University, Cairo, Egypt

<sup>8</sup>Department of Poultry Science, University of Arkansas, Fayetteville, AR, USA

<sup>9</sup>Dipartimento di Medicina Veterinaria, Laboratorio di Ispezione degli Alimenti di Origine Animale, University of Perugia, Perugia, Italy

<sup>10</sup>Department of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa

<sup>11</sup>Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa

<sup>12</sup>Faculty of Education, Asmarya University for Islamic Sciences, Zliten, Libya

<sup>13</sup>Department of Animal Production, Faculty of Agriculture, Al-Azhar University, Nasr City, Cairo, Egypt

<sup>14</sup>Department of Biological & Geological Sciences, Alexandria University, Alexandria, Egypt

<sup>15</sup>College of Science and Arts in Unaizah, Qassim University, Unaizah, Saudi Arabia

<sup>16</sup>College of Applied Health Sciences in Ar Rass, Qassim University, Ar Rass, Saudi Arabia

<sup>17</sup>National Animal Protozoa Laboratory & College of Veterinary Medicine, China Agricultural University, Beijing, China

## Correspondence

Saeed El-Ashram, College of Life Science and Engineering, Foshan University, 18 Jiangwan street, Foshan 528231, Guangdong Province, China.  
Email: saeed\_elashram@yahoo.com

Xun Suo, National Animal Protozoa Laboratory & College of Veterinary Medicine, China Agricultural University, Beijing, 100193, China.  
Email: suoxun@cau.edu.cn.

## Funding information

Start-up Research Grant Program provided by Foshan University, Foshan city,

## Abstract

This study describes a simple method for the large-scale isolation of pure *Toxoplasma gondii* tachyzoites and bradyzoites. *T. gondii* tachyzoites were obtained from infected human foreskin fibroblasts (HFFs) and peritoneal exudates of mice, while tissue cysts containing bradyzoites were collected from chronically infected mice. Harvested cells and brain tissues were incubated in Hanks balanced salt solution (HBSS), containing 0.25% trypsin and 0.5% taurodeoxycholic acid (TDC) for 5 min. Subsequent washes in phosphate buffered saline (PBS) were conducted, and the cell viability of the preparations was good, as determined by flow cytometry and ability to reinfect HFF cells and propagate in mice. The purification procedure allowed for a rapid preparation of

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Veterinary Medicine and Science* published by John Wiley & Sons Ltd

Guangdong province for distinguished researchers, Guangdong Science and Technology Plan Project (Grant No:1244 0600 4560 7389XC) and School of Life Science and Engineering fund (Grant No: KLPREAD201801-02).

pure *T. gondii* tachyzoites and bradyzoites in sufficient quantity that can be used for downstream procedures. The advantage of the new method is that it is convenient and inexpensive.

#### KEYWORDS

flow cytometry, purification, *T. gondii* tachyzoites and bradyzoites, trypsin, taurodeoxycholic acid

## 1 | INTRODUCTION

*Toxoplasma gondii* is an intracellular protozoan parasite. Felids are the definitive hosts and a wide range of warm-blooded animals, including humans, rodents, birds, livestock and marine mammals are considered intermediate hosts. Sexual reproduction only occurs in felids. Sporozoites attach and penetrate the felid enterocytes after oocyst excystation in the intestinal lumen. Sporozoites undergo multiplication by endodyogeny to form tachyzoites, while bradyzoites are able to invade the intestinal cells, entering into the enteric phase of the sexual cycle. Bradyzoites switch to tachyzoites and go through several rounds of asexual division. After the sporulated oocysts are being ingested by intermediate hosts, sporulated oocysts will be excysted into the intestinal lumen. Sporozoites will find their way into extraintestinal sites to develop into tissue cysts in visceral organs, muscular and neural tissues (El-Ashram et al., 2015). A substantial amount of research on separation of eimerian oocysts and nematode eggs from faecal materials has been published over the last decade. Techniques for the concentration and purification of oocysts and eggs from faecal samples include saturated salt solution floatation, sucrose density, zinc sulphate and Percoll discontinuous density gradient centrifugation (Kawazoe et al., 1992). For example, sucrose gradient ultracentrifugation has been exploited to separate subcellular fractions of *Eimeria tenella* sporozoites. Recently, attention has focused on the isolation of endocytic organelles from macrophages by sucrose gradient (Lamberti et al., 2015). Furthermore, discontinuous sucrose gradients and isopycnic Percoll gradients have been used to isolate *Cryptosporidium* spp. oocysts and sporozoites (Arrowood & Sterling, 1987). Discontinuous sucrose gradient has been reported to be a simple and rapid procedure to obtain viable *Cryptosporidium* spp. oocysts (Bautista et al., 1999). In addition, *T. gondii* tachyzoites, which were obtained by two consecutive discontinuous sucrose gradient separations maintain their biological activity (Garberi et al., 1990). However, the aforementioned procedures are cumbersome, and tachyzoite yield is low. Although, it is technically possible to obtain pure zoites from oocysts collected from cat faeces using the sucrose gradient method; however, the method remains complicated. Currently, in vivo research on *Toxoplasma* using cats is very expensive because cats are an expensive species that are difficult to breed as their yields are very low. Therefore, most research on *T. gondii* is carried out using tachyzoites, which can be harvested in significant quantities from experimentally infected human foreskin fibroblasts (HFFs), the peritoneal cavity/exudates of


### Highlights

- Purification of *Toxoplasma gondii* tachyzoites and bradyzoites
- Harvested cells and tissues were incubated in digestive solution
- Preparation viability was assessed by flow cytometry and re-infection
- The new method is convenient and inexpensive.

experimentally infected mice, brain tissues and HFF cells. However, to obtain *T. gondii* tachyzoites from purification of murine peritoneal exudates, brain tissues and HFF cells remains a serious challenge. In this short communication, we report a very simple approach for the isolation of highly purified tachyzoites and bradyzoites from infected human foreskin fibroblasts (HFFs) and peritoneal exudates of mice.

## 2 | MATERIALS AND METHODS

Six- to eight- week-old specific pathogen free (SPF) female BALB/c mice were purchased from the Institute of Laboratory Animal Sciences, Chinese Academy of Medical Sciences (Beijing, China). Two genotypes of *T. gondii* were used in this study: the avirulent type II *T. gondii* Prugniaud (Pru) strain and the virulent *T. gondii* RH strain. The avirulent type II *T. gondii* Prugniaud (Pru) strain maintained by passage every 4–5 weeks in mice by oral infection with five cysts where tachyzoites of *T. gondii* RH strain were kept stored in liquid nitrogen. The experimental infection was carried out by inoculating tachyzoites of the *T. gondii* RH strain to mice intraperitoneally. Successful *T. gondii* infection was achieved within 4–5 days after intraperitoneal infection, followed by recovery of the parasites from the peritoneal exudate. Three days after infection, mice were sacrificed by cervical dislocation. Furthermore, HFF cells were incubated in DMEM medium (HyClone) supplemented with 8% fetal bovine serum (HyClone) and infected with the *T. gondii* RH strain tachyzoites ( $1 \times 10^7$  parasites) for 72 hr. Tachyzoites and tachyzoites were collected from the peritoneal cavity and brain tissues, respectively in 5 ml phosphate buffered saline (PBS). Peritoneal cells and brain tissues




**FIGURE 1** Flow chart summarizing the purification steps of *T. gondii* tachyzoites and bradyzoites

were washed three times in PBS, placed in 5 volumes of digestive medium (i.e. 5 volumes of digestive medium/ 1 volume of cell- or tissue-containing zoites)—Hanks balanced salt solution (HBSS) containing 0.25% trypsin and 0.5% taurodeoxycholic acid (TDC). After their washing thrice with PBS, 5 ml of the trypsin-TDC medium was added to a T-75 flask containing *Toxoplasma*-infected HFF cells. The mixture was incubated at 40°C for 5 min with intermittent shaking and free zoites were monitored by the light microscopy. The zoite-containing solution was diluted 10 times in HBSS solution at 1:10 (v:v). Tachyzoites or bradyzoites were collected by centrifugation at 4,000 × g for 5 min, and supernatant was discarded (Figure 1). Tachyzoites were washed three times with PBS, centrifuged at 4,000 × g for 5 min and propagated in mice by intraperitoneal inoculation. HFFs with 80%–90% confluence were infected with tachyzoites. The viability of the purified zoites was detected by a flow cytometer (FCM) as follows: the isolated tachyzoites and bradyzoites were added to Eppendorf tubes containing 5 μl of Annexin V-FITC (fluorescein isothiocyanate conjugated), 5 μl propidium iodide (PI) or 1.5 ml PBS. Furthermore, Annexin<sup>+</sup>/PI<sup>-</sup> (early apoptosis) and Annexin<sup>+</sup>/PI<sup>+</sup> (late apoptosis or already dead) parasites were sorted by fluorescence-activated cell sorter (FACS) (Demchenko, 2013; Yin, et al., 2015; Yin et al., 2015).


### 3 | RESULTS AND DISCUSSION

HFF and murine peritoneal exudates of *T. gondii* RH strain-infected HFF cells (Figure 2a,b) and mice (Figure 2c,d) are shown below before and after purification, respectively. Additionally, the purified bradyzoites are displayed in Figure 3. The viability of tachyzoites and bradyzoites was established by flow cytometry and re-infectivity. As shown in Figure 4, more than 99% of collected parasites were Annexin<sup>-</sup>/PI<sup>-</sup>, indicating the high viability of parasites.

There are several published reports on purification of *Toxoplasma* tachyzoites by passing the parasite through a 25-gauge needle, filtration through a 3-μm pore size, sucrose gradient density centrifugation (Garberi et al., 1990; Khan & Grigg, 2017). *Toxoplasma* tachyzoite purification can also be achieved by sonic vibration and trypsin digestion (Tsunematsu, 1960) or by a sintered glass filter. These procedures are often complex and pose scale-up challenges. Additionally, recovery rates can be low, and some methods have failed to produce pure tachyzoite preparations. In this study, we developed a rapid and simple method for purifying tachyzoites and bradyzoites with optimal yield and purity. The new proposed method has the following advantages: (a) zoites can be obtained quickly and easily; (b) the technique can be performed in low resource settings and capable of handling




**FIGURE 2** Tachyzoite-infected human foreskin fibroblast (HFF) cell line (a) and purified tachyzoites from HFF cell line (b1 and b2); and tachyzoite-infected peritoneal cells (c) and purified tachyzoites from HFF cell line (d)



**FIGURE 3** Purified bradyzoites from infected murine brain

larger numbers of cells and tissues; (c) zoites obtained in this way are free of cellular debris; and (d) the viability and infectivity of tachyzoites and bradyzoites are maintained. A more detailed study is required to explore the effect of trypsin on surface antigens of parasites.



**FIGURE 4** Viability of *T. gondii* tachyzoites by flow cytometry

#### ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (No. 31502071), Youth Innovative Talents Project of Guangdong province Education Department (No. 2017KQNCX212), Guangdong province (2017GDK07), Start-up Research Grant Program

provided by Foshan University, Foshan city, Guangdong province for distinguished researchers, Guangdong Science and Technology Plan Project (Grant No: 1244060045607389XC), and School of Life Science and Engineering fund (Grant No: KLPREAD201801-02).

### CONFLICTS OF INTEREST

The author declares no conflicts of interest.

### AUTHOR CONTRIBUTIONS

Saeed El-Ashram: Conceptualization, Investigation and Methodology.

Saeed El-Ashram, Yu Zhang, Yongsheng Ji, Dina Salama, Kun Mei, Li Zhili, Huang Shujian, Haoji Zhang, Shawky M. Aboelhadid, Reem A. Alajmi, Dina M. Metwally, Manal F. El-Khadragy, Billy M. Hargis, Guillermo Tellez-Isaias, Beniamino T. Cenci-Goga, Musafiri Karama, Munyaradzi C. Marufu, Fathi Abouhajer, Gamal Ali Abdelhafez Hamady, Abeer El Wakil, Ibrahim Al Nasr and Xun Suo: Writing-original draft, Writing-review and editing.

### AUTHOR CONTRIBUTION

**Saeed El-Ashram:** Conceptualization; Investigation; Methodology; Writing-original draft; Writing-review & editing. **Yu Zhang:** Writing-original draft; Writing-review & editing. **Yongsheng Ji:** Writing-original draft; Writing-review & editing. **Dina Salama:** Writing-original draft; Writing-review & editing. **Kun Mei:** Writing-original draft; Writing-review & editing. **Li Zhili:** Writing-original draft; Writing-review & editing. **Huang Shujian:** Writing-original draft; Writing-review & editing. **Haoji Zhang:** Writing-original draft; Writing-review & editing. **Shawky M Aboelhadid:** Writing-original draft; Writing-review & editing. **Reem Reem Alajmi:** Writing-original draft; Writing-review & editing. **dina metwally:** Writing-original draft; Writing-review & editing. **Manal El-Khadragy:** Writing-original draft; Writing-review & editing. **Billy Hargis:** Writing-original draft; Writing-review & editing. **Guillermo Tellez:** Writing-original draft; Writing-review & editing. **Beniamino Cenci-Goga:** Writing-original draft; Writing-review & editing. **Musafiri Karama:** Writing-original draft; Writing-review & editing. **Munyaradzi Marufu:** Writing-original draft; Writing-review & editing. **Fathi Abouhajer:** Writing-original draft; Writing-review & editing. **gamal ali abdelhafez hamady:** Writing-original draft; Writing-review & editing. **Abeer El Wakil:** Writing-original draft; Writing-review & editing. **Xun Suo:** Writing-original draft; Writing-review & editing.

### ETHICAL STATEMENT

All experiments were conducted according to the ethical standards and protocols approved by the Committee of Animal Experimentation of College of Life Science and Engineering, Foshan University, China (permission number 2019- FOSU- 04).

### PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1002/vms3.364>.

### ORCID

Saeed El-Ashram  <https://orcid.org/0000-0003-0389-1980>

### REFERENCES

- Arrowood, M. J., & Sterling, C. R. (1987). Isolation of *Cryptosporidium* oocysts and sporozoites using discontinuous sucrose and isopycnic Percoll gradients. *Journal of Parasitology*, *73*, 314–319. <https://doi.org/10.2307/3282084>
- Bautista, E., Pezzani, B. C., Cordoba, A., De Luca, M. M., & Basualdo, J. A. (1999). Relationship between discontinuous sucrose gradient and viability of *Cryptosporidium* sp. oocysts. *Revista Argentina De Microbiologia*, *31*, 188–192.
- Demchenko, A. P. (2013). Beyond annexin V: Fluorescence response of cellular membranes to apoptosis. *Cytotechnology*, *65*, 157–172. <https://doi.org/10.1007/s10616-012-9481-y>
- El-Ashram, S., Yin, Q., Barta, J. R., Khan, J., Liu, X., & Suo, X. (2015). Immunoproteomic technology offers an extraordinary diagnostic approach for *Toxoplasma gondii* infection. *Journal of Microbiological Methods*, *119*, 18–30. <https://doi.org/10.1016/j.mimet.2015.09.011>
- Garber, J. C., Blanco, J. C., Angel, S. O., Pzsenny, V., Arakelian, M. C., & Pastini, A. (1990). *Toxoplasma gondii*: A rapid method for the isolation of pure tachyzoites. Preliminary characterization of its genome. *Memorias do Instituto Oswaldo Cruz*, *85*, 429–434. <https://doi.org/10.1590/S0074-0276199000400007>
- Kawazoe, U., Tomley, F. M., & Frazier, J. A. (1992). Fractionation and antigenic characterization of organelles of *Eimeria tenella* sporozoites. *Parasitology*, *104*(Pt 1), 1–9.
- Khan, A., & Grigg, M. E. (2017). *Toxoplasma gondii*: Laboratory Maintenance and Growth. *Curr Protoc Microbiol*, *44*, 20C.21.21–20C.21.17.
- Lamberti, G., de Araujo, M. E., & Huber, L. A. (2015). Isolation of macrophage early and late endosomes by latex bead internalization and density gradient centrifugation. *Cold Spring Harbor Protocols*, *2015*(12), pdb.prot083451.
- Tsunematsu, Y. (1960). Purification of *Toxoplasma* by means of sonic vibration and tryptic digestion. *The American Journal of Tropical Medicine and Hygiene*, *9*, 556–561. <https://doi.org/10.4269/ajtmh.1960.9.556>
- Yin, Q., El-Ashram, S., Liu, H., Sun, X., Zhao, X., Liu, X., & Suo, X. (2015). Interferon-gamma release assay: An effective tool to detect early *Toxoplasma gondii* infection in mice. *PLoS One*, *10*, e0137808. <https://doi.org/10.1371/journal.pone.0137808>
- Yin, Q., El-Ashram, S., Liu, X. Y., & Suo, X. (2015). Early detection of *Toxoplasma gondii*-infected cats by interferon-gamma release assay. *Experimental Parasitology*, *157*, 145–149. <https://doi.org/10.1016/j.exppara.2015.08.015>

**How to cite this article:** El-Ashram S, Zhang Y, Ji Y, et al. A rapid and simple single-step method for the purification of *Toxoplasma gondii* tachyzoites and bradyzoites. *Vet. Med. Sci.* 2020;00:1–5. <https://doi.org/10.1002/vms3.364>