



Article

Rabies Vaccination of 6-Week-Old Puppies Born to Immunized Mothers: A Randomized Controlled Trial in a High-Mortality Population of Owned, Free-Roaming Dogs

Sintayehu Arega ¹, Anne Conan ¹, Claude T. Sabeta ^{2,3}, Jan E. Crafford ³, Jeanette Wentzel ⁴, Bjorn Reininghaus ⁵, Louise Biggs ⁶, Andrew L. Leisewitz ⁷, Melvyn Quan ³, Felix Toka ⁸ and Darryn L. Knobel ^{1,3,*}

- ¹ Center for Conservation Medicine and Ecosystem Health, Ross University School of Veterinary Medicine, Basseterre, St Kitts and Nevis; SintayehuArega@students.rossu.edu (S.A.); AConan@rossvet.edu.kn (A.C.);
 - ² Agricultural Research Council-Onderstepoort Veterinary Institute, OIE Rabies Reference Laboratory, Onderstepoort 0110, South Africa; SabetaC@arc.agric.za
 - ³ Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Onderstepoort 0110, South Africa; jannie.crafford@up.ac.za (J.E.C.); melvyn.quan@up.ac.za (M.Q.)
 - ⁴ Hans Hoheisein Wildlife Research Station, Faculty of Veterinary Science, University of Pretoria, Orpen 1364, South Africa; jeanette.wentzel@up.ac.za
 - ⁵ Mpumalanga Veterinary Services, Department of Agriculture, Rural Development, Land and Environmental Affairs, Thulamahashe 1365, South Africa; bjorn.reininghaus@gmail.com
 - ⁶ Department of Production Animal Studies, University of Pretoria, Onderstepoort 0110, South Africa; louise.biggs@up.ac.za
 - ⁷ Department of Companion Animal Clinical Studies, University of Pretoria, Onderstepoort 0110, South Africa; andrew.leisewitz@up.ac.za
 - ⁸ Center for Integrative Mammalian Research, Ross University School of Veterinary Medicine, Basseterre, St Kitts and Nevis; FToka@rossvet.edu.kn
- * Correspondence: dknobel@rossvet.edu.kn

Table S1. Results of sensitivity analysis for survival analysis (six through 13 weeks of age), considering subjects reported as lost or stolen by owners as dead (n = 22).

	Sterile Water		Rabies Vaccine		HRs (95% CI) for RV within Strata of Sex
	Mortality Rate ^a (Deaths/Dog- Years)	HR (95% CI) ^b	Mortality Rate ^a (Deaths/Dog- Years)	HR (95% CI) ^b	
Females	1,808 (15/8.3)	1 (reference)	3750 (32/8.5)	1.71 (0.87–3.34)	1.71 (0.87–3.34)
Males	3,496 (30/8.6)	1.62 (0.80–3.29)	2744 (25/9.1)	1.36 (0.68–2.71)	0.84 (0.46–1.55)

Notes: HR = hazard ratio. CI = confidence intervals; ^a Mortality rate per 1,000 dog-years; ^b Compared to reference category (control females).

Table S2. Results of sensitivity analysis for survival analysis (six through 13 weeks of age), censoring subjects that reportedly died from accidents (n = 5).

	Sterile Water		Rabies Vaccine		HRs (95% CI) for RV within Strata of Sex
	Mortality Rate ^a (Deaths/Dog-Years)	HR (95% CI) ^b	Mortality Rate ^a (Deaths/Dog- Years)	HR (95% CI) ^b	
Females	844 (7/8.3)	1 (reference)	2696 (23/8.5)	2.83 (1.12– 7.15)	2.83 (1.12–7.15)

Males	2914 (25/8.6)	3.37 (1.29–8.79)	2,195 (20/9.1)	2.23 (0.89–5.60)	0.66 (0.33–.34)
-------	---------------	---------------------	----------------	------------------	-----------------

Notes: HR = hazard ratio. CI = confidence intervals; ^a Mortality rate per 1,000 dog-years; ^b Compared to reference category (control females).