

Supplementary Materials for
Fire and biodiversity in the Anthropocene

Luke T. Kelly*, Katherine M. Giljohann, Andrea Duane, Núria Aquilué, Sally Archibald, Enric Batllori, Andrew F. Bennett, Stephen T. Buckland, Quim Canelles, Michael F. Clarke, Marie-Josée Fortin, Virgilio Hermoso, Sergi Herrando, Robert E. Keane, Frank K. Lake, Michael A. McCarthy, Alejandra Morán-Ordóñez, Catherine L. Parr, Juli G. Pausas, Trent D. Penman, Adrián Regos, Libby Rumpff, Julianna L. Santos, Annabel L. Smith, Alexandra D. Syphard, Morgan W. Tingley, Lluís Brotons

*Corresponding author. Email: ltkelly@unimelb.edu.au







Published 20 November 2020, *Science* **370**, eabb0355 (2020)
DOI: [10.1126/science.abb0355](https://doi.org/10.1126/science.abb0355)

This PDF file includes:







Table S1
References

Table S1.

Details of linked changes in fire regimes and biodiversity documented in Figure 2.

 <p>1. Saprotrophic fungi A hotter, drier climate is projected to cause larger and more severe fires in the boreal forests of Alaska, USA. Within burned stands of Black Spruce forest and spruce-aspen forest, saprotrophic fungi decline in areas that are severely burnt (37).</p>	 <p>2. Whitebark Pine Whitebark Pine populations in subalpine forests of western North America are declining through mortality caused by the Mountain Pine Beetle and White Pine Blister Rust. Yet, some individual White Pine trees are putatively pest-resistant and provide the foundation for future restoration efforts. Loss of pest-resistant trees from wildfire could reduce Whitebark Pine regeneration. Wildfires are predicted to increase in subalpine forests because of global climate change and this may place Whitebark Pine populations at further risk of decline (84).</p>	 <p>3. White Spruce Extreme fire weather is increasing fire frequency in Canadian boreal forests. Short intervals between fires negatively affect recruitment of obligate seeders such as White Spruce. Global climate change is likely to increase fire activity in these forests and amplify the negative impact of drought-fire interactions on White Spruce (11).</p>
 <p>4. Lodgepole Pine Subalpine forests of Greater Yellowstone, USA, have been resilient to stand-replacing fires that historically burned</p>	 <p>5. Greater Sage-grouse Invasive Cheatgrass increases fuel loads and continuity in deserts and shrublands of western USA, which alters</p>	 <p>6. Chaparral In southern California, native shrublands known as chaparral support exceptionally high plant</p>




<p>at intervals of 100-300 years. However, fire intervals are projected to decrease as the climate warms. Short-interval stand-replacing fires in Lodgepole Pine forests reduce tree generation and biological legacies. There will likely be marked changes in forest structure and function if short-interval fires become more common in Lodgepole Pine forests (36).</p>	<p>regional fire regimes. In turn, an increase in fire occurrence reduces habitat for Greater Sage-grouse that prefers to forage in dense sagebrush (53).</p>	<p>diversity. Short intervals between fires, associated with proximity to trails and roads in recreation areas, are converting chaparral into vegetation dominated largely by exotic herbaceous cover. Landscape disturbances such as trail construction and fuel breaks also provide the means for exotic herbaceous species to colonize open vegetation (45).</p>
<div data-bbox="277 730 513 968" data-label="Image"> </div> <p>7. Eastern Collared Lizard Prior to European settlement, fires were common in the woodlands and savannas of midwestern USA. Fire suppression throughout much of the 20th century shifted open woodlands and savanna to a woodland with a dense woody understory. This regime shift fragmented habitats of exposed bedrock used by the Eastern Collared Lizard, followed by local extinction of the species (101).</p>	<div data-bbox="688 730 924 968" data-label="Image"> </div> <p>8. Great Plains grasslands The Great Plains of North America comprise extensive fire-prone grasslands. Exclusion of anthropogenic fire, including active fire suppression, has reduced the frequency of fire and led to the transition of species-rich grassland to areas dominated by woody plants such as Juniper species. Overgrazing by domestic livestock has also removed herbaceous species that contribute to grassland fires. Under global climate change, fire regimes in the Great Plains are expected to exhibit ranges of variability that have no historical analogue over millennia (16).</p>	<div data-bbox="1099 730 1334 968" data-label="Image"> </div> <p>9. Hylid frog The Endangered frog <i>Bromeliophyla melacaena</i> occurs in mountain pine forests and broadleaf cloud forests of Honduras. Fire is naturally rare in these forests but intentional use of fire for agriculture is reducing habitat for this species. Government incentivization for the cultivation of coffee in protected areas is one reason for increasing alteration of primary forest habitat. In addition, a woodboring pine beetle has recently caused high tree mortality, which encourages harvesting of timber and, in turn, use of intentional fires to convert areas harvested for timber into cropland. Climate change is expected to further reduce</p>

		suitable habitat for this hylid frog, and make populations more vulnerable to fire (24).
 <p>10. Ant communities In the cerrado ecosystems of Brazil, fire suppression is practiced in most nature reserves and protected areas because of the perception that fire is damaging to biodiversity. Reduction in fire occurrence is exacerbated by crops, pasture and forest plantations that fragment savanna landscapes. Fire suppression has transformed some savanna vegetation into forests, through woody encroachment, causing biodiversity losses and marked changes in ecological processes. Ant communities are less diverse when fire is excluded (43).</p>	 <p>11. ‘Pineapples’ Natural fire ignitions are rare in the shrubland ecosystem of central Chile called matorral. Since the 19th century, human-driven ignitions associated with agriculture and transportation have significantly increased fire frequency in matorral and are driving evolutionary changes in seed traits of the native herbaceous plant ‘pineapples’ (76).</p>	 <p>12. Forest butterflies Until recent decades, tropical broadleaf forests of the Amazon rarely experienced large fires. Extensive fires in the Amazon rainforest are a result of deforestation, habitat fragmentation and climate change, underpinned by social, political and economic changes. Forest specialists such as Leaf Wing Butterfly decline in areas burnt recurrently (41).</p>
 <p>13. Dartford Warbler In northern Spain, land abandonment continues to shift mosaics of farmland and</p>	 <p>14. Common Heather Common heather occurs across Europe in a variety of temperate, boreal and</p>	 <p>15. Spur-thighed Tortoise Fire frequency and extent is expected to increase in the Mediterranean woodlands</p>

<p>open forest to dense, fire-prone forests. A hotter and drier climate exacerbates the risk of very large fires. Higher tree cover has increased populations of some forest bird species, while causing declines in open-country species. Threatened bird species that prefer open habitat, such as Dartford Warbler, will benefit from open spaces created by wildfires (47).</p>	<p>montane ecosystems. In Norway, coastal populations of this shrub species have experienced higher fire frequency than other ecosystems, under millenia of traditional burning for agriculture. Smoke-induced germination, which helps plants exploit frequent fire, is found in populations from traditionally burnt shrublands but is lacking in ecosystems where fire is rare. Thus, anthropogenic fire has likely shaped the evolution of this species and conservation management will be enhanced by considering these adaptations (102).</p>	<p>and shrublands of northwest Africa because of global climate change, rural land abandonment and dense plantations of trees for timber production. Spur-thighed Tortoise can be killed by fires and areas recently burnt by wildfires do not provide suitable food and shelter resources. Short intervals between fires are expected to cause regional population crashes (103).</p>
<div data-bbox="272 978 509 1213" data-label="Image"> </div> <p>16. Savanna trees Humans are the main source of fires in the savannas of west and central Africa. In the savannas of Gabon's Bateke Plateaux there has been a change from annual to semi-annual fires over the past 40 years as governance has become less centralized. Fire are now lit several times a year by individual hunters to create pasture. Semi-annual fire regimes in the savannas of Gabon are increasing the survival and density of the dominant tree</p>	<div data-bbox="688 978 925 1213" data-label="Image"> </div> <p>17. Wild herbivores Reduction in fuel loads through increased livestock grazing has virtually excluded fire in savannah ecosystems of the Serengeti-Mara, Tanzania. Fire plays an important role in the function of this region and its exclusion could lead to an increase in bush encroachment and the displacement of wild herbivores such as wildebeest (18).</p>	<div data-bbox="1094 978 1331 1213" data-label="Image"> </div> <p>18. Cyprinid fish The fish <i>Enteromius seymouri</i> occurs in river systems in Malawi. The freshwater habitat of <i>E. seymouri</i> is declining due to sedimentation caused by agriculture. Fires used to clear land, or as part of agriculture, threaten wetlands in the northern tributaries of the species geographic range, through siltation from soil erosion (24).</p>

<p><i>Hymenocardia acida</i>. With less fuel to burn, semi-annual fires are patchier and cooler than the previous annual fires (104).</p>		
<div data-bbox="269 436 505 667" data-label="Image"> </div> <p>19. Red-fronted Brown Lemur The lemur <i>Eulemur rufifrons</i> occurs in moist lowland forests, montane forests and dry tropical forests in Madagascar. Populations of this species have declined due to habitat loss and fragmentation caused by deforestation fires, land clearing, and illegal logging. Hunting is also a threat. Uncontrolled fires are an ongoing threat, particularly in dry tropical forests in the western part of the species geographic range (24).</p>	<div data-bbox="688 436 924 667" data-label="Image"> </div> <p>20. Clanwilliam Cypress Conifers are disproportionately threatened by changes in fire regimes. In South Africa, Clanwilliam Cypress has declined in shrubland ecosystems called fynbos that, in some areas, are subject to high fire frequency. Increases in temperature and more frequent anthropogenic fires are projected to cause further declines of this species and highlight the need for conservation interventions (105).</p>	<div data-bbox="1097 436 1333 667" data-label="Image"> </div> <p>21. Eurasian steppe After the collapse of the Soviet Union, 12 million ha of cropland were abandoned in the steppe zone of Kazakhstan and livestock grazing ceased across large areas of fields and grasslands. In the absence of grazing, tall grasses dominate the recovering steppe grassland and increase fire frequency and severity. Dwarf shrubs, such as <i>Artemisia</i>, are tolerant of grazing but decline when fires are too frequent or intense (62).</p>
<div data-bbox="269 1423 505 1654" data-label="Image"> </div> <p>22. Scots Pine Fire is expected to become more frequent and more severe under a hotter and drier climate in the boreal</p>	<div data-bbox="688 1423 924 1654" data-label="Image"> </div> <p>23. Dragonfly <i>Asiagomphus coreanus</i> is a dragonfly known from a single geographic location on the Korean Peninsula. The</p>	<div data-bbox="1097 1423 1333 1654" data-label="Image"> </div> <p>24. Grasshopper communities In high-altitude grasslands of the Western Ghats, India, planned burning is used to</p>

<p>forests of Siberia, which will alter species composition, forest area and carbon storage. In particular, shifts from boreal forest, dominated by Scots pine, to grass-dominated vegetation are expected (12).</p>	<p>population occurs in a freshwater stream inside the demilitarized zone (DMZ) between South Korea and North Korea. Military forces of both countries use fire to reduce vegetation for increased visibility. Forest fires negatively impact the preferred stream habitat of <i>A. coreanus</i>, but fires cannot be extinguished due to administrative restrictions. Forest reduction is ongoing and is particularly severe around this species' habitat (24).</p>	<p>create habitat for an endangered ungulate. But uniform burning has negative effects on grasshopper communities, which prefer patchier fires that facilitate rapid population recovery. 'Patchy' burning could be used to promote both the flagship ungulate species as well as species-rich grasshopper communities (106).</p>
<div data-bbox="277 869 516 1108" data-label="Image"> </div> <p>25. Orangutan Large fires in tropical forest of Indonesia are primarily associated with land clearing for agriculture, particularly oil-palm and paper-pulp plantations, and threaten one of the world's most biodiverse ecosystems and endangered species including orangutan (8).</p>	<div data-bbox="683 869 922 1108" data-label="Image"> </div> <p>26. Obligate seeding conifer The invasive grass, Gamba <i>Andropogon gayanus</i>, was introduced to the savannas of northern Australia as food for cattle. The high biomass of this invasive grass led to a switch from low-intensity surface fires to more intense fires that burn midstorey vegetation. Successive fires of high intensity have led to the decline of the obligate seeder <i>Callitris intratropica</i> through tree mortality. In turn, the reduction in canopy cover facilitates further invasion by grass. (107).</p>	<div data-bbox="1084 869 1323 1108" data-label="Image"> </div> <p>27. Banksias Up to 90% of native vegetation has been cleared for cereal cropping and sheep grazing in a southwest Australian biodiversity hotspot. Consequently, shrubland ecosystems comprised of <i>Banksia</i> species occur in fragmented landscapes dominated by small patches. Interactions between fragmentation, fire and a drying climate are projected to increase local extinctions of <i>Banksia</i> species, especially in small, isolated patches that cannot be colonized post-fire through dispersal. Intervals between</p>

		fires that are either too short or too long can place <i>Banksia</i> species at risk of local extinction (73).
 <p>28. Leadbeater's possum Temperate Mountain Ash forests of southeastern Australia provide habitat for species-rich animal and plant assemblages. The cumulative impacts of logging and extensive wildfires have removed large trees, placing populations of arboreal mammals that nest in old trees with hollows, such as Leadbeater's Possum (<i>Gymnobelideus leadbeateri</i>), at increased risk of extinction. In addition, high temperatures and droughts associated with climate change can directly increase mortality of large, hollow-bearing Mountain Ash trees and amplify the size and frequency of wildfires (48).</p>	 <p>29. Southern temperate forests Invasive plant and animal species interact with fire to slow forest regeneration and cause shifts in vegetation types in New Zealand. Humans introduced fire to temperate forests that previously experienced low fire frequency, while seed predation by invasive mammals and the reduction of avian pollination and seed dispersal halted post-fire vegetation succession. Highly-flammable invasive plants can also contribute to regime shifts via higher fire frequency and by altering plant-soil feedbacks. Temperate forest is gradually replaced by more fire-prone shrubland (55).</p>	 <p>30. Endemic Tasmanian pine Populations of the pine <i>Athrotaxis selaginoides</i> collapsed following the transition from Indigenous to European fire management in temperate forests of Tasmania, Australia. The species is now restricted to topographic refugia that offer protection from more severe fires (64). More recently, large fires associated with global climate change are impacting these refugia and targeted fire management is required to protect the species (64).</p>

References and Notes

1. T. He, B. B. Lamont, J. G. Pausas, Fire as a key driver of Earth's biodiversity. *Biol. Rev. Camb. Philos. Soc.* **94**, 1983–2010 (2019). [doi:10.1111/brv.12544](https://doi.org/10.1111/brv.12544) [Medline](#)
2. P. W. Rundel, M. T. K. Arroyo, R. M. Cowling, J. E. Keeley, B. B. Lamont, J. G. Pausas, P. Vargas, Fire and plant diversification in Mediterranean-climate regions. *Front. Plant Sci.* **9**, 851 (2018). [doi:10.3389/fpls.2018.00851](https://doi.org/10.3389/fpls.2018.00851) [Medline](#)
3. L. T. Kelly, L. Brotons, Using fire to promote biodiversity. *Science* **355**, 1264–1265 (2017). [doi:10.1126/science.aam7672](https://doi.org/10.1126/science.aam7672) [Medline](#)
4. J. G. Pausas, W. J. Bond, Alternative biome states in terrestrial ecosystems. *Trends Plant Sci.* **25**, 250–263 (2020). [doi:10.1016/j.tplants.2019.11.003](https://doi.org/10.1016/j.tplants.2019.11.003) [Medline](#)
5. M. S. Fletcher, T. Hall, A. N. Alexandra, The loss of an indigenous constructed landscape following British invasion of Australia: An insight into the deep human imprint on the Australian landscape. *Ambio* (2020). [doi:10.1007/s13280-020-01339-3](https://doi.org/10.1007/s13280-020-01339-3) [Medline](#)
6. S. Archibald, Managing the human component of fire regimes: Lessons from Africa. *Philos. Trans. R. Soc. London Ser. B* **371**, 20150346 (2016). [doi:10.1098/rstb.2015.0346](https://doi.org/10.1098/rstb.2015.0346) [Medline](#)
7. D. M. J. S. Bowman, J. Balch, P. Artaxo, W. J. Bond, M. A. Cochrane, C. M. D'Antonio, R. Defries, F. H. Johnston, J. E. Keeley, M. A. Krawchuk, C. A. Kull, M. Mack, M. A. Moritz, S. Pyne, C. I. Roos, A. C. Scott, N. S. Sodhi, T. W. Swetnam, R. Whittaker, The human dimension of fire regimes on Earth. *J. Biogeogr.* **38**, 2223–2236 (2011). [doi:10.1111/j.1365-2699.2011.02595.x](https://doi.org/10.1111/j.1365-2699.2011.02595.x) [Medline](#)
8. R. A. Chisholm, L. S. Wijedasa, T. Swinfield, The need for long-term remedies for Indonesia's forest fires. *Conserv. Biol.* **30**, 5–6 (2016). [doi:10.1111/cobi.12662](https://doi.org/10.1111/cobi.12662) [Medline](#)
9. J. Barlow, E. Berenguer, R. Carmenta, F. França, Clarifying Amazonia's burning crisis. *Glob. Change Biol.* **26**, 319–321 (2020). [doi:10.1111/gcb.14872](https://doi.org/10.1111/gcb.14872) [Medline](#)
10. F. S. Hu, P. E. Higuera, P. Duffy, M. L. Chipman, A. V. Rocha, A. M. Young, R. Kelly, M. C. Dietze, Arctic tundra fires: Natural variability and responses to climate change. *Front. Ecol. Environ.* **13**, 369–377 (2015). [doi:10.1890/150063](https://doi.org/10.1890/150063)
11. E. Whitman, M. A. Parisien, D. K. Thompson, M. D. Flannigan, Short-interval wildfire and drought overwhelm boreal forest resilience. *Sci. Rep.* **9**, 18796 (2019). [doi:10.1038/s41598-019-55036-7](https://doi.org/10.1038/s41598-019-55036-7) [Medline](#)
12. K. Barrett, R. Baxter, E. Kukavskaya, H. Balzter, E. Shvetsov, L. Buryak, Postfire recruitment failure in Scots pine forests of southern Siberia. *Remote Sens. Environ.* **237**, 111539 (2020). [doi:10.1016/j.rse.2019.111539](https://doi.org/10.1016/j.rse.2019.111539)
13. T. Schoennagel, J. K. Balch, H. Brenkert-Smith, P. E. Dennison, B. J. Harvey, M. A. Krawchuk, N. Mietkiewicz, P. Morgan, M. A. Moritz, R. Rasker, M. G. Turner, C. Whitlock, Adapt to more wildfire in western North American forests as climate changes. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 4582–4590 (2017). [doi:10.1073/pnas.1617464114](https://doi.org/10.1073/pnas.1617464114) [Medline](#)

14. M. M. Boer, V. Resco de Dios, R. A. Bradstock, Unprecedented burn area of Australian mega forest fires. *Nat. Clim. Chang.* **10**, 171–172 (2020). [doi:10.1038/s41558-020-0716-1](https://doi.org/10.1038/s41558-020-0716-1)
15. F. Moreira, D. Ascoli, H. Safford, M. A. Adams, J. M. Moreno, J. M. C. Pereira, F. X. Catry, J. Armesto, W. Bond, M. E. González, T. Curt, N. Koutsias, L. McCaw, O. Price, J. G. Pausas, E. Rigolot, S. Stephens, C. Tavsanoglu, V. R. Vallejo, B. W. Van Wilgen, G. Xanthopoulos, P. M. Fernandes, Wildfire management in Mediterranean-type regions: Paradigm change needed. *Environ. Res. Lett.* **15**, 011001 (2020). [doi:10.1088/1748-9326/ab541e](https://doi.org/10.1088/1748-9326/ab541e)
16. D. Twidwell, C. H. Bielski, R. Scholtz, S. D. Fuhlendorf, Advancing fire ecology in 21st century rangelands. *Rangeland Ecol. Manag.* (2020). [doi:10.1016/j.rama.2020.01.008](https://doi.org/10.1016/j.rama.2020.01.008)
17. N. Andela, D. C. Morton, L. Giglio, Y. Chen, G. R. van der Werf, P. S. Kasibhatla, R. S. DeFries, G. J. Collatz, S. Hantson, S. Kloster, D. Bachelet, M. Forrest, G. Lasslop, F. Li, S. Mangeon, J. R. Melton, C. Yue, J. T. Randerson, A human-driven decline in global burned area. *Science* **356**, 1356–1362 (2017). [doi:10.1126/science.aal4108](https://doi.org/10.1126/science.aal4108) [Medline](#)
18. J. R. Probert, C. L. Parr, R. M. Holdo, T. M. Anderson, S. Archibald, C. J. Courtney Mustaphi, A. P. Dobson, J. E. Donaldson, G. C. Hopcraft, G. P. Hempson, T. A. Morrison, C. M. Beale, Anthropogenic modifications to fire regimes in the wider Serengeti-Mara ecosystem. *Glob. Change Biol.* **25**, 3406–3423 (2019). [doi:10.1111/gcb.14711](https://doi.org/10.1111/gcb.14711) [Medline](#)
19. A. M. Gill, Fire and the Australian flora: A review. *Aust. For.* **38**, 4–25 (1975). [doi:10.1080/00049158.1975.10675618](https://doi.org/10.1080/00049158.1975.10675618)
20. R. G. Miller, R. Tangney, N. J. Enright, J. B. Fontaine, D. J. Merritt, M. K. J. Ooi, K. X. Ruthrof, B. P. Miller, Mechanisms of fire seasonality effects on plant populations. *Trends Ecol. Evol.* **34**, 1104–1117 (2019). [doi:10.1016/j.tree.2019.07.009](https://doi.org/10.1016/j.tree.2019.07.009) [Medline](#)
21. J. E. Keeley, J. G. Pausas, Distinguishing disturbance from perturbations in fire-prone ecosystems. *Int. J. Wildland Fire* **28**, 282–287 (2019). [doi:10.1071/WF18203](https://doi.org/10.1071/WF18203)
22. J. E. Keeley, W. J. Bond, R. A. Bradstock, J. G. Pausas, P. W. Rundel, *Fire in Mediterranean Ecosystems: Ecology, Evolution and Management* (Cambridge: Cambridge Univ. Press, 2012).
23. D. G. Nimmo, S. Avitabile, S. C. Banks, R. Bliege Bird, K. Callister, M. F. Clarke, C. R. Dickman, T. S. Doherty, D. A. Driscoll, A. C. Greenville, A. Haslem, L. T. Kelly, S. A. Kenny, J. J. Lahoz-Monfort, C. Lee, S. Leonard, H. Moore, T. M. Newsome, C. L. Parr, E. G. Ritchie, K. Schneider, J. M. Turner, S. Watson, M. Westbrooke, M. Wouters, M. White, A. F. Bennett, Animal movements in fire-prone landscapes. *Biol. Rev. Camb. Philos. Soc.* **94**, 981–998 (2019). [doi:10.1111/brv.12486](https://doi.org/10.1111/brv.12486) [Medline](#)
24. International Union for the Conservation of Nature (IUCN), “The IUCN red list of threatened species,” Version 2019-3 (IUCN, 2019); www.iucnredlist.org.
25. International Union for the Conservation of Nature (IUCN), “The IUCN red list of ecosystems,” 14 April 2020 (IUCN, 2020); <https://iucnrle.org/>.

26. B. A. Wintle, S. Legge, J. C. Z. Woinarski, After the megafires: What next for Australian wildlife? *Trends Ecol. Evol.* **35**, 753–757 (2020). [doi:10.1016/j.tree.2020.06.009](https://doi.org/10.1016/j.tree.2020.06.009) [Medline](#)
27. W. Steffen, J. Rockström, K. Richardson, T. M. Lenton, C. Folke, D. Liverman, C. P. Summerhayes, A. D. Barnosky, S. E. Cornell, M. Crucifix, J. F. Donges, I. Fetzer, S. J. Lade, M. Scheffer, R. Winkelmann, H. J. Schellnhuber, Trajectories of the Earth System in the Anthropocene. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 8252–8259 (2018). [doi:10.1073/pnas.1810141115](https://doi.org/10.1073/pnas.1810141115) [Medline](#)
28. D. M. J. S. Bowman, J. K. Balch, P. Artaxo, W. J. Bond, J. M. Carlson, M. A. Cochrane, C. M. D’Antonio, R. S. Defries, J. C. Doyle, S. P. Harrison, F. H. Johnston, J. E. Keeley, M. A. Krawchuk, C. A. Kull, J. B. Marston, M. A. Moritz, I. C. Prentice, C. I. Roos, A. C. Scott, T. W. Swetnam, G. R. van der Werf, S. J. Pyne, Fire in the Earth system. *Science* **324**, 481–484 (2009). [doi:10.1126/science.1163886](https://doi.org/10.1126/science.1163886) [Medline](#)
29. S. Archibald, C. E. R. Lehmann, J. L. Gómez-Dans, R. A. Bradstock, Defining pyromes and global syndromes of fire regimes. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 6442–6447 (2013). [doi:10.1073/pnas.1211466110](https://doi.org/10.1073/pnas.1211466110) [Medline](#)
30. B. Rogers, J. K. Balch, S. J. Goetz, C. E. R. Lehmann, M. Turetsky, Focus on changing fire regimes: Interactions with climate, ecosystems, and society. *Environ. Res. Lett.* **15**, 030201 (2018). [doi:10.1088/1748-9326/ab6d3a](https://doi.org/10.1088/1748-9326/ab6d3a)
31. S. Díaz, J. Settele, E. S. Brondízio, H. T. Ngo, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. R. Chowdhury, Y.-J. Shin, I. Visseren-Hamakers, K. J. Willis, C. N. Zayas, Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science* **366**, eaax3100 (2019). [doi:10.1126/science.aax3100](https://doi.org/10.1126/science.aax3100) [Medline](#)
32. R. A. Bradstock, A biogeographic model of fire regimes in Australia: Current and future implications. *Glob. Ecol. Biogeogr.* **19**, 145–158 (2010). [doi:10.1111/j.1466-8238.2009.00512.x](https://doi.org/10.1111/j.1466-8238.2009.00512.x)
33. E. Batllori, M. De Cáceres, L. Brotons, D. D. Ackerly, M. A. Moritz, F. Lloret, Compound fire-drought regimes promote ecosystem transitions in Mediterranean ecosystems. *J. Ecol.* **107**, 1187–1198 (2019). [doi:10.1111/1365-2745.13115](https://doi.org/10.1111/1365-2745.13115)
34. P. M. Brando, L. Paolucci, C. C. Ummenhofer, E. M. Ordway, H. Hartmann, M. E. Cattau, L. Rattis, V. Medjibe, M. T. Coe, J. Balch, Droughts, wildfires, and forest carbon cycling: A pantropical synthesis. *Annu. Rev. Earth Planet. Sci.* **47**, 555–581 (2019). [doi:10.1146/annurev-earth-082517-010235](https://doi.org/10.1146/annurev-earth-082517-010235)
35. T. A. Fairman, C. R. Nitschke, L. T. Bennett, Too much, too soon? A review of the effects of increasing wildfire frequency on tree mortality and regeneration in temperate eucalypt forests. *Int. J. Wildland Fire* **25**, 831–848 (2016). [doi:10.1071/WF15010](https://doi.org/10.1071/WF15010)
36. M. G. Turner, K. H. Braziunas, W. D. Hansen, B. J. Harvey, Short-interval severe fire erodes the resilience of subalpine lodgepole pine forests. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 11319–11328 (2019). [doi:10.1073/pnas.1902841116](https://doi.org/10.1073/pnas.1902841116) [Medline](#)

37. S. R. Holden, B. M. Rogers, K. K. Treseder, J. T. Randerson, Fire severity influences the response of soil microbes to a boreal forest fire. *Environ. Res. Lett.* **11**, 035004 (2016). [doi:10.1088/1748-9326/11/3/035004](https://doi.org/10.1088/1748-9326/11/3/035004)
38. K. Joly, P. A. Duffy, T. S. Rupp, Simulating the effects of climate change on fire regimes in Arctic biomes: Implications for caribou and moose habitat. *Ecosphere* **3**, 1–18 (2012). [doi:10.1890/ES12-00012.1](https://doi.org/10.1890/ES12-00012.1)
39. M. A. Cochrane, C. P. Barber, Climate change, human land use and future fires in the Amazon. *Glob. Change Biol.* **15**, 601–612 (2009). [doi:10.1111/j.1365-2486.2008.01786.x](https://doi.org/10.1111/j.1365-2486.2008.01786.x)
40. L. N. Paolucci, J. H. Schoereder, P. M. Brando, A. N. Andersen, Fire-induced forest transition to derived savannas: Cascading effects on ant communities. *Biol. Conserv.* **214**, 295–302 (2017). [doi:10.1016/j.biocon.2017.08.020](https://doi.org/10.1016/j.biocon.2017.08.020)
41. R. B. de Andrade, J. K. Balch, J. Y. O. Carreira, P. M. Brando, A. V. L. Freitas, The impacts of recurrent fires on diversity of fruit-feeding butterflies in a south-eastern Amazon forest. *J. Trop. Ecol.* **33**, 22–32 (2017). [doi:10.1017/S0266467416000559](https://doi.org/10.1017/S0266467416000559)
42. M. P. Veldhuis, M. E. Ritchie, J. O. Ogutu, T. A. Morrison, C. M. Beale, A. B. Estes, W. Mwakilema, G. O. Ojwang, C. L. Parr, J. Probert, P. W. Wargute, J. G. C. Hopcraft, H. Olf, Cross-boundary human impacts compromise the Serengeti-Mara ecosystem. *Science* **363**, 1424–1428 (2019). [doi:10.1126/science.aav0564](https://doi.org/10.1126/science.aav0564) [Medline](#)
43. R. C. R. Abreu, W. A. Hoffmann, H. L. Vasconcelos, N. A. Pilon, D. R. Rossatto, G. Durigan, The biodiversity cost of carbon sequestration in tropical savanna. *Sci. Adv.* **3**, e1701284 (2017). [doi:10.1126/sciadv.1701284](https://doi.org/10.1126/sciadv.1701284) [Medline](#)
44. D. A. Driscoll, M. Bode, R. A. Bradstock, D. A. Keith, T. D. Penman, O. F. Price, Resolving future fire management conflicts using multicriteria decision making. *Conserv. Biol.* **30**, 196–205 (2016). [doi:10.1111/cobi.12580](https://doi.org/10.1111/cobi.12580) [Medline](#)
45. A. D. Syphard, T. J. Brennan, J. E. Keeley, Drivers of chaparral type conversion to herbaceous vegetation in coastal Southern California. *Divers. Distrib.* **25**, 90–101 (2019). [doi:10.1111/ddi.12827](https://doi.org/10.1111/ddi.12827)
46. F. Moreira, G. Pe'er, Agricultural policy can reduce wildfires. *Science* **359**, 1001 (2018). [Medline](#)
47. A. Regos, V. Hermoso, M. D'Amen, A. Guisan, L. Brotons, Trade-offs and synergies between bird conservation and wildfire suppression in the face of global change. *J. Appl. Ecol.* **55**, 2181–2192 (2018). [doi:10.1111/1365-2664.13182](https://doi.org/10.1111/1365-2664.13182)
48. D. B. Lindenmayer, C. Sato, Hidden collapse is driven by fire and logging in a socioecological forest ecosystem. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 5181–5186 (2018). [doi:10.1073/pnas.1721738115](https://doi.org/10.1073/pnas.1721738115) [Medline](#)
49. G. T. Pecl, M. B. Araújo, J. D. Bell, J. Blanchard, T. C. Bonebrake, I.-C. Chen, T. D. Clark, R. K. Colwell, F. Danielsen, B. Evengård, L. Falconi, S. Ferrier, S. Frusher, R. A. Garcia, R. B. Griffis, A. J. Hobday, C. Janion-Scheepers, M. A. Jarzyna, S. Jennings, J. Lenoir, H. I. Linnetved, V. Y. Martin, P. C. McCormack, J. McDonald, N. J. Mitchell, T. Mustonen, J. M. Pandolfi, N. Pettorelli, E. Popova, S. A. Robinson, B. R. Scheffers, J. D.

- Shaw, C. J. B. Sorte, J. M. Strugnell, J. M. Sunday, M.-N. Tuanmu, A. Vergés, C. Villanueva, T. Wernberg, E. Wapstra, S. E. Williams, Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355**, eaai9214 (2017). [doi:10.1126/science.aai9214](https://doi.org/10.1126/science.aai9214) [Medline](#)
50. J. G. Pausas, J. E. Keeley, Abrupt climate-independent fire regime changes. *Ecosystems (N. Y.)* **17**, 1109–1120 (2014). [doi:10.1007/s10021-014-9773-5](https://doi.org/10.1007/s10021-014-9773-5)
 51. M. L. Brooks, C. M. D’Antonio, D. M. Richardson, J. B. Grace, J. Keeley, J. M. DiTOMASO, R. J. Hobbs, M. Pellant, D. Pyke, Effects of invasive alien plants on fire regimes. *Bioscience* **54**, 677–688 (2004). [doi:10.1641/0006-3568\(2004\)054\[0677:EOIAP0\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0677:EOIAP0]2.0.CO;2)
 52. E. J. Fusco, J. T. Finn, J. K. Balch, R. C. Nagy, B. A. Bradley, Invasive grasses increase fire occurrence and frequency across US ecoregions. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 23594–23599 (2019). [doi:10.1073/pnas.1908253116](https://doi.org/10.1073/pnas.1908253116) [Medline](#)
 53. C. Riginos, T. A. Monaco, K. E. Veblen, K. Gunnell, E. Thacker, D. Dahlgren, T. Messmer, Potential for post-fire recovery of greater sage-grouse habitat. *Ecosphere* **10**, e02870 (2019). [doi:10.1002/ecs2.2870](https://doi.org/10.1002/ecs2.2870)
 54. C. N. Foster, S. C. Banks, G. J. Cary, C. N. Johnson, D. B. Lindenmayer, L. E. Valentine, Animals as agents in fire regimes. *Trends Ecol. Evol.* **35**, 346–356 (2020). [doi:10.1016/j.tree.2020.01.002](https://doi.org/10.1016/j.tree.2020.01.002) [Medline](#)
 55. G. L. W. Perry, J. M. Wilmschurst, J. Ogden, N. J. Enright, Exotic mammals and invasive plants alter fire-related thresholds in southern temperate forested landscapes. *Ecosystems* **18**, 1290–1305 (2015). [doi:10.1007/s10021-015-9898-1](https://doi.org/10.1007/s10021-015-9898-1)
 56. B. A. Hradsky, Conserving Australia’s threatened native mammals in predator-invaded, fire-prone landscapes. *Wildl. Res.* **47**, 1–15 (2020). [doi:10.1071/WR19027](https://doi.org/10.1071/WR19027)
 57. C. N. Johnson, L. D. Prior, S. Archibald, H. M. Poulos, A. M. Barton, G. J. Williamson, D. M. J. S. Bowman, Can trophic rewilding reduce the impact of fire in a more flammable world? *Philos. Trans. R. Soc. London B.* **373**, 20170443 (2018). [doi:10.1098/rstb.2017.0443](https://doi.org/10.1098/rstb.2017.0443) [Medline](#)
 58. C. M. Ryan, R. J. Hobbs, L. E. Valentine, Bioturbation by a reintroduced digging mammal reduces fuel loads in an urban reserve. *Ecol. Appl.* **30**, e02018 (2020). [doi:10.1002/eap.2018](https://doi.org/10.1002/eap.2018) [Medline](#)
 59. M. A. Moritz, E. Batllori, R. A. Bradstock, A. M. Gill, J. Handmer, P. F. Hessburg, J. Leonard, S. McCaffrey, D. C. Odion, T. Schoennagel, A. D. Syphard, Learning to coexist with wildfire. *Nature* **515**, 58–66 (2014). [doi:10.1038/nature13946](https://doi.org/10.1038/nature13946) [Medline](#)
 60. A. D. Syphard, J. E. Keeley, A. H. Pfaff, K. Ferschweiler, Human presence diminishes the importance of climate in driving fire activity across the United States. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 13750–13755 (2017). [doi:10.1073/pnas.1713885114](https://doi.org/10.1073/pnas.1713885114) [Medline](#)
 61. C. A. Nobre, G. Sampaio, L. S. Borma, J. C. Castilla-Rubio, J. S. Silva, M. Cardoso, Land-use and climate change risks in the Amazon and the need of a novel sustainable development paradigm. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 10759–10768 (2016). [doi:10.1073/pnas.1605516113](https://doi.org/10.1073/pnas.1605516113) [Medline](#)

62. A. Brinkert, N. Hölzel, T. V. Sidorova, J. Kamp, Spontaneous steppe restoration on abandoned cropland in Kazakhstan: Grazing affects successional pathways. *Biodivers. Conserv.* **25**, 2543–2561 (2016). [doi:10.1007/s10531-015-1020-7](https://doi.org/10.1007/s10531-015-1020-7)
63. M. J. Liebmann, J. Farella, C. I. Roos, A. Stack, S. Martini, T. W. Swetnam, Native American depopulation, reforestation, and fire regimes in the Southwest United States, 1492-1900 CE. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E696–E704 (2016). [doi:10.1073/pnas.1521744113](https://doi.org/10.1073/pnas.1521744113) [Medline](#)
64. A. Holz, S. W. Wood, C. Ward, T. T. Veblen, D. M. J. S. Bowman, Population collapse and retreat to fire refugia of the Tasmanian endemic conifer *Athrotaxis selaginoides* following the transition from Aboriginal to European fire management. *Glob. Change Biol.* **26**, 3108–3121 (2020). [doi:10.1111/gcb.15031](https://doi.org/10.1111/gcb.15031) [Medline](#)
65. R. E. Keane, D. McKenzie, D. A. Falk, E. A. H. Smithwick, C. Miller, L.-K. B. Kellogg, Representing climate, disturbance, and vegetation interactions in landscape models. *Ecol. Modell.* **309–310**, 33–47 (2015). [doi:10.1016/j.ecolmodel.2015.04.009](https://doi.org/10.1016/j.ecolmodel.2015.04.009)
66. M. C. Urban, G. Bocedi, A. P. Hendry, J.-B. Mihoub, G. Pe'er, A. Singer, J. R. Bridle, L. G. Crozier, L. De Meester, W. Godsoe, A. Gonzalez, J. J. Hellmann, R. D. Holt, A. Huth, K. Johst, C. B. Krug, P. W. Leadley, S. C. F. Palmer, J. H. Pantel, A. Schmitz, P. A. Zollner, J. M. J. Travis, Improving the forecast for biodiversity under climate change. *Science* **353**, aad8466 (2016). [doi:10.1126/science.aad8466](https://doi.org/10.1126/science.aad8466) [Medline](#)
67. P. H. Verburg, J. A. Dearing, J. G. Dyke, S. Leeuw, S. Seitzinger, W. Steffen, J. Syvitski, Methods and approaches to modelling the Anthropocene. *Glob. Environ. Change* **39**, 328–340 (2016). [doi:10.1016/j.gloenvcha.2015.08.007](https://doi.org/10.1016/j.gloenvcha.2015.08.007)
68. C. M. Beale, C. J. Courtney Mustaphi, T. A. Morrison, S. Archibald, T. M. Anderson, A. P. Dobson, J. E. Donaldson, G. P. Hempson, J. Probert, C. L. Parr, Pyrodiversity interacts with rainfall to increase bird and mammal richness in African savannas. *Ecol. Lett.* **21**, 557–567 (2018). [doi:10.1111/ele.12921](https://doi.org/10.1111/ele.12921) [Medline](#)
69. L. C. Ponisio, K. Wilkin, L. K. M'Gonigle, K. Kulhanek, L. Cook, R. Thorp, T. Griswold, C. Kremen, Pyrodiversity begets plant-pollinator community diversity. *Glob. Change Biol.* **22**, 1794–1808 (2016). [doi:10.1111/gcb.13236](https://doi.org/10.1111/gcb.13236) [Medline](#)
70. M. W. Tingley, V. Ruiz-Gutiérrez, R. L. Wilkerson, C. A. Howell, R. B. Siegel, Pyrodiversity promotes avian diversity over the decade following forest fire. *Proc. Biol. Sci.* **283**, 20161703 (2016). [doi:10.1098/rspb.2016.1703](https://doi.org/10.1098/rspb.2016.1703) [Medline](#)
71. P. D. Bentley, T. D. Penman, Is there an inherent conflict in managing fire for people and conservation? *Int. J. Wildland Fire* **26**, 455–468 (2017). [doi:10.1071/WF16150](https://doi.org/10.1071/WF16150)
72. J. Henzler, H. Weise, N. J. Enright, S. Zander, B. Tietjen, A squeeze in the suitable fire interval: Simulating the persistence of fire-killed plants in a Mediterranean-type ecosystem under drier conditions. *Ecol. Modell.* **389**, 41–49 (2018). [doi:10.1016/j.ecolmodel.2018.10.010](https://doi.org/10.1016/j.ecolmodel.2018.10.010)
73. A. I. T. Tulloch, J.-B. Pichancourt, C. R. Gosper, A. Sanders, I. Chadès, Fire management strategies to maintain species population processes in a fragmented landscape of fire-interval extremes. *Ecol. Appl.* **26**, 2175–2189 (2016). [doi:10.1002/eap.1362](https://doi.org/10.1002/eap.1362) [Medline](#)

74. H. M. Regan, C. I. Bohórquez, D. A. Keith, T. J. Regan, K. E. Anderson, Implications of different population model structures for management of threatened plants. *Conserv. Biol.* **31**, 459–468 (2017). [doi:10.1111/cobi.12831](https://doi.org/10.1111/cobi.12831) [Medline](#)
75. T. C. Bonebrake, A. D. Syphard, J. Franklin, K. E. Anderson, H. R. Akçakaya, T. Mizerek, C. Winchell, H. M. Regan, Fire management, managed relocation, and land conservation options for long-lived obligate seeding plants under global changes in climate, urbanization, and fire regime. *Conserv. Biol.* **28**, 1057–1067 (2014). [doi:10.1111/cobi.12253](https://doi.org/10.1111/cobi.12253) [Medline](#)
76. S. Gómez-González, C. Torres-Díaz, C. Bustos-Schindler, E. Gianoli, Anthropogenic fire drives the evolution of seed traits. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 18743–18747 (2011). [doi:10.1073/pnas.1108863108](https://doi.org/10.1073/pnas.1108863108) [Medline](#)
77. M. C. Castellanos, S. C. González-Martínez, J. G. Pausas, Field heritability of a plant adaptation to fire in heterogeneous landscapes. *Mol. Ecol.* **24**, 5633–5642 (2015). [doi:10.1111/mec.13421](https://doi.org/10.1111/mec.13421) [Medline](#)
78. S. Archibald, C. E. R. Lehmann, C. M. Belcher, W. J. Bond, R. A. Bradstock, A.-L. Daniau, K. G. Dexter, E. J. Forrester, M. Greve, T. He, S. I. Higgins, W. A. Hoffmann, B. B. Lamont, D. J. McGlenn, G. R. Moncrieff, C. P. Osborne, J. G. Pausas, O. Price, B. S. Ripley, B. M. Rogers, D. W. Schwilk, M. F. Simon, M. R. Turetsky, G. R. Van der Werf, A. E. Zanne, Biological and geophysical feedbacks with fire in the Earth System. *Environ. Res. Lett.* **13**, 033003 (2017). [doi:10.1088/1748-9326/aa9ead](https://doi.org/10.1088/1748-9326/aa9ead)
79. D. M. J. S. Bowman, G. L. W. Perry, S. I. Higgins, C. N. Johnson, S. D. Fuhlendorf, B. P. Murphy, Pyrodiversity is the coupling of biodiversity and fire regimes in food webs. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **371**, 20150169 (2016). [doi:10.1098/rstb.2015.0169](https://doi.org/10.1098/rstb.2015.0169) [Medline](#)
80. T. A. Spies, E. White, A. Ager, J. D. Kline, J. P. Bolte, E. K. Platt, K. A. Olsen, R. J. Pabst, A. M. G. Barros, J. D. Bailey, S. Charnley, A. T. Morzillo, J. Koch, M. M. Steen-Adams, P. H. Singleton, J. Sulzman, C. Schwartz, B. Csuti, Using an agent-based model to examine forest management outcomes in a fire-prone landscape in Oregon, USA. *Ecol. Soc.* **22**, art25 (2017). [doi:10.5751/ES-08841-220125](https://doi.org/10.5751/ES-08841-220125)
81. G. Boisramé, S. Thompson, B. Collins, S. Stephens, Managed wildfire effects on forest resilience and water in the Sierra Nevada. *Ecosystems (N. Y.)* **20**, 717–732 (2017). [doi:10.1007/s10021-016-0048-1](https://doi.org/10.1007/s10021-016-0048-1)
82. S. L. Stephens, L. N. Kobziar, B. M. Collins, R. Davis, P. Z. Fulé, W. Gaines, J. Ganey, J. M. Guldin, P. F. Hessburg, K. Hiers, S. Hoagland, J. J. Keane, R. E. Masters, A. E. McKellar, W. Montague, M. North, T. A. Spies, Is fire “for the birds”? How two rare species influence fire management across the US. *Front. Ecol. Environ.* **17**, 391–399 (2019). [doi:10.1002/fee.2076](https://doi.org/10.1002/fee.2076)
83. A. Morton, “Dinosaur trees: Firefighters save endangered Wollemi pines from NSW bushfires,” *The Guardian*, 15 January 2020; <https://www.theguardian.com/australia-news/2020/jan/15/dinosaur-trees-firefighters-save-endangered-wollemi-pines-from-nsw-bushfires>.

84. R. E. Keane, Managing wildfire for whitebark pine ecosystem restoration in western North America. *Forests* **9**, 648 (2018). [doi:10.3390/f9100648](https://doi.org/10.3390/f9100648)
85. G. Granath, P. A. Moore, M. C. Lukenbach, J. M. Waddington, Mitigating wildfire carbon loss in managed northern peatlands through restoration. *Sci. Rep.* **6**, 28498 (2016). [doi:10.1038/srep28498](https://doi.org/10.1038/srep28498) [Medline](#)
86. T. E. Lovejoy, C. Nobre, Amazon tipping point: Last chance for action. *Sci. Adv.* **5**, eaba2949 (2019). [doi:10.1126/sciadv.aba2949](https://doi.org/10.1126/sciadv.aba2949) [Medline](#)
87. I. P. J. Smit, G. P. Asner, N. Govender, N. R. Vaughn, B. W. van Wilgen, An examination of the potential efficacy of high-intensity fires for reversing woody encroachment in savannas. *J. Appl. Ecol.* **53**, 1623–1633 (2016). [doi:10.1111/1365-2664.12738](https://doi.org/10.1111/1365-2664.12738)
88. C. M. Sgrò, A. J. Lowe, A. A. Hoffmann, Building evolutionary resilience for conserving biodiversity under climate change. *Evol. Appl.* **4**, 326–337 (2011). [doi:10.1111/j.1752-4571.2010.00157.x](https://doi.org/10.1111/j.1752-4571.2010.00157.x) [Medline](#)
89. A. J. H. Meddens, C. A. Kolden, J. A. Lutz, A. M. S. Smith, C. A. Cansler, J. T. Abatzoglou, G. W. Meigs, W. M. Downing, M. A. Krawchuk, Fire refugia: What are they, and why do they matter for global change? *Bioscience* **68**, 944–954 (2018). [doi:10.1093/biosci/biy103](https://doi.org/10.1093/biosci/biy103)
90. O. D. Bassett, L. D. Prior, C. M. Slijkerman, D. Jamieson, D. M. J. S. Bowman, Aerial sowing stopped the loss of alpine ash (*Eucalyptus delegatensis*) forests burnt by three short-interval fires in the Alpine National Park, Victoria, Australia. *For. Ecol. Manage.* **342**, 39–48 (2015). [doi:10.1016/j.foreco.2015.01.008](https://doi.org/10.1016/j.foreco.2015.01.008)
91. F. K. Lake, A. C. Christianson, “Indigenous fire stewardship,” in *Encyclopedia of Wildfires and Wildland-Urban Interface (WUI) Fires*, Manzello S, Ed. (Springer, 2019); https://doi.org/10.1007/978-3-319-51727-8_225-1.
92. R. Bliege Bird, D. Nimmo, Restore the lost ecological functions of people. *Nat. Ecol. Evol.* **2**, 1050–1052 (2018). [doi:10.1038/s41559-018-0576-5](https://doi.org/10.1038/s41559-018-0576-5) [Medline](#)
93. J. Mistry, B. A. Bilbao, A. Berardi, Community owned solutions for fire management in tropical ecosystems: Case studies from Indigenous communities of South America. *Philos. Trans. R. Soc. London B.* **371**, 20150174 (2016). [doi:10.1098/rstb.2015.0174](https://doi.org/10.1098/rstb.2015.0174) [Medline](#)
94. J. R. Welch, E. S. Brondízio, S. S. Hetrick, C. E. Coimbra Jr., Indigenous burning as conservation practice: Neotropical savanna recovery amid agribusiness deforestation in Central Brazil. *PLOS ONE* **8**, e81226 (2013). [doi:10.1371/journal.pone.0081226](https://doi.org/10.1371/journal.pone.0081226) [Medline](#)
95. X. Cui, M. A. Alam, G. L. W. Perry, A. M. Paterson, S. V. Wyse, T. J. Curran, Green firebreaks as a management tool for wildfires: Lessons from China. *J. Environ. Manage.* **233**, 329–336 (2019). [doi:10.1016/j.jenvman.2018.12.043](https://doi.org/10.1016/j.jenvman.2018.12.043) [Medline](#)
96. A. D. Barnosky, E. A. Hadly, P. Gonzalez, J. Head, P. D. Polly, A. M. Lawing, J. T. Eronen, D. D. Ackerly, K. Alex, E. Biber, J. Blois, J. Brashares, G. Ceballos, E. Davis, G. P. Dietl, R. Dirzo, H. Doremus, M. Fortelius, H. W. Greene, J. Hellmann, T. Hickler, S. T. Jackson, M. Kemp, P. L. Koch, C. Kremen, E. L. Lindsey, C. Looy, C. R. Marshall, C. Mendenhall, A. Mulch, A. M. Mychajliw, C. Nowak, U. Ramakrishnan, J. Schnitzler, K. Das Shrestha, K. Solari, L. Stegner, M. A. Stegner, N. C. Stenseth, M. H. Wake, Z.

- Zhang, Merging paleobiology with conservation biology to guide the future of terrestrial ecosystems. *Science* **355**, eaah4787 (2017). [doi:10.1126/science.aah4787](https://doi.org/10.1126/science.aah4787) [Medline](#)
97. J. G. Pausas, J. E. Keeley, Wildfires as an ecosystem service. *Front. Ecol. Environ.* **17**, 289–295 (2019). [doi:10.1002/fee.2044](https://doi.org/10.1002/fee.2044)
98. Conservation Biology Institute, The Nature Conservancy, Paradise Parks and Recreation, “Paradise nature-based fire risk resilience project” (Corvallis, OR, 2020).
99. Convention on Biological Diversity, “Preparations for the post-2020 biodiversity framework” (CBD, 2020); <https://www.cbd.int/conferences/post2020>.
100. B. W. van Wilgen, N. Govender, I. P. J. Smit, S. MacFadyen, The ongoing development of a pragmatic and adaptive fire management policy in a large African savanna protected area. *J. Environ. Manage.* **132**, 358–368 (2014). [doi:10.1016/j.jenvman.2013.11.003](https://doi.org/10.1016/j.jenvman.2013.11.003) [Medline](#)
101. A. R. Templeton, H. Brazeal, J. L. Neuwald, The transition from isolated patches to a metapopulation in the eastern collared lizard in response to prescribed fires. *Ecology* **92**, 1736–1747 (2011). [doi:10.1890/10-1994.1](https://doi.org/10.1890/10-1994.1) [Medline](#)
102. V. Vandvik, J. P. Töpper, Z. Cook, M. I. Daws, E. Heegaard, I. E. Måren, L. G. Velle, Management-driven evolution in a domesticated ecosystem. *Biol. Lett.* **10**, 20131082 (2014). [10.1098/rsbl.2013.1082](https://doi.org/10.1098/rsbl.2013.1082) [Medline](#)
103. B. Chergui, R. C. Rodríguez-Caro, E. Graciá, S. Fahd, X. Santos, Population density of the spur-thighed tortoise *Testudo graeca* declines after fire in north-western Africa. *PLOS ONE* **14**, e0220969 (2019). [doi:10.1371/journal.pone.0220969](https://doi.org/10.1371/journal.pone.0220969) [Medline](#)
104. G. Walters, Customary fire regimes and vegetation structure in Gabon’s Bateke Plateaux. *Hum. Ecol.* **40**, 943–955 (2011). [doi:10.1007/s10745-012-9536-x](https://doi.org/10.1007/s10745-012-9536-x)
105. J. D. M. White, S. L. Jack, M. T. Hoffman, J. Puttick, D. Bonora, V. Visser, E. C. February, Collapse of an iconic conifer: Long-term changes in the demography of *Widdringtonia cedarbergensis* using repeat photography. *BMC Ecol.* **16**, 53 (2016). [doi:10.1186/s12898-016-0108-6](https://doi.org/10.1186/s12898-016-0108-6) [Medline](#)
106. D. Bhaskar, P. S. Easa, K. A. Sreejith, J. Skejo, A. Hochkirch, Large scale burning for a threatened ungulate in a biodiversity hotspot is detrimental for grasshoppers (Orthoptera: Caelifera). *Biodivers. Conserv.* **28**, 3221–3237 (2019). [doi:10.1007/s10531-019-01816-6](https://doi.org/10.1007/s10531-019-01816-6)
107. D. M. Bowman, H. J. MacDermott, S. C. Nichols, B. P. Murphy, A grass-fire cycle eliminates an obligate-seeding tree in a tropical savanna. *Ecol. Evol.* **4**, 4185–4194 (2014). [doi:10.1002/ece3.1285](https://doi.org/10.1002/ece3.1285) [Medline](#)