

CORRECTED PROOF

## Rapid Communication

**First records of *Amaranthus palmeri*, a new emerging weed in southern Africa with further notes on other poorly known alien amaranths in the continent**Alexander P. Sukhorukov<sup>1,2,\*</sup>, Maria Kushunina<sup>3</sup>, Carl F. Reinhardt<sup>4,5</sup>, Hugo Bezuidenhout<sup>6,7</sup> and Barend J. Vorster<sup>5</sup><sup>1</sup>Department of Higher Plants, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russian Federation<sup>2</sup>Laboratory Herbarium (TK), Tomsk State University, Lenin St. 36, 634050, Tomsk, Russian Federation<sup>3</sup>Department of Plant Physiology, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russian Federation<sup>4</sup>Unit for Environmental Sciences and Management, School of Geo- and Spatial Sciences, North-West University, Potchefstroom, Republic of South Africa<sup>5</sup>Department of Plant and Soil Sciences, University of Pretoria, Pretoria, Republic of South Africa<sup>6</sup>Applied Behavioral Ecology and Ecosystem Research Unit, UNISA, Private Bag X6, Florida 1717, Republic of South Africa<sup>7</sup>Scientific Services, Kimberley, SANParks, PO Box 110040, Hadison Park, Kimberley 8306, Republic of South AfricaAuthor e-mails: [suchor@mail.ru](mailto:suchor@mail.ru) (APS), [mkushunina@gmail.com](mailto:mkushunina@gmail.com) (MK), [37079573@nwu.ac.za](mailto:37079573@nwu.ac.za) (CFR), [dr.charlie.reinhardt@gmail.com](mailto:dr.charlie.reinhardt@gmail.com) (CFR), [hugo.bez24@gmail.com](mailto:hugo.bez24@gmail.com), [hugo.bezuidenhout@sanparks.org](mailto:hugo.bezuidenhout@sanparks.org) (HB), [juan.vorster@fabi.up.ac.za](mailto:juan.vorster@fabi.up.ac.za) (BJV)

\*Corresponding author

**Citation:** Sukhorukov AP, Kushunina M, Reinhardt CF, Bezuidenhout H, Vorster BJ (2021) First records of *Amaranthus palmeri*, a new emerging weed in southern Africa with further notes on other poorly known alien amaranths in the continent. *BioInvasions Records* 10 (in press)

**Received:** 4 August 2020**Accepted:** 3 October 2020**Published:** 1 December 2020**Handling editor:** Giuseppe Brundu**Thematic editor:** Stelios Katsanevakis**Copyright:** © Sukhorukov et al.This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International - CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

OPEN ACCESS

**Abstract**

*Amaranthus palmeri* is native to Mexico and the south-eastern parts of the USA, and is reported as alien in subtropical regions of the Old World. Previous records from Africa were from the northern parts of the continent. This species was first found in South Africa in March 2018 with further records in different regions of the country as well as in northern Botswana in March 2020. We consider it as naturalized weed which invades both ruderal and segetal plant communities. Mechanical control of *A. palmeri* plants may be effective to minimize its spreading, because at the flowering stage these tend to be taller compared to other amaranths. Those species of the genus alien in Africa are discussed in terms of their distribution and possible invasive status, particularly *A. dubius* and *A. standleyanus*.

**Key words:** Amaranths, naturalized species, tropical Africa, weeds**Introduction**

The order Caryophyllales Berchtold & J.Presl encompasses ~ 6% of all flowering plants and includes at least 30 families with the major diversity centers in hot and cold deserts of the World (Brockington et al. 2009; Hernández-Ledesma et al. 2015). Some members from different families of the order became noxious weeds in the tropics and temperate regions as a stable component of diverse disturbed plant communities (see e.g., Esser et al. 2013; Sukhorukov 2014). Only a minor part of them are cultivated plants. Among them, many Cactaceae provide the most negative impact on the landscapes in arid regions of Africa (Novoa et al. 2014; Shackleton 2017; van Wilgen et al. 2020). Other well-known intentionally introduced species and naturalized in the continent are some shrubby Chenopodiaceae s.str., or Amaranthaceae s.l. (e.g., *Atriplex nummularia* Lindl., *A. canescens*

(Pursh) Nutt.) used as forage plants in the deserts (Barnard et al. 1992; Heshmati and Squires 2013). However, the vast majority of naturalized or invasive Caryophyllales are accidentally introduced plants. Recently, *Sesuvium verrucosum* Raf. (Aizoaceae), *Mollugo verticillata* L. (Molluginaceae), *Bassia indica* (Wight) A.J.Scott (Amaranthaceae s.l.), and *Oxygonum sinuatum* (Hochst. & Steud.) Dammer (Polygonaceae) were described as naturalized or invasive xenophytes in different regions of continental Africa and Macaronesia (Sukhorukov et al. 2017, 2018a, 2019).

Many amaranths are widespread aliens that can be competitive weeds in both ruderal and segetal plant communities (Costea et al. 2004). In Europe, *Amaranthus powellii* S.Watson is a recently established neophyte invading ruderal habitats and agricultural areas of Europe (Forman 2003; Sukhorukov 2011) and foothills of East Kazakhstan (Sukhorukov 2009). Another well-established species in southern Europe and eastern Mediterranean is *A. palmeri* S.Watson (Zohary 1966; di Castri et al. 2012; Iamónico 2015; Sukhorukov pers. obs. in the Mediterranean part of Israel), a dioecious amaranth originating from Mexico and south-eastern part of the USA (Sauer 1955; Ward et al. 2013). In Africa, this species was recently reported in Egypt (Boulos 1995) and Tunisia (Iamónico and El Mokni 2017). Further records were reported in Ethiopia (APD 2019) and Morocco (EPPO 2020+). In Morocco, it is included in the quarantine pest list (EPPO 2020+). In the present article, we report the first documented records of *A. palmeri* in South Africa and Botswana, with further notes on the most successful amaranths in tropical Africa.


## Materials and methods

### *Study species*

*Amaranthus palmeri* (Palmer amaranth, coded as AMAPA in the EPPO Global Database: EPPO 2020+) is an annual herb with C<sub>4</sub> photosynthetic pathway. The most distinct characteristics of this species are tall stems up to 300 cm high; one type (male or female) of flower in each individual (plants are dioecious); large bracteate synflorescences reaching 60 cm in length; mucronate bracts longer than perianth; 5 spatulate perianth segments with clearly subulate outer segments; flowers with 5 stamens (male individuals) and 2 pistils (female); fruits dehiscent with smooth pericarp (Sauer 1955). A detailed morphological description and taxonomy of *A. palmeri* can be obtained from different references (e.g., Sauer 1955; Wetzal et al. 1999; Ward et al. 2013; Iamónico and El Mokni 2017). The species can thrive under different climatic conditions, but requires full sun and high temperatures (26–35 °C) that enable rapid growth (Ward et al. 2013).

### *Data sources and analysis*

The weed composition in Republic of South Africa has been investigated by Carl F. Reinhardt during the last three decades. The studies of Alexander



**Figure 1.** Locations of the new records of *Amaranthus palmeri* in southern Africa.


P. Sukhorukov on invasive Caryophyllales were carried out in 2009+ in different countries including Egypt (2009), Cape Verde (2015+), Namibia (2017, 2018), South Africa (2018), Tanzania (2019), Zimbabwe, Zambia, and Botswana (2020). The material collected during the field studies is kept in the herbaria G, K, MW, PRE, and W. During the last trip by APS in the first half of March 2020, the area under investigation included the surroundings of Victoria Falls (Zimbabwe), Livingstone (Zambia), and Kasane (Botswana) towns. The records of *Amaranthus palmeri* were mapped using [simplemapp.net](http://simplemapp.net) online tool.

## Results

### *Records and population size*

In South Africa, *Amaranthus palmeri* (Figure 1) was recorded for the first time in a farm in the Douglas district, Northern Cape Province (NC) [29°02'60.00"S; 23°45'59.99"E, March 2018, *legit* H. Bezuidenhout 871HB, SANBI Pretoria (PRE) & Kimberley South African National Parks Herbarium (KSAN)], where plants infest maize, cotton, and alfalfa fields (Figure 2A). On those crop fields *A. palmeri* was observed growing together with *A. hybridus* L. long since naturalized in South Africa. On that farm, *A. palmeri* only drew attention because it showed resistance against herbicides that usually provide good control of other amaranths. Recently (2019), *A. palmeri* was observed along a main road running past the farm on which it was first recorded in 2018 to the town of Douglas ( $\pm 7$  km distance), as well as in the Douglas town. Another record originates in the Kruger National Park, Limpopo Province (22°22'26.8"S; 31°12'38.4"E, 12 April 2019,





**Figure 2.** *Amaranthus palmeri*. A: South Africa, Northern Cape Province, Douglas district, cotton field, March 2018, photo by C.F. Reinhardt; B: Botswana, North-West district, Kasane town, March 2020, photo by A. Sukhorukov.

*legit* G. Zambatis & N. Zambatis GZ1563, KNP0011482/1) as occasional herb up to 2 m high in disturbed habitats on the floodplain and riverbank of the Limpopo River. Subsequently, samples collected from a farm near Howick, Kwa-Zulu Natal (KZN) were identified as *A. palmeri* (29°29'19.0"S; 30°10'52.0"E, March 2020, *legit* S. Friis, PRE).

In addition to herbarium vouchers, various living amaranths were collected from the locations in NC and KZN and additional verification was done by sequencing the ITS region using the method developed by Murphy and Tranel (2018). ITS sequencing identified *A. palmeri* in NC (NCBI: MT811920) and KZN (MT811923), *A. hybridus* in NC (MT811921) and KZN (MT811924), and *A. standleyanus* (MT811922) from the NC region. Subsequent sequence characterization of herbicide resistance genes has shown the two *A. palmeri* populations to be distinct from each other.

Another record for *Amaranthus palmeri* in southern Africa is from Botswana (Figure 1) where it was found in enormous quantity along the roads in Kasane town, North-West district of Botswana (17°47'31.96"S; 25°10'49.93"E, March 2020, *legit* A. Sukhorukov 482, G, K, MW, W). Both male and female individuals were present in flowering and fruiting [female individuals] stages (Figure 2B). A survey of the local people showed that the plant has been present in the town for at least 10 years. *A. palmeri* has invaded ruderal places distributed across an area of 10 km<sup>2</sup> westwards from the border with Zimbabwe. Along the main road in Kasane, the number of plants ranges from 40–50 per 100 m. Further observations are needed to determine whether *A. palmeri* is an invasive plant, since this status requires

that the offspring are found “at considerable distances from the parent plants” (Pyšek et al. 2004). The species was not observed in the investigated areas in Zimbabwe and Zambia, including the road Kasane – Victoria Falls.

### *Habitat preferences*

*Amaranthus palmeri* is a plant adapted to both ruderal and segetal communities. It was not found in the natural plant communities but in a degraded area on the floodplain of the Limpopo River. It is plausible that the KNP population originated from propagules carried downstream in the Limpopo River from farmland adjacent to this river. Our observations are in agreement with other data showing the same habitat preferences (Zohary 1966; Smith et al. 2000). In Kasane town (Botswana), most individuals were highly branched and remarkably tall compared with other weedy plants, reaching 250 cm in height. The co-occurring species found there were typical alien weeds (e.g., *Amaranthus spinosus* L., *Boerhavia repens* L., *B. diffusa* L., *Tridax procumbens* L.). On the Douglas farm (South Africa), in cotton crop that reaches height of  $\pm 100$  cm, *A. palmeri* grows to about as tall as the crop, and in this setting the weed tends to be multi-stemmed or bushy. In alfalfa crop on the same farm, *A. palmeri* is regularly cut together with the crop during harvest, but the weed survives and produces fruits and seeds below the cut line, or by growing horizontally under the alfalfa canopy. In contrast, on said farm in the taller growing maize crop, *A. palmeri* tends to grow as tall and even taller than the crop, i.e. attains 200 cm height and taller. When growing in maize crop this weed tends to have a single stem. The aforementioned growth characteristics of *A. palmeri* are in agreement with the data from USA and Argentina (e.g., Berger et al. 2016, with references therein).

## **Discussion**

### *Possible impact of Amaranthus palmeri*

Like many other amaranths that are prolific seed producers (Weaver 1984; Mohler and Callaway 1995), a single individual of *Amaranthus palmeri* is able to produce up to 600,000 seeds in favorable conditions (Keeley et al. 1987). The seeds are both non-dormant and dormant and can successfully germinate after 3 years (Korres et al. 2018). This indicates that *A. palmeri* establishes a long lasting seed bank. Rapid growth and high density of *A. palmeri* induce competition for water, nutrients, and light and reduce the harvesting efficiency of agricultural crops (Smith et al. 2000, with references therein). The spread can be further accelerated if the plant is cultivated for its edible leaves (Shurtleff and Aoyagi 2010; Grubben and Denton 2004) and may have negative impacts on both ruderal and segetal plant communities.

The effective methods of population control and local eradication of *Amaranthus palmeri* have not yet been established in the new territory, but control programs that rely heavily on various herbicides are utilized in the USA, and herbicide type vary from crop to crop (Lindsay et al. 2017). *Amaranthus palmeri* is well adapted to withstand certain common active principles (a.p.) through the rapid evolution of resistance to at least eight very important a.p. sites of action (Ward et al. 2013; Berger et al. 2016, with references therein). However, we believe that selective mechanical removal of younger plants may be effective, because these are much taller than other similar looking amaranths. Unfortunately, selective mechanical removal will require weeding by hand which is time-consuming and relatively expensive. Mechanical control by means of soil tillage operations can be considered either as substitute or supplement for chemical control. Failure to control this weed effectively and timely in crops will likely result in catastrophic crop yield losses.

Based on the presence and persistence of *A. palmeri* populations for long periods, its invasive status in southern Africa can be considered as “naturalized”. However, this status may be changed to “invasive” if further observations will allow to conclude and quantify its negative impacts and its rate of spread and consequent distribution in adjacent regions.

#### *Notes on distribution of some alien Amaranthus species in Africa*

The taxonomic composition of *Amaranthus* in the temperate and tropical regions is different. For instance, *A. retroflexus* L., widely distributed in the temperate regions of Eurasia, has not been found in the lowlands of African tropics. In addition, many records of the “tropical” amaranths in temperate European Russia are clearly erroneous (Sukhorukov and Kushunina 2012). According to the observations of the first author, the most common naturalized amaranths in Africa are the American *A. spinosus* L. and, especially, *A. viridis* L. Other frequently encountered species are *A. blitum* L. and *A. hybridus* L. [sub *A. patulus* Bertol. in the treatment of Amaranthaceae by Baker and Clarke (1909)]. *Amaranthus blitum* prefers arable lands, while *A. hybridus* is a typical ruderal species which can also grow on volcanic soil at elevation up to 1500 m a.s.l. (caldera of Fogo volcano, Fogo Island, Cape Verde; legit A. Sukhorukov, G, MW).

*Amaranthus dubius* Mart. was not listed in the earlier African floras (e.g., Baker and Clarke 1909; Hutchinson and Dalziel 1927; Maire 1962), but is reported from different African countries in many recent floristic accounts (e.g., Germishuizen and Meyer 2003; Grubben and Denton 2004; Mapaura and Timberlake 2004; Arechavaleta et al. 2005; Phiri 2005). Despite the fact that *A. dubius* is now widely cultivated in many tropical countries in Africa as a pot herb along with other species of the same genus (Grubben and Denton 2004), its intentional introduction in Africa is



doubtful. Surprisingly, this species is considered to be a “protected weed” (Grubben and Denton 2004), although it does not need any protection, and cultivation as a food crop clearly increases its spread. The current scattered distribution of *A. dubius* in Africa shown in APD (2019) is clearly underestimated. In fact, *A. dubius* is frequently found in waste and grassy places in different African countries (Sukhorukov, field observations and revision of the herbarium material at BR and K). In the countries with arid tropical climate, it thrives in any irrigated areas and sewage (e.g., on Sal Island, Cape Verde), and is abundant in disturbed habitats in the inlands of Santo Antão Island (*pers. obs.* of AS in 2018). We consider *A. dubius* a naturalized species according to Pyšek et al. (2004) in the tropics with humid climate.

Another South American amaranth, *Amaranthus standleyanus* Parodi, is mostly known from the arid subtropical parts of Africa (Germishuizen and Meyer 2003; Klaassen and Kwembeya 2013; APD 2019, with references herein) as well as from Macaronesia (Cape Verde) with a pending invasive status (Arechavaleta et al. 2005; Sukhorukov et al. 2018b). It is still rare in Namibia and Cape Verde, and in these countries *A. standleyanus* can be considered an alien introduced species in human-made habitats.

### Acknowledgements

We cordially thank the anonymous reviewers and Associate Editor Giuseppe Brundu for the valuable comments to the first draft of the paper. CFR thanks herbarium staff (Pieter Winter & Erich van Wyk) of South African National Biodiversity Institute for the confirmation of the identification of *Amaranthus palmeri* in South Africa. APS thanks Alison Heath and Roger Heath (Kew Botanical Gardens) for discussing of the record of *A. palmeri* in Botswana.

### Funding declaration

The study design of APS was supported by the scientific programme AAAA-A16-116021660045-2 of the Department of Higher Plants (Moscow State University) and the Tomsk State University competitiveness improvement programme. CFR and BJV are project leaders in the South African Herbicide Resistance Initiative (SAHRI) based at the University of Pretoria – this project is managed and funded by Enterprises University of Pretoria (Pty) Ltd.

### References

- APD (2019) African Plant Database, version 3.4.0. <http://www.ville-ge.ch/musinfo/bd/cjb/africa/> (accessed 07 June 2020)
- Arechavaleta M, Zurita MN, Marrero MC, Martín JL (2005) Lista preliminar de especies silvestres de Cabo Verde (hongos, plantas y animales terrestres). Litografía A. Romero, Santa Cruz de Tenerife, 155 pp
- Baker JG, Clarke CB (1909) Amaranthaceae. In: Thiselton-Dyer WT (ed), Flora of Tropical Africa, Volume 6(1). Reeve & Co, London, pp 14–75
- Barnard SA, van Heerden JM, Gerber HS (1992) Evaluation of shrub species for sheep grazing in the Strandveld of the Cape west coast of South Africa. *African Journal of Range and Forage Science* 9(3): 111–113, <https://doi.org/10.1080/02566702.1992.9648309>
- Berger S, Madeira PT, Ferrell J, Gettys L, Morichetti S, Cantero JJ, Nunes C (2016) Palmer amaranth (*Amaranthus palmeri*) identification and documentation of ALS-resistance in Argentina. *Weed Science* 64: 312–320, <https://doi.org/10.1614/WS-D-15-00125.1>
- Boulos L (1995) Flora of Egypt. Checklist. Al Hadara Publishing, Cairo, 287 pp
- Brockington SF, Alexandre R, Ramdial J, Moore MJ, Crawley S, Dhingra A, Hilu K, Soltis DE, Soltis PS (2009) Phylogeny of the Caryophyllales sensu lato: revisiting hypotheses on pollination biology and perianth differentiation in the core Caryophyllales. *International Journal of Plant Sciences* 170: 627–643, <https://doi.org/10.1086/597785>

- Costea M, Weaver S, Tardif FJ (2004) The biology of Canadian weeds. 130. *Amaranthus retroflexus* L., *A. powellii* S.Watson and *A. hybridus* L. (update). *Canadian Journal of Plant Sciences* 84: 631–668, <https://doi.org/10.4141/P02-183>
- di Castri F, Hansen AJ, Debussche M (2012) Biological invasions in Europe and the Mediterranean basin. Kluwer Academic Publishers, Dordrecht-Boston-London, 463 pp
- EPPO (2020) European and Mediterranean Plant Protection Organization. <https://gd.eppo.int/taxon/AMAPA> (accessed 07 June 2020)
- Esser K, Kadereit JW, Lüttge U, Runge M (2013) Progress in Botany: Genetics, physiology, systematics, Ecology. Springer Science & Business Media, Berlin-Heidelberg, 374 pp
- Forman J (2003) The introduction of American plant species into Europe: Issues and consequences. In: Child LE, Bröck JH, Brundu G, Prach K, Pyšek P, Wade PM, Williamson M (eds), Plant invasions. Ecological threats and management solutions. Backhuys Publ., Leiden, The Netherlands, pp 17–39
- Germishuizen G, Meyer NL (2003) Plants of southern Africa: an annotated checklist. Strelitzia 14. Pretoria, National Botanical Institute, 1232 pp
- Grubben GJH, Denton OA (2004) Vegetables [Plant resources of Africa, 2]. PROTA, Wageningen, 668 pp
- Hernández-Ledesma P, Berendsohn WG, Borsch T, von Mering S, Akhiani H, Arias S, Castañeda-Noa I, Eggli U, Eriksson R, Flores-Olvera H, Fuentes-Bazán S, Kadereit G, Klak C, Korotkova N, Nyffeler R, Ocampo G, Ochoterena H, Oxelman B, Rabeler RK, Sanchez A, Schlumberger BO, Uotila P (2015) A taxonomic backbone for the global synthesis of species diversity in the angiosperm order Caryophyllales. *Willdenowia* 39: 281–383, <https://doi.org/10.3372/wi.45.45301>
- Heshmati GA, Squires VR (2013) Combating desertification in Asia, Africa and the Middle East: Proven practices. Springer Science & Business Media, Dordrecht, 476 pp, <https://doi.org/10.1007/978-94-007-6652-5>
- Hutchinson JL, Dalziel J (1927) Flora of West Tropical Africa, Volume 1. Crown Agents, London, 960 pp
- Iamonico D (2015) Taxonomic revision of the genus *Amaranthus* (Amaranthaceae) in Italy. *Phytotaxa* 199: 1–84, <https://doi.org/10.11646/phytotaxa.199.1.1>
- Iamonico D, El Mokni R (2017) *Amaranthus palmeri*, a second record for Africa and notes on *A. sonoriensis* nom. nov. *Bothalia* 47: a2100, <https://doi.org/10.4102/abc.v47i1.2100>
- Keeley PE, Carter CH, Thullen RJ (1987) Influence of planting date on growth of Palmer amaranth (*Amaranthus palmeri*). *Weed Science* 35: 199–204, <https://doi.org/10.1017/S0043174500079054>
- Klaassen E, Kwembeya E (2013) A checklist of Namibian indigenous and naturalised plants. National Botanical Research Institute, Windhoek, 597 pp
- Korres NE, Norsworthy JK, Young BG, Reynolds DB, Johnson WG, Conley SP, Smeda RJ, Mueller TC, Spaunhorst DJ, Gage KL, Loux M, Kruger GR, Bagavathiannan MV (2018) Seedbank persistence of Palmer Amaranth (*Amaranthus palmeri*) and Waterhemp (*Amaranthus tuberculatus*) across diverse geographical regions in the United States. *Weed Science* 66: 446–456, <https://doi.org/10.1017/wsc.2018.27>
- Lindsay K, Popp M, Norsworthy J, Bagavathiannan M, Powles S, Lacoste M (2017) Decision support for long-term Palmer Amaranth (*Amaranthus palmeri*) control. *Weed Technology* 31: 915–927, <https://doi.org/10.1017/wet.2017.69>
- Maire R (1962) Flore de l’Afrique du Nord, Volume 8. Lechevalier, Paris, 303 pp
- Mapaura A, Timberlake J (2004) A checklist of Zimbabwean vascular plants [Southern African Botanical Diversity Network Report No. 33]. SABONET, Pretoria & Harare, 148 pp
- Mohler CL, Callaway BM (1995) Effects of tillage and mulch on weed seed production and seed banks in sweet corn. *Journal of Applied Ecology* 32: 627–639, <https://doi.org/10.2307/2404658>
- Murphy BP, Tranel PJ (2018) Identification and validation of *Amaranthus* species-specific SNPs within the ITS region: applications in quantitative species identification. *Crop Science* 58: 304–311, <https://doi.org/10.2135/cropsci2017.06.0359>
- Novoa A, Le Roux JJ, Robertson MP, Wilson JR, Richardson DM (2014) Introduced and invasive cactus species: a global review. *AoB Plants* 7: plu078, <https://doi.org/10.1093/aobpla/plu078>
- Phiri PSM (2005) A checklist of Zambian vascular plants [Southern African Botanical Diversity Network Report No. 32]. SABONET, Pretoria, 169 pp
- Pyšek P, Richardson DM, Rejmánek M, Webster GL, Williamson M, Kirscher J (2004) Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon* 53: 131–143, <https://doi.org/10.2307/4135498>
- Sauer JD (1955) Revision of the dioecious amaranths. *Madroño* 13(1): 5–46
- Shackleton RT, Witt ABR, Piroris FM, van Wilgen BW (2017) Distribution and socio-ecological impacts of the invasive alien cactus *Opuntia stricta* in eastern Africa. *Biological Invasions* 19: 2427–2441, <https://doi.org/10.1007/s10530-017-1453-x>



- Shurtleff W, Aoyagi A (2010) History of soybeans and soyfoods in Southeast Asia (13<sup>th</sup> century to 2010): Extensive annotated bibliography and sourcebook. Soyinfo Center, Lafayette, 1031 pp
- Smith DT, Baker RV, Steele GR (2000) Palmer amaranth (*Amaranthus palmeri*) impacts on yield, harvesting, and ginning in dryland cotton (*Gossypium hirsutum*). *Weed Technology* 14: 122–126, [https://doi.org/10.1614/0890-037X\(2000\)014\[0122:PAAPIO\]2.0.CO;2](https://doi.org/10.1614/0890-037X(2000)014[0122:PAAPIO]2.0.CO;2)
- Sukhorukov AP (2009) Additions to the flora of Kazakhstan Republic. *Bulletin of Moscow Society of Naturalists (MOIP), sect. biol.* 114(3): 72 [in Russian]
- Sukhorukov AP (2011) New invasive alien plant species in the forest-steppe and northern steppe subzones of European Russia: secondary range patterns, ecology and causes of fragmentary distribution. *Feddes Repertorium* 122: 287–304, <https://doi.org/10.1002/fedr.201100004>
- Sukhorukov AP (2014) The carpology of Chenopodiaceae with reference to the phylogeny, systematics and diagnostics of its representatives. Grif & K, Tula, 400 pp [In Russian with English summary]
- Sukhorukov AP, Kushunina MA (2012) Some new information about the flora of western districts of Saratov Region. *Bulletin of the Botanical Garden of Saratov University* 10: 21–28 [in Russian]
- Sukhorukov AP, Verloove F, Alonso MA, Belyaeva IV, Chapano C, Crespo MB, Aouni M, El Mokni R, Maroyi A, Shekede MD, Vicente A, Dreyer A, Kushunina M (2017) Chorological and taxonomic notes on African plants, 2. *Botany Letters* 164: 135–153, <https://doi.org/10.1080/23818107.2017.1311281>
- Sukhorukov AP, Nilova MV, Erst AS, Kushunina M, Baider C, Verloove F, Salas-Pascual M, Belyaeva IV, Krinitsina AA, Bruyns PV, Klak C (2018a) Diagnostics, taxonomy, nomenclature and distribution of perennial Sesuvium (Aizoaceae) in Africa. *PhytoKeys* 92: 45–88, <https://doi.org/10.3897/phytokeys.92.22205>
- Sukhorukov AP, Kushunina M, El Mokni R, Goñalons LS, El Aouni MH, Daniel TF (2018b) Chorological and taxonomic notes on African plants, 3. *Botany Letters* 165: 228–240, <https://doi.org/10.1080/23818107.2018.1465467>
- Sukhorukov AP, Kushunina M, El Mokni R, Ardenghi NMG, Verloove F, Uotila P, Baider C, Bruyns PV, Klak C (2019) Chorological and taxonomic notes on African plants, 4: Caryophyllales. *Botany Letters* 166: 401–416, <https://doi.org/10.1080/23818107.2019.1652848>
- van Wilgen BW, Measey J, Richardson DM, Wilson JR, Zengeya TA (2020) Biological invasions in South Africa. Springer, Cham, 975 pp, <https://doi.org/10.1007/978-3-030-32394-3>
- Ward SM, Webster TM, Steckel LE (2013) Palmer amaranth (*Amaranthus palmeri*): a review. *Weed Technology* 27: 12–27, <https://doi.org/10.1614/WT-D-12-00113.1>
- Weaver SE (1984) Differential growth and competitive ability of *Amaranthus retroflexus*, *A. powellii* and *A. hybridus*. *Canadian Journal of Plant Sciences* 64: 715–724, <https://doi.org/10.4141/cjps84-098>
- Wetzel DK, Horak MJ, Skinner DZ, Kulakow PA (1999) Transfer of herbicide resistance traits from *Amaranthus palmeri* to *Amaranthus rudis*. *Weed Science* 47: 538–543, <https://doi.org/10.1017/S0043174500092237>
- Zohary M (1966) Flora Palaestina, Volume. 1 (text). The Israel Academy of Sciences and Humanities, Jerusalem, 364 pp