

Human Eltonian niche

A FRAMEWORK FOR THE ELTONIAN NICHE OF HUMANS

Remington J. Moll^{1*}, remington.moll@unh.edu

Alexander K. Killion², killion@umich.edu

Matt W. Hayward^{3,4}, matthew.hayward@newcastle.edu.au

Robert A. Montgomery⁵, robert.montgomery@zoo.ox.ac.uk

¹Department of Natural Resources and the Environment, University of New Hampshire, 56 College Road, Durham, NH 03824 USA

²School for Environment & Sustainability, University of Michigan, 440 Church St, Ann Arbor, MI 48109 USA

³Conservation Biology Research Group, School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, Australia

⁴Mammal Research Centre, University of Pretoria, Tshwane, South Africa, X001; and Centre for African Conservation Ecology, Nelson Mandela University, Port Elizabeth, South Africa 6213

⁵Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Tubney House, Abingdon Road, Tubney, Oxon OX13 5QL, United Kingdom

*Corresponding Author

ABSTRACT

Recent research has highlighted several influential roles that humans play in ecosystems, including that of a super predator, hyperkeystone species, and niche constructor. This work has begun to describe the Eltonian niche of humans, which encompasses humanity's cumulative ecological and evolutionary roles in trophic systems. Yet, we lack a unifying framework that brings together these strands of research, links them to eco-evolutionary and social-cultural theory, and identifies current research needs. Here, we present such a framework in hope of facilitating a more holistic approach to operationalizing human roles in trophic systems across an increasingly anthropogenic biosphere. The framework underscores how humans play numerous nuanced roles in trophic systems, from top-down to bottom-up, that entail not only pernicious effects but also benefits for many nonhuman species. Such a nuanced view of the Eltonian niche of humans is important for understanding complex social-ecological system functioning and enacting effective policies and conservation measures.

KEYWORDS

Anthropocene; anthrome; community ecology; *Homo sapiens*; social-ecological systems

INTRODUCTION

Ostensibly, the study of ecology, which encompasses the relationships between organisms and their environments, should include humans within its purview, since *Homo sapiens* is a species that interacts with both biotic and abiotic components of ecosystems. Yet historically ecology has maintained a disciplinary tradition that has tended to place humans “outside” of ecosystems, or even of nature more broadly (Tansley 1935, Alberti et al. 2003). This dualistic view dominated the formative early years of ecology in Western societies (Tansley 1935, McDonnell and Pickett 1993). In the modern era, however, humanity's local, regional, and global effects on

ecosystems have begun to render such human-dominated versus “natural” dichotomies untenable (Alberti et al. 2003). Although humans have strongly influenced ecosystems for millennia (Jones 1969, Dunne et al. 2016, Trant et al. 2016, Fisher et al. 2019), the scale and magnitude of anthropogenic effects have soared in recent decades (Ellis 2011). As it has become increasingly clear that humans act as a dominant species across Earth’s ecosystems (Vitousek et al. 1997), ecologists have begun incorporating humans as a functional species into ecological thought and theory (McDonnell and Pickett 1993, Alberti et al. 2003, Dorresteijn et al. 2015, Dunne et al. 2016).

Only recently, however, have ecologists explored how humans act as an intrinsic, functional species within trophic systems. Hereafter, we use the term *trophic systems* in a general sense to refer to ecological communities consisting of more than one trophic level. Trophic systems are therefore inclusive of the classical ecological notion of *food webs* (cf. Polis and Winemiller 1995), but extend to include relationships beyond consumer-resource dynamics. Contemporary research has now documented how humans influence trophic systems in a variety of ways. For example, humans moderate the effects from apex predators by lethally exploiting them, modifying their behavior, and altering landscapes, and such effects can have cascading influence over broader trophic system dynamics in terrestrial and aquatic contexts alike (Ferretti et al. 2010, Estes et al. 2011, Ordiz et al. 2013, Oriol-Cotterill et al. 2015). Accordingly, humans have often been envisioned as affecting trophic systems from top-down positions (i.e., as an apex predator; Roopnarine 2014, Bugir et al. 2021). This top-down perspective has been reinforced by recent work classifying humans as *super predators* (i.e., a species that exploits another species at high rates or targets adult individuals with certain "trophy" characteristics; Darimont et al. 2015)

Human Eltonian niche

and a *hyperkeystone species* (i.e., a high-order species that regulates other keystone species; Worm and Paine 2016).

However, humans should not only be envisioned to affect trophic systems through top-down pathways but also via those that are bottom-up. For instance, humans fix more atmospheric nitrogen than all other nonhuman sources combined (Vitousek et al. 1997), have caused the atmospheric carbon dioxide concentration to rise to a level that would be sustained for a millennium even if all human-related emissions immediately ceased (Solomon et al. 2009), and have transformed vast landscapes across the globe via agricultural practices and urbanization (Grimm et al. 2008, Pickett et al. 2011). In these and other ways, humans shape trophic systems by acting as a *niche constructor* that exerts intense selective pressure on nonhuman species via bottom-up pathways (Palumbi 2001, Smith 2007, Albuquerque et al. 2017). Moreover, humans not only shape trophic systems but also respond to them. Given that anthropogenic effects on such systems can feedback to affect human behavior, attitudes, and decision-making, humans must be viewed as an interactive agent that both affects and is affected by trophic system dynamics (Ostrom 2007; Box 1).

What is presently lacking is a unifying framework that brings together these strands of research and links them to eco-evolutionary theory and the social-cultural dimensions of human action and behavior. Here, we present such a framework by drawing upon the classical ecological concept of the *Eltonian niche*, which encompasses a species' functional roles within a given system (Sales et al. 2021, Elton 1927, Mittelbach 2012). Importantly, the Eltonian niche of modern humans is unlike that of nonhuman species in that it is more complex, multi-faceted, and system-dependent. As outlined above, humans should be expected to exert positive or negative direct influence on many – and perhaps all – species in a trophic system along multiple

Human Eltonian niche

pathways, with complex indirect modifications to a variety of interactions between organisms. These human-triggered and multi-faceted effects can in turn alter system-level processes and feedback to influence human decision-making and behavior. Further, the intensities of such effects and their feedbacks vary considerably from system to system. For instance, North American and European hunters remove prey at rates roughly ten times that of their African counterparts, thereby acting as very different kinds of super predators, each with system-specific effects (Darimont et al. 2015). Given the potential for this complexity, ecologists must strive to take a multi-faceted approach to define the ecological roles of humans in a given system, operationalize them, analyze them systematically to determine their relative importance, and link the most salient anthropogenic effects stemming from these roles with policy and conservation efforts (Montgomery et al. 2018, 2020).

Below, we first describe a general Eltonian framework aimed at facilitating this goal and then we illustrate how it clarifies several prominent *Eltonian roles* humans play within trophic systems. To borrow classical terms from Hutchinson's definition of the niche (Hutchinson 1957), this effort can be viewed as first defining the fundamental Eltonian niche of humans writ large as all of the actions, roles, and trophic effects of humanity as a whole, and then highlighting several aspects of humans' realized Eltonian niche on a case- and system-specific basis (cf. Sales et al. 2021). We conclude by discussing the implications of our analysis for future research, conservation, and policy. We acknowledge that the concept of the niche has evolved over time and presently many different, occasionally conflicting, viewpoints are held (Sales et al. 2021). Here, we focus on the Eltonian concept, which emphasizes a species' role or "what it is doing" in a trophic system (Elton 1927). This focus is roughly analogous to the ecosystem ecology view as described by Chase and Liebold (2003) which emphasizes how species' functional traits impact

systems rather than how environmental factors govern species' distributions (cf. Sales et al. 2021). Our focus also reflects the classical Hutchinsonian view that the niche is a property of a species rather than the environment (Hutchinson 1957). This species-centric aspect of the niche is especially pertinent to humans, as the functional ecological roles of humans have changed – and will continue to change – over time.

FRAMEWORK DESCRIPTION

At the highest and most general level, we structure the Eltonian framework of humans around five central, interconnected components (Fig. 1): *human decision-making* drives *human actions* that result in *ecological and evolutionary effects*. Human actions and their effects are characterized by several key *qualities*. Finally, the combination of human actions, ecological and evolutionary effects, and their qualities can lead to humans playing one or more of five unique *Eltonian roles* in a given system. Below we describe each of these components in turn.

At a fundamental level, *human decision-making* drives human actions and is responsive to reciprocal feedbacks emerging from the ecological and evolutionary effects of those actions (Fig. 2). Importantly, the cognitive capabilities and socio-cultural nature of humans are unlike any other species on Earth and translate to trophic effects that are not merely ecological in nature (Manfredo 2008, Ellis 2015). We would be remiss not to mention that human actions are driven by normative attributes (e.g., values, attitudes, beliefs, interests, etc.) and social phenomena and processes (e.g., markets, governance, culture, etc.; Fig. 1), which are ultimately responsive to both ecological and non-ecological changes in social-ecological systems (Fulton et al. 1996, Manfredo 2008, Manfredo et al. 2014, Bennett et al. 2017). These drivers of human action are the subject of vast amounts of literature and research across disciplines beyond ecology, such as economics and philosophy, and thus fall outside the scope of this article. Nonetheless, drawing

Human Eltonian niche

connections between ecological phenomena and the diverse socio-cultural drivers of human decision-making is crucial, and has given rise to interdisciplinary efforts such as the social-ecological systems framework (Ostrom 2007; Box 1). Although included in our framework, these drivers are not strictly a part of the Eltonian niche of humans, which is ecological in nature. Rather, they are the socio-cultural constructs and processes that give rise to particular realizations of the Eltonian niche of humans in a given system and location (Ellis 2015).

Thus, our framework focuses on the ecologically-relevant *human actions* that result from individual and collective human decision-making and their effects on trophic systems. We have organized these actions into four categories, including lethal exploitation (hunting, poaching, or culling), nonlethal activities (movement, travel, or recreation), landscape modification (infrastructure development, agricultural production, or activities associated with livestock), and translocation of nonhuman species (Fig. 2). This is because each of these actions have particular *ecological and evolutionary effects*. Ecological effects include direct effects on individual animals (e.g., via altered traits such as behavior) and populations (e.g., reductions, establishment, or persistence; Fig. 2). These direct effects can lead to indirect effects that can propagate through a trophic system vertically or horizontally, and can take any form of indirect effects described in classical ecological theory (Strauss 1991, Wootton 1994; Fig. 2), namely, exploitative competition (e.g., humans competing with marine predators for fish; Fig. 3a; Hansson et al. 2018), trophic cascades (e.g., humans lethally exploiting carnivores, thus benefiting their prey; Fig. 3b; Henke and Bryant 1999), apparent competition (e.g., human hunters targeting one species that leads to exploitation of a sympatric species; Fig. 3c; Clayton et al. 1997), and indirect commensalisms and mutualisms (e.g., the "human shield" effect whereby humans benefit prey by repelling their predators; Fig. 3d; Hebblewhite et al. 2005, Berger 2007).

The aggregate consequences of direct and indirect effects are ecosystem-scale outcomes involving species' persistence, community dynamics, and ecosystem function and services (Fig. 2), which can result in singular, pulsed, or long-lasting changes to trophic systems (cf. Williams et al. 2020). For example, local, regulated hunting of wildlife species can result in pulsed compensatory mortality across seasons while altered soil chemistry due to agricultural practices can trigger rapid changes in microbial community structure that persist for many decades (Skalski et al. 2005, Fichtner et al. 2014). The intensity and temporal nature largely determine the evolutionary effects of these human-induced ecological dynamics. Finally, ecological outcomes related to both direct effects and their systems-level consequences feedback into human decision-making (Fig. 2; Box 1). For example, the ecosystem-level effects of apex carnivores include the regulation of nutrient cycles, hydrological dynamics, and community stability (Estes et al. 2011, Marshall et al. 2013, Ripple et al. 2014). These processes subsequently influence ecosystem services such as carbon storage (Schmitz et al. 2018). Thus, ecologically-focused human actions (e.g., reintroductions of these carnivores into systems where they have been extirpated) can have positive, trophic-mediated feedback effects on society, which in turn can influence human decision-making regarding land management, animal conservation, and carbon storage (Nelson et al. 2008, Schmitz et al. 2018). In such feedbacks, the lines between the ecology and sociology of humans become interactive. Much recent work in social-ecological systems has focused on understanding such connections, and the framework presented here serves to operationalize human-related roles in trophic systems from an ecological perspective (Box 1). Similarly, human actions can trigger direct and indirect ecological effects that feedback upon themselves (human-mediated ecological feedbacks; Fig. 2). For instance, in many areas across North America, white-tailed deer (*Odocoileus virginianus*) have become

Human Eltonian niche

overabundant due to a lack of natural predators (DeYoung 2011). This species is a keystone herbivore that strongly influences vegetation structure and function through foraging pressure, and its overabundance can reduce plant diversity (Waller and Alverson 1997, Greenwald et al. 2008). Controlling white-tailed deer populations through lethal exploitation has been shown to not only effectively reduce their abundance (a direct effect), but also promote native plant biodiversity (an indirect effect) that in turn supports native bird communities (a human-mediated ecological feedback; McShea and Rappole 2000, Côté et al. 2004).

Human actions and their associated effects can be further characterized by three central *qualities*, namely their spatial scale (ranging from local to global), temporal nature (i.e., singular, occasional, pulsed, or continuous), and intensity (Fig. 2; see examples in Fig. 4). Often the qualities of human actions and their ecological effects will be tightly coupled. For example, more intense lethal exploitation of a given species is likely to have more intense ecological effects in the form of population decline (Packer et al. 2009, Fuller 2014, Darimont et al. 2015). Similarly, as human actions occur over larger spatial scales and become more continuous, their resultant ecological effects are likely to share these qualities. For instance, the burning of fossil fuels occurs near-continuously on a worldwide spatial scale, resulting in global ecological changes that will likely persist for millennia (Vitousek et al. 1997, Solomon et al. 2009). The intensity and scales of these ecological effects in turn drive evolutionary responses (Johnson and Munshi-South 2017, Des Roches et al. 2021). There are, of course, exceptions to these patterns. Low intensity, small scale, and singular human actions could have strong, large scale, and long-lasting ecological consequences, such as a single campfire igniting a large swath of prairie or forest (Balch et al. 2017). Likewise, high intensity human actions do not always translate to uniformly intense ecological or evolutionary effects. For example, the effects of herbicide applications can

Human Eltonian niche

range from relatively innocuous to highly pernicious depending on the toxicity and environmental persistence of the chemical employed (Pratt et al. 1997, Newman 2014). Adding to these complexities are the non-linear changes that frequently occur in both human actions and their effects over time. For example, Native Americans in North America hunted the passenger pigeon (*Ectopistes migratorius*) at sustainable intensities for centuries, but European colonization and the hunting technologies that accompanied it non-linearly intensified and expanded hunting pressure to the point where the once-hyperabundant species was exterminated over several decades (Fuller 2014, Murray et al. 2017). Moreover, given the globalized nature of human economic activity and travel, local effects can be driven by human actions originating at vast distance. Such telecoupled relationships are common in social-ecological processes related to agriculture, ecotourism, commercial forestry, and exotic species translocation (Liu et al. 2007, 2013). Cataloguing all combinations of human actions, their effects, and their associated qualities is clearly beyond the scope of this paper (but see examples in Fig. 4). Nonetheless, the Eltonian framework presented here serves to help systematically identify these components so that they can be more effectively operationalized and examined for a given system or ecological issue (cf. Moll et al. 2019).

ELTONIAN ROLES PLAYED BY HUMANS

The Eltonian framework components described above provide context to clarify the nature of several prominent *Eltonian roles* that humans play in trophic systems and how they connect to the broader view of the Eltonian niche of humans. Stated another way, the broad components above define the fundamental Eltonian niche of humans in general terms that capture all the ways humans can function in trophic systems, while what follows focuses on the realized Eltonian niche of humans in particular cases (cf. Hutchinson 1957). Below we highlight three

Human Eltonian niche

Eltonian roles that have been previously identified (super predators, hyperkeystone species, and niche constructors), and two which have been less formally articulated in the literature (pseudomutualists and risk-responders; Fig. 2).

Super predators

Humans can act as *super predators* when exploitative or nonlethal actions induce strong effects on species' populations or traits (Fig. 2). Notably, humans exploit wild animals at rates higher – sometimes much higher – than those documented among nonhuman species. In aquatic systems, for instance, humans exploit fish species at rates 14 times greater than those associated with nonhuman marine predators (Fig. 5a; Darimont et al., 2015). In terrestrial systems, human hunting of certain species can nearly double mortality from nonhuman predators (Darimont et al. 2015). Indeed, there are numerous examples of continent-wide defaunation events that exemplify humans' super predatory role over terrestrial fauna (Schwartz et al. 2003). Humans also function as super predators by removing predators and individuals that face virtually no threat from nonhuman predators. For example, unlike any other species, human hunters often target the healthiest and largest adult individuals or those possessing “trophy” characteristics (e.g., large antlers; Darimont et al. 2015). Note that the above super predatory rates calculated by Darimont et al. (2015) examined only the mortality of adults in prey populations rather than all individuals, further underscoring humans' unique focus on large, reproductively mature individuals. On broader timescales, both humans and related hominin ancestors have uniquely exerted strong top-down control over Earth's megafauna, leading to declines and extinctions of modern and historical species alike (Baker and Clapham 2004, Estes et al. 2011, Smith et al. 2018). Although unconstrained super predation typically leads to overexploitation and deleterious population-level effects (Darimont et al. 2009, 2015), humans' removal of adult individuals can also benefit

Human Eltonian niche

ecosystems by controlling overabundant prey populations where nonhuman apex predators are absent (e.g., regulated hunting of herbivorous ungulates; McShea and Rappole 2000, Côté et al. 2004, Greenwald et al. 2008).

Initially, the term *super predators* focused on humans' lethal capabilities (Darimont et al. 2015). However, humans can also non-lethally alter species' behavior via the pathway of fear. A variety of species, from waterfowl to large ungulates, make behavioral adjustments to avoid encountering humans and vehicles (Frid and Dill 2002, Montgomery et al. 2020). These non-lethal risk effects even influence species situated in apex trophic positions (Clinchy et al. 2016, Smith et al. 2017, Montgomery et al. 2020). For example, many apex carnivores alter their spatio-temporal movement, activity, and foraging behavior to avoid humans, including pumas (*Puma concolor*; Smith et al. 2017; Fig. 5b), African lions (*Panthera leo*; Loveridge et al., 2017), brown bears (*Ursus arctos*; Ordiz et al., 2011), tigers (*Panthera tigris*; Carter et al., 2012) and Eurasian lynx (*Lynx lynx*; Gehr et al., 2017). Thus, humans function as super predators along lethal and nonlethal pathways. This role becomes increasingly prominent as lethal exploitation and nonlethal human activity grows more intense, spatially expansive, and continuous in nature (Darimont et al. 2015, Montgomery et al. 2020).

Hyperkeystone species

Humans can also play the role of a high-order *hyperkeystone species* when anthropogenic effects influence nonhuman keystone species, that is, those species whose trophic effects are disproportionately large to their abundance (Power et al. 1996, Worm and Paine 2016). Such cascading effects can result in dramatic consequences for ecosystem functioning, with the potential to restructure species assemblages and trophic system dynamics (Gill et al. 2009, Worm and Paine 2016). With respect to the broader Eltonian niche of humans (Fig. 2), this role can

Human Eltonian niche

emerge any time that human actions begins to affect nonhuman keystone species in ways that modify the degree to which such species influence overall trophic dynamics. Examples of humans functioning as a hyperkeystone species include local extirpations of keystone species in both aquatic (e.g., sea otters *Enhydra lutis*; Estes and Palmisano 1974) and terrestrial systems (e.g., large carnivores; Fig. 5c; Darimont et al., 2015; Estes et al., 2011; Ripple et al., 2014). Such extirpations can release herbivore populations from predation pressure, triggering impacts that cascade down to vegetative communities (Estes et al. 2011). The hyperkeystone conception of humans as a functional species in trophic systems provides a bridge between research approaches focused on human behavior and those emphasizing classic food web theory (Worm and Paine 2016).

Niche constructor

A third Eltonian role humans can occupy is that of a *niche constructor* (Albuquerque et al., 2017; Ellis, 2015). Niche construction occurs when a species alters the environment in ways that initiate selective pressure inter- or intra-specifically (Odling-Smee 2003). Within this context, ecosystem engineering (Jones et al. 1994, Smith 2007) falls under the broader umbrella of niche construction (see Odling-Smee et al., 2013). In nonhuman species, classic examples of niche construction include the soil chemistry-altering effects of earthworms (Lee 1985) and “home construction” activities performed by numerous species (e.g., nest-building, burrowing, dam building; Odling-Smee et al., 2013). Humans are prolific niche constructors, with urbanization and animal domestication representing clear cases (Jones et al. 1994, Odling-Smee et al. 2013). In many instances, humans have created niches for species that in turn benefit society, a dynamic that has resulted in the coevolution of humans and agricultural crops and livestock (Ellis 2015).

Humans' role as a niche constructor often results when landscape modification is intense and alters ecosystems through bottom-up pathways (Fig. 2; Box 2). For example, urban development replaces native habitat with artificial structures and landscaped vegetation, leading to a variety of effects on the species inhabiting a given system. In contexts with low basal resource levels (e.g., those in desert biomes), such urban development can translate to increased plant productivity and biodiversity due to water provisioning and landscaping (Faeth et al. 2005, Grimm et al. 2008, McKinney 2008), especially in wealthy areas of cities (i.e., a "luxury" effect; Hope et al. 2003). These changes can intensify bottom-up competition for resources by urban specialist species while reducing top-down pressures through the elimination of fragmentation-sensitive predators (Faeth et al. 2005). Urban areas in particular and landscape development more generally also elevate environmental noise and light to levels unprecedented in evolutionary time (Swaddle et al. 2015, Shannon et al. 2016). This anthropogenic sensory pollution alters selective pressures on a variety of ecological traits and processes, including physiology, mate selection, communication, and predator-prey interactions (Swaddle et al. 2015, Roca et al. 2016, Shannon et al. 2016; Fig. 5d). These pressures can drive rapid phenotypic changes over contemporary timescales (i.e., years to decades) and might result in eco-evolutionary feedbacks that impact ecosystem services and human health, although the potential for such dynamics are understudied (Francis et al. 2009, Alberti 2015, Swaddle et al. 2015, Alberti et al. 2017). On the whole, the acceleration of urbanization and other forms of landscape modification is resulting in a proliferation of emergent niches within evolutionarily novel "anthromes" across the planet (Ellis 2011). The spatial scales, temporal dynamics, and ecological characteristics of such anthromes vary considerably, but all are generally characterized by novel

Human Eltonian niche

selective pressures driven by human actions that result in strong changes in eco-evolutionary dynamics and feedbacks (Ellis 2011, Alberti 2015, Miles et al. 2019).

Pseudo-mutualists

The roles above have been formalized in the literature and have tended to focus on the negative effects of humans on trophic systems. Such effects are certainly deserving of research attention. However, humans can also positively influence animal populations, and such positive influences can have reciprocal benefits for humans. Biocontrol efforts exemplify this phenomenon, whereby humans intentionally augment one species to control another that is considered undesirable from an anthropogenic perspective. For instance, human promotion of pest-controlling vertebrates (e.g., raptors) via nest box provision can benefit agricultural production by reducing seed and fruit predation (Lindell et al. 2018; Fig. 5e). Ecotourism is another example of how humans intentionally benefit species that are actively managed (e.g., via space and resource provision for species within a reserve) while receiving benefits in the form of pleasure derived from wildlife viewing and interaction (Skibins et al. 2012; Box 2). Consumptive human activities, such as hunting and fishing, are also often mutualistic when financial resources and enjoyment from recreation feedback to support the sustainable management of animal populations (Organ and Fritzell 2000, Granek et al. 2008). Finally, recent research has highlighted how human habitation, particularly by indigenous groups, over long periods (e.g., centuries to millennia) can enhance rather than degrade forest productivity and maintain a diverse mosaic of vegetation, often through the judicious application of fire (DeGraaf and Yamasaki 2001, Trant et al. 2016, Hoffman et al. 2017).

In the above and other instances, humans act as *pseudo-mutualists* that benefit other species while furthering human well-being. Notably, these beneficial human actions are often

Human Eltonian niche

intentional and direct rather than mere incidental, auxiliary outcomes of human decision-making. In describing this role, the prefix *pseudo-* indicates that the benefits accrued to humans do not necessarily reflect the positive effects traditionally implied by mutualisms, such as enhanced fitness (Bronstein 2009). This, however, does not imply that such mutualism categorically fails to affect human fitness in the classic sense. For example, benefits accrued from wildlife in the form of psychological well-being improve human health (Miller 2005, Shanahan et al. 2016) and ecosystem services provided by numerous species (e.g., organic waste removal by scavengers; Whelan et al. 2008) likely have indirect effects on human life expectancy and population dynamics. However, studies on these human demographic responses are rare (Butler et al. 2005) and represent an intriguing line of future work.

Risk-responders

Humans are not only initiators of effects on trophic systems, but also recipients. For instance, humans regularly make behavioral decisions to avoid risks associated with perceived danger from non-human animals (Slovic and Peters 2006). Humans' evolutionary ancestors (i.e., species in the Primate order) exhibited a range of antipredator behaviors, including alarm calls, mobbing, and spatial avoidance (Gursky and Nekaris 2007). Similarly, humans' closest ancestors, Plio-Pleistocene hominins (6.0-1.8 Ma), mitigated predation risk from predators via vigilance, flight, and counterattack (Treves and Palmqvist 2007, Hart and Sussman 2009). These behaviors played a crucial role in hominid daily life in as recently as one million years ago (Hart and Sussman 2009).

Of course, most modern humans engage in lifestyles that are very different from those of our hominid ancestors. Nonetheless, evolutionarily-embedded antipredator behaviors linger in human consciousness, influencing individual and collective behavior (Hart and Sussman 2009),

Human Eltonian niche

meaning humans can play the Eltonian role of *risk-responders*. For example, even though risk of injury from large carnivores is at a historic low, most people today still strongly express fear and intensify vigilance when presented with scenarios involving large carnivores (Jacobs et al. 2014). Humans also become more protective of household members (e.g., keeping children or pets indoors) when living in areas where carnivores are regularly encountered (Crooks and Soulé 1999, Collard 2012). Beyond such anti-predator responses, humans adjust behavior to avoid risk of discomfort or disease from many invertebrates and microbes (Curtis 2014). For example, Lyme disease, the most common vector-borne zoonotic in North America, is transmitted to humans via the bite of *Ixodes* ticks that carry an infectious bacterium (Nelson et al. 2015, Schwartz et al. 2017). Humans act as risk-responders and reduce exposure to ticks by changing their behavior, such as avoiding brushy areas that are favorable for ticks and reducing off-trail hiking during outdoor recreation (Fig. 5f; Center for Disease Control and Prevention 2020). Such altered behavior can reduce humans' ecological impact in forests and grasslands by spatially constraining human activity that would otherwise disturb animals and damage vegetation (Stankowich 2008, Barros and Pickering 2017).

Despite these behavioral responses to risk, the human role of a risk-responder has rarely been integrated into studies of human-trophic system interactions (Prokop and Fančovičová 2010). Doing so is crucial because the risk of negative interactions with nonhuman species not only influences human behavior (Killion et al. 2020), but also plays a role in shaping policy, conservation, and management by influencing human attitudes and values. For example, fear-inducing species, such as sharks or snakes, receive lower amounts of conservation support compared to neutral or aesthetically-pleasing species (Knight 2008, Liordos et al. 2017). Therefore, as in pseudo-mutualism, risk-responses entail feedbacks between humans and

nonhuman species (Fig. 2). These feedback pathways can quickly become complex as the number of species related to a particular risk increase. For instance, although Lyme disease is transmitted to humans by ticks, as noted above, its overall prevalence is related to numerous abiotic and biotic ecological factors. One of these factors is the presence and abundance of certain mesocarnivores (e.g., foxes and mustelids), which are hypothesized to reduce Lyme disease via predation on the rodents that serve as a natural reservoir for the infectious bacteria (Levi et al. 2012, Hofmeester et al. 2017, Moll et al. 2020). Wildlife authorities typically allow for regulated harvest of such mesocarnivores. Recently, a citizen group petitioned for protection of these mesocarnivores with the aim of an indirect reduction of rodent populations and Lyme disease risk (Royar 2018). In this instance, one group's risk-response to ticks took the form of protecting other species in the trophic system, which created tension with a second group's value for harvesting those species. This example highlights how ecological and social values combine with complex trophic processes to create situations where humans play multiple Eltonian roles that can be in tension with one another, underscoring the need to take a social-ecological approach to finding resolutions for complex issues (Box 1).

IMPLICATIONS FOR FUTURE RESEARCH, CONSERVATION, AND POLICY

Ecologists are currently situated in a unique and somewhat perplexing position whereby virtually all systems under current study are strongly shaped by a species that has often been omitted from classical ecological theory and inquiry: humans (Tansley 1935, McDonnell and Pickett 1993). Today, this historical tendency is no longer defensible, and ecologists have increasingly sought to understand the myriad of ways that humans shape trophic systems (Alberti et al. 2003, Worm and Paine 2016). At the same time, social scientists studying human and natural systems have taken a broader approach to understand linked social and environmental outcomes (Ostrom 2007,

Human Eltonian niche

Binder et al. 2013). The framework that we have developed here (Fig. 1, Fig. 2) provides a connection between these approaches unified under the classical ecological concept of the Eltonian niche.

Perhaps more than anything else, our analysis emphasizes how humans play numerous interconnected roles in trophic systems. As the above examples have shown, many ecologically- and evolutionarily-relevant human actions occur simultaneously, meaning that humans can play multiple influential Eltonian roles within a given system. An important first step to examine such complexity is to operationalize human actions by identifying their categorical classifications (i.e., lethal exploitation, landscape modification, nonlethal activity, and translocation) and characteristic qualities (i.e., their spatial scales, temporal nature, and intensity) within a given system (Box 2; Fig. 4). Such operationalization helps crystallize what is meant by the often-invoked but typically vague concept of “anthropogenic effects” (Moll et al. 2019, Montgomery et al. 2020). Humans’ Eltonian roles vary between and within systems, and are propelled by diverse social and cultural forces that underlie human value systems and governance (Ellis 2015). We therefore discourage monolithic conceptualizations of modern humans as being a particular type of species, or as playing singular ecological roles. Casting the human species as predominately playing any one of the five roles discussed above – or any other singular ecological role – oversimplifies the manifold and context-specific ways in which humans affect trophic systems (Montgomery et al. 2020). Such oversimplifications undermine efforts to develop a clear understanding of the complexity of human-natural systems, which in turn hinders the development of effective policy and management for a given ecosystem (Liu et al. 2007, Montgomery et al. 2018).

Rather, we recommend that ecologists endeavor to capture the complexity of human effects in trophic systems by operationalizing multiple human-related variables in models and analyzing them systematically to determine which anthropogenic roles matter most in a given system (Dorresteijn et al. 2015, Dunne et al. 2016). Given that much research on human effects on trophic systems has focused on direct effects, there is a particularly pressing need to examine the extent to which such direct actions lead to indirect and system-level consequences (cf. Worm and Paine 2016) and resultant socio-eco-evolutionary dynamics (Des Roches et al. 2021). Importantly, multiple human-related variables should be integrated into broader interdisciplinary efforts that analyze social-ecological models by connecting humans' realized Eltonian niche to corresponding social drivers (Box 1). Certainly, efforts are currently being pursued towards these ends (Carter and Linnell 2016, Montgomery et al. 2018). One aim of our analysis is to make such endeavors more effective by organizing the manifold effects of humans on trophic systems in a coherent, ecologically-based framework.

Lastly, it is crucial to note that conservation policy and management will fail if they do not comprehensively account for the multifarious impacts humans have on trophic systems. For example, harvest management that only considers lethal effects of hunters while omitting their nonlethal effects on species' behavior will likely underestimate the overall impact of harvest regulations (Cromsigt et al. 2013). In contrast, policy that neglects the positive roles humans play in trophic systems will overestimate potential harm due to anthropogenic activity. For instance, policy aimed at bird conservation (e.g., the United States' Migratory Bird Treaty Act) must consider not only pernicious anthropogenic effects such as land conversion, but also human actions such as feeder provision that result in enormous benefits for many species (Davies et al. 2009). Therefore, the formation of effective policy and management requires taking a holistic

view of the Eltonian niche of humans, cataloguing the roles played by humans in a given system, and identifying which of those are most prominent. Indeed, conservation policy has evolved over the last several decades from taking a species- and problem-specific approach to more holistic efforts focused on ecosystems (Nesshöver et al. 2017). We hope the framework presented here will advance these efforts by providing ecologists a common structure to account for human roles in trophic systems and a common language regarding these roles that can be shared by researchers, managers, policymakers, and conservation organizations.

ACKNOWLEDGEMENTS

We thank the handling editor, C. Darimont, and two anonymous reviewers for their constructive feedback.

REFERENCES

- Adams CE. 2016. *Urban Wildlife Management*. 3rd ed. CRC Press.
- Alberti M. 2015. Eco-evolutionary dynamics in an urbanizing planet. *Trends in Ecology and Evolution* 30: 114–126.
- Alberti M, Correa C, Marzluff JM, Hendry AP, Palkovacs EP, Gotanda KM, Hunt VM, Apgar TM, Zhou Y. 2017. Global urban signatures of phenotypic change in animal and plant populations. *Proceedings of the National Academy of Sciences of the United States of America* 114: 8951–8956.
- Alberti M, Marzluff JM, Shulenberger E, Bradley G, Ryan C, Zumbrunnen C. 2003. Integrating human into ecology: Opportunities and challenges for studying urban ecosystems. *BioScience* 53: 1169–1179.
- Albuquerque UP, Gonçalves PHS, Ferreira Júnior WS, Chaves LS, Oliveira RC da S, Silva TLL da, Santos GC dos, Araújo E de L. 2017. Humans as niche constructors: Revisiting the concept of chronic anthropogenic disturbances in ecology. *Perspectives in Ecology and Conservation* 16: 1–11.
- Baker CS, Clapham PJ. 2004. Modelling the past and future of whales and whaling. *Trends in Ecology and Evolution* 19: 365–371.
- Balch JK, Bradley BA, Abatzoglou JT, Chelsea Nagy R, Fusco EJ, Mahood AL. 2017. Human-started wildfires expand the fire niche across the United States. *Proceedings of the National Academy of Sciences of the United States of America* 114: 2946–2951.
- Barros A, Pickering CM. 2017. How networks of informal trails cause landscape level damage to

- vegetation. *Environmental Management* 60: 57–68.
- Bateman PW, Fleming PA. 2012. Big city life: carnivores in urban environments. *Journal of Zoology* 287: 1–23.
- Bennett NJ, Roth R, Klain SC, Chan K, Christie P, Clark DA, Cullman G, Curran D, Durbin TJ, Epstein G, Greenberg A, Nelson MP, Sandlos J, Stedman R, Teel TL, Thomas R, Verissimo D, Wyborn C. 2017. Conservation social science: Understanding and integrating human dimensions to improve conservation. *Biological Conservation* 205: 93–108.
- Berger J. 2007. Fear, human shields and the redistribution of prey and predators in protected areas. *Biology Letters* 3: 620–623.
- Beschta RL, Ripple WJ. 2016. Riparian vegetation recovery in Yellowstone: The first two decades after wolf reintroduction. *Biological Conservation* 198: 93–103.
- Binder CR, Hinkel J, Bots PWG, Pahl-Wostl C. 2013. Comparison of frameworks for analyzing social-ecological systems. *Ecology and Society* 18: 26.
- Bojarska K, Kwiatkowska M, Skórka P, Gula R, Theuerkauf J, Okarma H. 2017. Anthropogenic environmental traps: Where do wolves kill their prey in a commercial forest? *Forest Ecology and Management* 397: 117–125.
- Bronstein JL. 2009. The evolution of facilitation and mutualism. *Journal of Ecology* 97: 1160–1170.
- Bugir CK, Peres CA, White KS, Montgomery RA, Griffin AS, Rippon P, Clulow J, Hayward MW. 2021. Prey preferences of modern human hunter-gatherers. *Food Webs* 26.
- Bull JW, Ejrnæs R, Macdonald DW, Svenning J, Sandom CJ. 2018. Fences can support restoration in human-dominated ecosystems when rewilding with large predators. *Restoration Ecology* 1–12.
- Bullock JM, Bonte D, Pufal G, da Silva Carvalho C, Chapman DS, García C, García D, Matthysen E, Delgado MM. 2018. Human-mediated dispersal and the rewiring of spatial networks. *Trends in Ecology and Evolution* 33: 958–970.
- Butler CD, Corvalan CF, Koren HS. 2005. Human health, well-being, and global ecological scenarios. *Ecosystems* 8: 153–162.
- Carter NH, Linnell JDC. 2016. Co-adaptation is key to coexisting with large carnivores. *Trends in Ecology and Evolution* 31: 575–578.
- Carter NH, Shrestha BK, Karki JB, Man N, Pradhan B, Liu J. 2012. Coexistence between wildlife and humans at fine spatial scales. *Proceeding of the Royal Society London Biological Sciences* 109: 15360–15365.
- Center for Disease Control and Prevention. 2020. Preventing tick bites. Lyme disease. (24 December 2020; <https://www.cdc.gov/lyme/prev/index.html>).
- Chase JM, Leibold MA. 2003. *Ecological niches: Linking classical and contemporary approaches*. University of Chicago Press.

- Clayton L, Keeling M, Milner-Gulland EJ. 1997. Bringing home the bacon: A spatial model of wild pig hunting in Sulawesi, Indonesia. *Ecological Applications* 7: 642–652.
- Clinchy M, Zanette LY, Roberts D, Justin P, Newman C, Macdonald DW. 2016. Fear of the human “super predator” far exceeds the fear of large carnivores in a model mesocarnivore. *Behavioral Ecology* 27: 1826–1832.
- Collard RC. 2012. Cougar-human entanglements and the biopolitical un/making of safe space. *Environment and Planning D: Society and Space* 30: 23–42.
- Cote M, Nightingale AJ. 2012. Resilience thinking meets social theory: Situating social change in socio-ecological systems (SES) research. *Progress in Human Geography* 36: 475–489.
- Côté SD, Rooney TP, Tremblay J-P, Dussault C, Waller DM. 2004. Ecological impacts of deer overabundance. *Annual Review of Ecology, Evolution, and Systematics* 35: 113–147.
- Cromsigt JPGM, Kuijper DPJ, Adam M, Beschta RL, Churski M, Eycott A, Kerley GIH, Mysterud A, Schmidt K, West K. 2013. Hunting for fear: Innovating management of human-wildlife conflicts. *Journal of Applied Ecology* 50: 544–549.
- Crooks KR, Soulé ME. 1999. Mesopredator release and avifaunal extinctions in a fragmented system. *Nature* 400: 563–566.
- Curtis VA. 2014. Infection-avoidance behaviour in humans and other animals. *Trends in Immunology* 35: 457–464.
- Darimont CT, Carlson SM, Kinnison MT, Paquet PC, Reimchen TE, Wilmers CC. 2009. Human predators outpace other agents of trait change in the wild. *Proceedings of the National Academy of Sciences* 106: 8–10.
- Darimont CT, Fox CH, Bryan HM, Reimchen TE. 2015. The unique ecology of human predators. *Science* 349: 858–861.
- Davies ZG, Fuller RA, Loram A, Irvine KN, Sims V, Gaston KJ. 2009. A national scale inventory of resource provision for biodiversity within domestic gardens. *Biological Conservation* 142: 761–771.
- DeGraaf RM, Yamasaki M. 2001. *New England wildlife*. University Press of New England.
- Delcourt PA. 2004. *Prehistoric Native Americans and ecological change: Human ecosystems in eastern North America since the Pleistocene*. 1st ed. Cambridge University Press.
- DeYoung C. 2011. Population dynamics. Pages 147–180 in Hewitt DG, ed. *Biology and management of white-tailed deer*. Taylor & Francis.
- Dorresteijn I, Schultner J, Nimmo DG, Fischer J, Hanspach J, Kuemmerle T, Kehoe L, Ritchie EG. 2015. Incorporating anthropogenic effects into trophic ecology: predator–prey interactions in a human-dominated landscape. *Proceedings of the Royal Society B: Biological Sciences* 282: 20151602.
- Dunne JA, Maschner H, Betts MW, Huntly N, Russell R, Williams RJ, Wood SA. 2016. The roles and impacts of human hunter-gatherers in North Pacific marine food webs. *Scientific Reports* 6: 1–9.

- Ellis EC. 2011. Anthropogenic transformation of the terrestrial biosphere. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 369: 1010–1035.
- Ellis EC. 2015. Ecology in an anthropogenic biosphere. *Ecological Monographs* 85: 287–331.
- Elton CS. 1927. *Animal Ecology*. University of Chicago Press.
- Estes JA, Palmisano JF. 1974. Sea otters: Their role in structuring nearshore communities. *Science* 185: 1058–1060.
- Estes JA, Terborgh J, Brashares JS, Power ME, Berger J, Bond WJ, Carpenter SR, Essington TE, Holt RD, Jackson JBC, Marquis RJ, Oksanen L, Oksanen T, Paine RT, Pikitch EK, Ripple WJ, Sandin S a, Scheffer M, Schoener TW, Shurin JB, Sinclair ARE, Soulé ME, Virtanen R, Wardle D a. 2011. Trophic downgrading of planet Earth. *Science* 333: 301–306.
- Faeth SH, Marussich WA, Shochat E, Warren PS. 2005. Trophic dynamics in urban communities. *BioScience* 55: 399.
- Ferretti F, Worm B, Britten GL, Heithaus MR, Lotze HK. 2010. Patterns and ecosystem consequences of shark declines in the ocean. *Ecology Letters* 13: 1055–1071.
- Fichtner A, von Oheimb G, Härdtle W, Wilken C, Gutknecht JLM. 2014. Effects of anthropogenic disturbances on soil microbial communities in oak forests persist for more than 100 years. *Soil Biology and Biochemistry* 70: 79–87.
- Fisher JA, Shackelford N, Hocking MD, Trant AJ, Starzomski BM. 2019. Indigenous peoples' habitation history drives present-day forest biodiversity in British Columbia's coastal temperate rainforest. *People and Nature* 1: 103–114.
- Francis CD, Ortega CP, Cruz A. 2009. Noise pollution changes avian communities and species interactions. *Current Biology* 19: 1415–1419.
- Frid A, Dill L. 2002. Human-caused disturbance stimuli as a form of predation risk. *Ecology and Society* 6: 11.
- Fuller E. 2014. *The passenger pigeon*. Princeton University Press.
- Fulton DC, Manfredo MJ, Lipscomb J. 1996. Wildlife value orientations: A conceptual and measurement approach. *Human Dimensions of Wildlife* 1: 24–47.
- Gaynor KM, Hojnowski CE, Carter NH, Brashares JS. 2018. The influence of human disturbance on wildlife nocturnality. *Science* 360: 1232–1235.
- Gehr B, Hofer EJ, Muff S, Ryser A, Vimercati E, Vogt K, Keller LF. 2017. A landscape of coexistence for a large predator in a human dominated landscape. *Oikos* 1–11.
- Gill JL, Williams JW, Jackson ST, Lininger KB, Robison GS. 2009. Pleistocene megafaunal collapse, novel plant communities, and enhanced fire regimes in North America. *Science* 326: 1100–1103.
- Granek EF, Madin EMP, Brown MA, Figueira W, Cameron DS, Hogan Z, Kristianson G, de Villiers P, Williams JE, Post J, Zahn S, Arlinghaus R. 2008. Engaging recreational fishers

- in management and conservation: Global case studies. *Conservation Biology* 22: 1125–1134.
- Greenwald KR, Petit LJ, Waite TA. 2008. Indirect effects of a keystone herbivore elevate local animal diversity. *Journal of Wildlife Management* 72: 1318–1321.
- Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM. 2008. Global change and the ecology of cities. *Science* 319: 756–760.
- Gursky, S. L., and K. A. I. Nekaris, eds. 2007. *Primate anti-predator strategies*. Springer.
- Hansson S, Bergström U, Bonsdorff E, Härkönen T, Jepsen N, Kautsky L, Lundström K, Lunneryd SG, Ovegård M, Salmi J, Sendek D, Vetemaa M. 2018. Competition for the fish - Fish extraction from the Baltic Sea by humans, aquatic mammals, and birds. *ICES Journal of Marine Science* 75: 999–1008.
- Hart D, Sussman RW. 2009. *Man the hunted*. Expanded Ed. Westview Press.
- Hayward MW, Kerley GIH, Adendorff J, Moolman LC, O'Brien J, Sholto-Douglas A, Bissett C, Bean P, Fogarty A, Howarth D, Slater R. 2007. The reintroduction of large carnivores to the Eastern Cape, South Africa: an assessment. *Oryx* 41: 205.
- Hebblewhite M, White CA, Nietvelt CG, McKenzie JA, Hurd TE, Fryxell JM, Bayley SE, Paquet PC. 2005. Human activity mediates a trophic cascade caused by wolves. *Ecology* 86: 2135–2144.
- Henke SE, Bryant FC. 1999. Effects of coyote removal on the faunal community in western Texas. *Journal of Wildlife Management* 63: 1066–1081.
- Hoffman KM, Lertzman KP, Starzomski BM. 2017. Ecological legacies of anthropogenic burning in a British Columbia coastal temperate rain forest. *Journal of Biogeography* 44: 2903–2915.
- Hofmeester TR, Jansen PA, Wijnen HJ, Coipan EC, Fonville M, Prins HHT, Sprong H, van Wieren SE. 2017. Cascading effects of predator activity on tick-borne disease risk. *Proceedings of the Royal Society B: Biological Sciences* 284: 20170453.
- Hope D, Gries C, Zhu W, Fagan WF, Redman CL, Grimm NB, Nelson AL, Martin C, Kinzig A. 2003. Socioeconomics drive urban plant diversity. *Proceedings of the National Academy of Sciences* 100: 8788–8792.
- Hutchinson GE. 1957. Concluding remarks. *Cold Spring Harbor Symposia on Quantitative Biology* 22: 415–427.
- Jacobs MH, Vaske JJ, Dubois S, Fehres P. 2014. More than fear: Role of emotions in acceptability of lethal control of wolves. *European Journal of Wildlife Research* 60: 589–598.
- Johnson MTJ, Munshi-South J. 2017. Evolution of life in urban environments. *Science* 358: eaam8327.
- Jones CG, Lawton JH, Shachak M. 1994. Organisms as ecosystem engineers. *Oikos* 69: 373–386.

- Jones R. 1969. Fire-stick farming. *Australian Natural History* 16: 224–228.
- Killion AK, Ramirez JM, Carter NH. 2020. Human adaptation strategies are key to cobenefits in human–wildlife systems. *Conservation Letters* e12769.
- Knight AJ. 2008. ‘Bats, snakes and spiders, Oh my!’ How aesthetic and negativistic attitudes, and other concepts predict support for species protection. *Journal of Environmental Psychology* 28: 94–103.
- Lee KE. 1985. *Earthworms: their ecology and relation with soil and land use*. Academic Press.
- Levi T, Kilpatrick AM, Mangel M, Wilmers CC. 2012. Deer, predators, and the emergence of Lyme disease. *Proceedings of the National Academy of Sciences* 109: 10942–10947.
- Lindell C, Eaton RA, Howard PH, Roels SM, Shave ME. 2018. Enhancing agricultural landscapes to increase crop pest reduction by vertebrates. *Agriculture, Ecosystems and Environment* 257: 1–11.
- Liordos V, Kotsiotis VJ, Anastasiadou M, Karavasiyas E. 2017. Effects of attitudes and demography on public support for endangered species conservation. *Science of the Total Environment* 595: 25–34.
- Liu J, Dietz T, Carpenter SR, Alberti M, Folke C, Moran E, Pell AN, Deadman P, Kratz T, Lubchenco J, Ostrom E, Ouyang Z, Provencher W, Redman CL, Schneider SH, Taylor WW. 2007. Complexity of coupled human and natural systems. *Science* 317: 1513–1516.
- Liu J, Hull V, Batistella M, DeFries R, Dietz T, Fu F, Hertel TW, Izaurrealde RC, Lambin EF, Li S, Marinelli LA, McConnell WJ, Moran EF, Naylor R, Ouyang Z, Polenske KR, Reenberg A, de Miranda Rocha G, Simmons CS, Verburg PH, Vitousek PM, Zhu C. 2013. Framing sustainability in a telecoupled world. *Ecology and Society* 18: 26.
- Loveridge AJ, Valeix M, Elliot NB, Macdonald DW. 2017. The landscape of anthropogenic mortality: how African lions respond to spatial variation in risk. *Journal of Applied Ecology* 54: 815–825.
- Manfredo MJ. 2008. *Who cares about wildlife?* Springer.
- Manfredo MJ, Teel TL, Gavin MC, Fulton D. 2014. Consideration in representing human individuals in social-ecological models. Pages 137–158 in Manfredo MJ, Vaske JJ, Rechkemmer A, and Duke EA, eds. *Understanding society and natural resources: Forging new strands of integration across the social sciences*. Springer.
- Marshall KN, Hobbs NT, Cooper DJ. 2013. Stream hydrology limits recovery of riparian ecosystems after wolf reintroduction. *Proceedings. Biological sciences / The Royal Society* 280: 20122977.
- Marzluff JM, Ewing K. 2001. Restoration of fragmented landscapes for the conservation of birds: A general framework and specific recommendations for urbanizing landscapes. *Restoration Ecology* 9: 280–292.
- McDonnell, M. J., and S. T. A. Pickett, eds. 1993. *Humans as components of ecosystems*. Springer-Verlag.

- McKinney ML. 2002. Urbanization, biodiversity, and conservation. *BioScience* 52: 883–890.
- McKinney ML. 2006. Urbanization as a major cause of biotic homogenization. *Biological Conservation* 127: 247–260.
- McKinney ML. 2008. Effects of urbanization on species richness: A review of plants and animals. *Urban Ecosystems* 11: 161–176.
- McShea WJ, Rappole JH. 2000. Managing the abundance and diversity of breeding bird populations through manipulation of deer populations. *Conservation Biology* 14: 1161–1170.
- Miles LS, Breitbart ST, Wagner HH, Johnson MTJ. 2019. Urbanization shapes the ecology and evolution of plant-arthropod herbivore interactions. *Frontiers in Ecology and Evolution* 7: 1–14.
- Miles LS, Johnson JC, Dyer RJ, Verrelli BC. 2018. Urbanization as a facilitator of gene flow in a human health pest. *Molecular Ecology* 27: 3219–3230.
- Miller JR. 2005. Biodiversity conservation and the extinction of experience. *Trends in Ecology and Evolution* 20: 430–434.
- Mittelbach GG. 2012. *Community ecology*. Sinauer Associates, Inc.
- Moll RJ, Cepek JD, Lorch PD, Dennis PM, Robison T, Millspaugh JJ, Montgomery RA. 2018. Humans and urban development mediate the sympatry of competing carnivores. *Urban Ecosystems* 21: 765–778.
- Moll RJ, Cepek JD, Lorch PD, Dennis PM, Tans E, Robison T, Millspaugh JJ, Montgomery RA. 2019. What does urbanization actually mean? A framework for urban metrics in wildlife research. *Journal of Applied Ecology* 56: 1289–1300.
- Moll RJ, Eaton JT, Cepek JD, Lorch PD, Dennis PM, Robinson T, Tsao J, Montgomery RA. 2020. Dynamic rodent behavioral response to predation risk: Implications for disease ecology. *Oecologia* 19: 67–78.
- Montgomery RA, Elliott KC, Hayward MW, Gray SM, Millspaugh JJ, Riley SJ, Kissui BM, Kramer DB, Moll RJ, Mudumba T, Tans ED, Muneza AB, Abade L, Beck JM, Hoffmann CF, Booher CR, Macdonald DW. 2018. Examining evident interdisciplinarity among prides of lion researchers. *Frontiers in Ecology and Evolution* 6: 1–13.
- Montgomery RA, Macdonald DW, Hayward MW. 2020. The inducible defences of large mammals to human lethality. *Functional Ecology* 1–16.
- Murray GGR, Soares AER, Novak BJ, Schaefer NK, Cahill JA, Baker AJ, Demboski JR, Doll A, da Fonseca RR, Fulton TL, Gilbert MTP, Heintzman PD, Letts B, McIntosh G, O’Connell BL, Peck M, Pipes ML, Rice ES, Santos KM, Sohrweide AG, Vohr SH, Corbett-Detig RB, Green RE, Shapiro B. 2017. Natural selection shaped the rise and fall of passenger pigeon genomic diversity. *Science* 954: 951–954.
- Nelson CA, Saha S, Kugeler KJ, Delorey MJ, Shankar MB, Hinckley AF, Mead PS. 2015. Incidence of clinician-diagnosed lyme disease, United States, 2005–2010. *Emerging*

Infectious Diseases 21: 1625–1631.

- Nelson E, Polasky S, Lewis DJ, Plantinga AJ, Lonsdorf E, White D, Bael D, Lawler JJ. 2008. Efficiency of incentives to jointly increase carbon sequestration and species conservation on a landscape. *Proceedings of the National Academy of Sciences of the United States of America* 105: 9471–9476.
- Nesshöver C, Assmuth T, Irvine KN, Rusch GM, Waylen KA, Delbaere B, Haase D, Jones-Walters L, Keune H, Kovacs E, Krauze K, Külvik M, Rey F, van Dijk J, Vistad OI, Wilkinson ME, Wittmer H. 2017. The science, policy and practice of nature-based solutions: An interdisciplinary perspective. *Science of the Total Environment* 579: 1215–1227.
- Newman MC. 2014. *Fundamentals of ecotoxicology: The science of pollution*. CRC Press.
- Newsome D, Moore SA, Dowling RK. 2012. *Natural area tourism: Ecology, impacts and management*. 2nd ed. Channel View Publications.
- Odling-Smee FJ. 2003. *Niche construction: The neglected process in evolution*. Princeton University Press.
- Odling-Smee J, Erwin DH, Palkovacs EP, Feldman MW, Laland KN. 2013. Niche construction theory: A practical guide for ecologists. *The Quarterly Review of Biology* 88: 3–28.
- Ordiz A, Bischof R, Swenson JE. 2013. Saving large carnivores, but losing the apex predator? *Biological Conservation* 168: 128–133.
- Ordiz A, Støen OG, Delibes M, Swenson JE. 2011. Predators or prey? Spatio-temporal discrimination of human-derived risk by brown bears. *Oecologia* 166: 59–67.
- Organ JF, Fritzell EK. 2000. Trends in consumptive recreation and the wildlife profession. *Wildlife Society Bulletin* 28: 780–787.
- Oriol-Cotterill A, Valeix M, Frank LG, Riginos C, Macdonald DW. 2015. Landscapes of Coexistence for terrestrial carnivores: The ecological consequences of being downgraded from ultimate to penultimate predator by humans. *Oikos* 124: 1263–1273.
- Ostrom E. 2007. A diagnostic approach for going beyond panaceas. *Proceedings of the National Academy of Sciences* 104: 15181–15187.
- Packer C, Kosmala M, Cooley HS, Brink H, Pintea L, Garshelis D, Purchase G, Strauss M, Swanson A, Balme G, Hunter L, Nowell K. 2009. Sport hunting, predator control and conservation of large carnivores. *PLoS ONE* 4: e5941.
- Palumbi SR. 2001. Human as the world's greatest evolutionary force. *Science* 293: 1786–1790.
- Parsons AW, Forrester T, Baker-Whetton MC, McShea WJ, Rota CT, Schuttler SG, Millspaugh JJ, Kays R. 2018. Mammal communities are larger and more diverse in moderately developed areas. *eLife* 7: 1–13.
- Pickett STA, Cadenasso ML, Grove JM. 2005. Biocomplexity in coupled natural-human systems: A multidimensional framework. *Ecosystems* 8: 225–232.

- Pickett STA, Cadenasso ML, Grove JM, Boone CG, Groffman PM, Irwin E, Kaushal SS, Marshall V, McGrath BP, Nilon CH, Pouyat R V., Szlavecz K, Troy A, Warren P. 2011. Urban ecological systems: Scientific foundations and a decade of progress. *Journal of Environmental Management* 92: 331–362.
- Polis GA, Winemiller KO. 1995. *Food webs: Integration of patterns and dynamics*. Springer.
- Power ME, Tilman D, Estes JA, Menge BA, Bond WJ, Mills LS, Daily G, Castilla JC, Lubchenco J, Paine RT. 1996. Challenges in the quest for keystones. *BioScience* 46: 609–620.
- Pratt JR, Melendez AE, Barreiro R, Bowers NJ. 1997. Predicting the ecological effects of herbicides. *Ecological Applications* 7: 1117–1124.
- Prokop P, Fančovičová J. 2010. Perceived body condition is associated with fear of a large carnivore predator in humans. *Annales Zoologici Fennici* 2450: 417–425.
- Pulver S, Ulibarri N, Sobocinski KL, Alexander SM, Johnson ML, Mccord PF. 2018. Frontiers in socio-environmental research: components, connections, scale, and context. *Ecology and Society* 23: 23.
- Ripple WJ, Estes JA, Beschta RL, Wilmers CC, Ritchie EG, Hebblewhite M, Berger J, Elmhagen B, Letnic M, Nelson MP, Schmitz OJ, Smith DW, Wallach AD, Wirsing AJ. 2014. Status and ecological effects of the world’s largest carnivores. *Science* 343: 1241484.
- Rissman AR, Gillon S. 2017. Where are ecology and biodiversity in social–ecological systems research? A review of research methods and applied recommendations. *Conservation Letters* 10: 86–93.
- Roca IT, Desrochers L, Giacomazzo M, Bertolo A, Bolduc P, Deschesnes R, Martin CA, Rainville V, Rheault G, Proulx R. 2016. Shifting song frequencies in response to anthropogenic noise: A meta-analysis on birds and anurans. *Behavioral Ecology* 27: 1269–1274.
- Des Roches S, Brans KI, Lambert MR, Rivkin LR, Savage AM, Schell CJ, Correa C, De Meester L, Diamond SE, Grimm NB, Harris NC, Govaert L, Hendry AP, Johnson MTJ, Munshi-South J, Palkovacs EP, Szulkin M, Urban MC, Verrelli BC, Alberti M. 2021. Socio-eco-evolutionary dynamics in cities. *Evolutionary Applications* 14: 248–267.
- Roopnarine PD. 2014. Humans are apex predators. *Proceedings of the National Academy of Sciences* 111: E796–E796.
- Royar K. 2018. Red Fox and Lyme Disease – Is there a connection in Vermont ?
- Sales LP, Hayward MW, Loyola R. 2021. What do you mean by “niche”? Modern ecological theories are not coherent on rhetoric about the niche concept. *Acta Oecologica* 110: 103701.
- Schmitz OJ, Wilmers CC, Leroux SJ, Doughty CE, Atwood TB, Galetti M, Davies AB, Goetz SJ. 2018. Animals and the zoogeochemistry of the carbon cycle. *Science* 362: eaar3213.
- Schwartz AM, Hinckley AF, Mead PS, Hook SA, Kugeler KJ. 2017. Surveillance for Lyme Disease — United States, 2008–2015. *MMWR. Surveillance Summaries* 66: 1–12.

- Schwartz CC, Swenson JE, Miller SD. 2003. Large carnivores, moose, and humans: a changing paradigm of predator management in the 21st century. *Alces* 39: 41–63.
- Shanahan DF, Bush R, Gaston KJ, Lin BB, Dean J, Barber E, Fuller RA. 2016. Health benefits from nature experiences depend on dose. *Scientific Reports* 6: 1–10.
- Shannon G, Mckenna MF, Angeloni LM, Crooks KR, Fristrup KM, Brown E, Warner KA, Nelson MD, White C, Briggs J, Mcfarland S, Wittemyer G. 2016. A synthesis of two decades of research documenting the effects of noise on wildlife. *Biological Reviews* 91: 985–1005.
- Skalski JR, Ryding KE, Millspaugh JJ. 2005. *Wildlife demography: Analysis of sex, age, and count data*. Elsevier Academic Press.
- Skibins JC, Hallo JC, Sharp JL, Manning RE. 2012. Quantifying the role of viewing the Denali ‘Big 5’ in visitor satisfaction and awareness: Conservation implications for flagship recognition and resource management. *Human Dimensions of Wildlife* 17: 112–128.
- Slovic P, Peters E. 2006. Risk perception and affect. *Current Directions in Psychological Science* 15: 322–325.
- Smith BD. 2007. The ultimate ecosystem engineers. *Science* 315: 1797–1798.
- Smith FA, Smith REE, Lyons SK, Payne JL. 2018. Body size downgrading of mammals over the late Quaternary. *Science* 360: 310–313.
- Smith JA, Suraci JP, Clinchy M, Crawford A, Roberts D, Zanette LY, Wilmers CC. 2017. Fear of the human ‘super predator’ reduces feeding time in large carnivores. *Proceedings of the Royal Society B: Biological Sciences* 284: 20170433.
- Solomon S, Plattner G-K, Knutti R, Friedlingstein P. 2009. Irreversible climate change due to carbon dioxide emissions. *Proceedings of the National Academy of Sciences of the United States of America* 106: 1704–9.
- Stankowich T. 2008. Ungulate flight responses to human disturbance: A review and meta-analysis. *Biological Conservation* 141: 2159–2173.
- Strauss SY. 1991. Indirect effects in community ecology: Their definition, study and importance. *Trends in Ecology and Evolution* 6: 206–210.
- Swaddle JP, Francis CD, Barber JR, Cooper CB, Kyba CCM, Dominoni DM, Shannon G, Aschehoug E, Goodwin SE, Kawahara AY, Luther D, Spoelstra K, Voss M, Longcore T. 2015. A framework to assess evolutionary responses to anthropogenic light and sound. *Trends in Ecology and Evolution* 30: 550–560.
- Tansley AG. 1935. The use and abuse of vegetational concepts and terms. *Ecology* 16: 284–307.
- Trant AJ, Nijland W, Hoffman KM, Mathews DL, McLaren D, Nelson TA, Starzomski BM. 2016. Intertidal resource use over millennia enhances forest productivity. *Nature Communications* 7: 1–8.
- Treves A, Palmqvist P. 2007. Reconstructing hominin interactions with mammalian carnivores (6.0–1.8 Ma). Pages 355–381 in Gursky SL and Nekaris KAI, eds. *Primate Anti-Predator*

Strategies. Springer.

- Valeix M, Fritz H, Loveridge AJ, Davidson Z, Hunt JE, Murindagomo F, Macdonald DW. 2009. Does the risk of encountering lions influence African herbivore behaviour at waterholes? *Behavioral Ecology and Sociobiology* 63: 1483–1494.
- Vitousek PM, Mooney H a, Lubchenco J, Melillo JM. 1997. Human domination of earth's ecosystems. *Science* 277: 494–499.
- Waller DM, Alverson WS. 1997. The white-tailed deer: A keystone herbivore. *Wildlife Society Bulletin* 25: 217–226.
- Whelan CJ, Wenny DG, Marquis RJ. 2008. Ecosystem services provided by birds. *Annals of the New York Academy of Sciences* 1134: 25–60.
- Williams JW, Ordonez A, Svenning JC. 2020. A unifying framework for studying and managing climate-driven rates of ecological change. *Nature Ecology and Evolution* 5: 17–26.
- Wootton JT. 1994. The nature and consequences of indirect effects in ecological communities. *Annual Review of Ecology and Systematics* 25: 443–466.
- Worm B, Paine RT. 2016. Humans as a hyperkeystone species. *Trends in Ecology and Evolution* 31: 600–607.
- Wright GJ, Peterson RO, Smith DW, Lemke TO. 2006. Selection of Northern Yellowstone elk by gray wolves and hunters. *Journal of Wildlife Management* 70: 1070–1078.

Box 1 Connections to social-ecological systems science

Humans and the environment are inherently connected. Several frameworks have been developed to address the various interactions, feedbacks, and complexities between these two systems (Pickett et al. 2005, Liu et al. 2007, Binder et al. 2013). Here we use the term social-ecological systems (SES) science to encompass this research. These paradigms position humans and the environment as interfacing parts of a larger, integrated, complex system to measure the effects of many multidirectional relationships. SES research relies on methodologies from diverse disciplines, where various epistemologies are applied to understand particular interactions between humans and the environment (Binder et al. 2013). For example, many SES approaches use social science theories to understand human decision-making processes in response to environmental changes (e.g., climate change adaptation, human-wildlife conflict, environmental hazard preparedness; Cote and Nightingale 2012). In these instances, traditional ecological models are paired with social data to simultaneously assess the effects of feedbacks between social and ecological units across spatial and temporal scales. Despite their promise, such approaches have been hindered by failing to adequately incorporate and operationalize ecological rules and theory (Rissman and Gillon 2017, Pulver et al. 2018)

Our ecologically-based evaluation of humans as a functional trophic species enhances a SES approach by clarifying how to operationalize humans' multifaceted effects on trophic processes as well as the feedbacks that emerge from such effects. Rather than reconstructing social processes as ecological ones, the roles described herein provide researchers with an ecological foundation to account for how humans interact with and impact trophic systems. For example, social surveys and interviews could be used to measure land owner attitudes and beliefs to estimate the probability of a land use decision, and the impact of that decision on local

Human Eltonian niche

ungulate populations could be measured within an ecological framework using established principles (e.g., niche construction theory; Odling-Smee 2003, Smith 2007; Fig. 1). In this manner, the result of human behaviors is converted to ecological terms to better measure ecological effects, which forms a component of a larger SES analysis. Additionally, the framework can help operationalize the drivers of human behaviors, which are both ecological and social in nature. For example, an urban area with a nearby bear population might propose the use of secure garbage receptacles. The local community's acceptance of such a proposal could be analyzed using both ecological (e.g., human fear of physical harm) and socio-economic variables (e.g., a desire to protect property and belongings). In turn, the ecological effects of such a decision on bear behavior on lower trophic levels could be analyzed under the paradigm of anthropogenic niche construction. Thereby, such an analysis would evaluate the full feedback cycle of the ecological and social drivers of human decision-making as well as the trophic processes affected by those decisions.

Box 2 Applying the framework to contrasting systems

Although the framework can be applied to any trophic system, we illustrate its utility here by comparing the Eltonian niche of humans in contrasting systems: 1) that of a populous city with several hundred thousand inhabitants and 2) that of a large (i.e., thousands of km²) nature reserve or national park (hereafter *reserve*). Of course, specific realizations of such systems will vary according to many factors such as regional abiotic conditions and the socio-economic characteristics of states or nations. These differences notwithstanding, comparing humans' Eltonian niche across these prototypical systems helps clarify its context-dependence.

In the city, intense landscape modification will result in a heterogeneous mosaic of land covers ranging from city centers to greenways to less densely populated residential areas. The ecological consequences of these bottom-up trophic changes are manifold: connectivity and gene flow declines for most species but can increase for a select few well-adapted to fragmentation (Marzluff and Ewing 2001, Miles et al. 2018); overall biodiversity may be elevated compared to the city's surrounding area due to habitat heterogeneity, although non-native species will abound (McKinney 2002, Grimm et al. 2008, Parsons et al. 2018); lethal exploitation is typically highly regulated, limiting humans' super predatory role to particular species (e.g., pest species; Adams 2016); top predators are rare, resulting in increased bottom-up control of trophic dynamics (Faeth et al. 2005, Estes et al. 2011); resource availability is elevated and seasonally stabilized due to landscaping and water provisioning, which intensifies resource competition and favors generalists species tolerant of humans (Faeth et al. 2005, McKinney 2006). Additionally, nonlethal human activity in cities is very high, especially along transportation routes and areas of congregation such as commercial shopping centers. This activity triggers behavioral adaptations in wildlife (e.g., increased nocturnality; Gaynor et al. 2018, Moll et al. 2018), and vehicular


Human Eltonian niche

traffic acts as a major source of animal mortality (Bateman and Fleming 2012) and a conduit for translocation events (Bullock et al. 2018). Thus, the predominant Eltonian role played by humans in this system is that of a niche constructor via intense and long-lasting landscape modification in the form of the built environment and the high-volume, daily-pulsed nonlethal human activity it supports (Pickett et al. 2011).

The Eltonian niche of humans in a large reserve differs markedly from that described above. Assuming effective regulation exists, lethal exploitation in the reserve is low overall but occasionally present and targeted (e.g., hunting a closely managed species; Wright et al. 2006). Landscape development is likewise relatively minimal, typically taking the form of road development, fencing, or waterhole provisioning (Valeix et al. 2009, Bojarska et al. 2017, Bull et al. 2018). Nonlethal human activity depends on recreation opportunity and interest, but is much lower than an urban context and exhibits temporal pulsing across tourist seasons (Newsome et al. 2012). Translocation events are infrequent but could be ecologically influential when exotic species are removed or keystone species are reintroduced (Hayward et al. 2007, Beschta and Ripple 2016). The gestalt ecological result of these collective actions is a relatively “intact” trophic structure where species interactions and food web dynamics more closely resemble those expected prior to human settlement of an area (although this will vary according to ecological and socio-cultural context; Jones 1969, Delcourt 2004). In contrast to the city, the predominant Eltonian role played by humans in such a reserve is that of a pseudo-mutualist via providing space for wildlife and plants that benefit of human well-being through the mechanisms of tourism, recreation, existence value, and ecosystem services.

FIGURES


Figure 1. A framework for the Eltonian niche of humans: *human decision-making* drives *human actions* that result in *ecological and evolutionary effects*. Human actions and their effects are characterized by several key *qualities*. The combination of human actions, ecological effects, and their qualities often lead to humans playing one or more of five predominant and unique *Eltonian roles* in a given system.



Human Eltonian niche

Figure 2. A detailed view of the Eltonian niche of humans. Arrows indicate causal processes.

Eltonian roles (1-5) are special cases of broader categories in the framework, as follows: 1) *super predation* occurs when lethal exploitation or human-induced fear is of large magnitude, 2) landscape modification entails *niche construction* if resultant selective pressures persist across generations, 3) humans act as a *hyperkeystone* species when trophic effects are propagated via keystone species, 4) *risk-response* occurs when fear modifies human behavior, and 5) *pseudo-mutualism* occurs when humans benefit species that provides desirable services or value to humanity.



Human Eltonian niche

Figure 3. Incorporating humans into classic food web modules. Diagrams depict direct lethal and nonlethal effects (solid and double-lined arrows) and positive (+) and negative (-) indirect effects (dashed arrows). H = Human, C = Consumer Species, R = Resource Species. Panels: (a) human fishing of herring (*Clupea harengus*) results in exploitative competition with grey seals (*Halichoerus grypus*); (b) human exploitation of coyotes (*Canis latrans*) positively affects their prey, black-tailed jackrabbits (*Lepus californicus*); (c) apparent competition from human exploitation of sympatric ungulate species, white-tailed (*Odocoileus virginianus*) and mule deer (*Odocoileus hemionus*); and (d) a human shield effect provides elk (*Cervus canadensis*) protection from human-averse wolves (*Canis lupus*). Images: Public Domain.





Figure 4. Examples of human actions across categories and qualities. In each colored graph, the spatial scale of the action increases vertically while intensity increases horizontally. The temporal nature of actions include singular or occasional, pulsed across various timescales, or relatively continuous. Colored arrows depict human actions that bridge categories.



Human Eltonian niche

Figure 5. Humans' Eltonian roles. Dashed arrows indicate effects that are behaviorally rather than numerically-mediated. Panels: (a) humans can act as super predators via lethal exploitation or (b) by inducing fear in apex predators (e.g., mountain lion *Puma concolor*); (c) humans can act as hyperkeystone species, as in gray wolf (*Canis lupus*) reintroduction into Yellowstone National Park, USA; (d) humans act as niche constructors by altering selective pressures via urbanization; (e) humans act as pseudo-mutualists by providing nest boxes for raptors that hunt agricultural pests; (f) humans act as risk-responders that avoid threatening species (e.g., the *Ixodes* ticks that act as a vector for Lyme disease). Images: Panel e – Florida Fish and Wildlife; others – Public Domain.

