

Routine surgical management of PAS without previa

	USA & Canada n=21	Latin & South America n=60	Oceania n=4	Europe =24	Western & Central Asia n=4	Southern, Eastern & Southeastern Asia n=15	Africa n=6
If PAS is confirmed intraoperatively, is there routinely an attempt to remove the placenta?							
Yes	1 (4.8)	12 (20.0)	0 (0.0)	3 (12.5)	0 (0.0)	6 (40.0)	3 (50.0)
If PAS is confirmed intraoperatively, is FIGO grading performed?							
Yes	6 (28.6)	38 (63.3)	1 (25.0)	14 (58.3)	2 (50.0)	12 (80.0)	4 (66.7)
Are preoperative vascular occlusion devices used?							
Yes	1 (4.8)	4 (6.7)	0 (0.0)	1 (4.2)	0 (0.0)	1 (6.7)	1 (16.7)
No	19 (90.5)	54 (90.0)	4 (100.0)	20 (83.3)	4 (100.0)	13 (86.7)	5 (83.3)
Are intraoperative vascular occlusion devices used?							
Yes	2 (9.5)	8 (13.3)	0 (0.0)	2 (8.3)	1 (25.0)	1 (6.7)	1 (16.7)
No	18 (85.7)	48 (80.0)	4 (100.0)	20 (83.3)	3 (75.0)	13 (86.7)	5 (83.3)
Are preoperative ureteral stents/catheters used?							
Yes	5 (23.8)	18 (30.0)	1 (25.0)	0 (0.0)	2 (50.0)	0 (0.0)	0 (0.0)
No	15 (71.4)	37 (61.7)	3 (75.0)	20 (83.3)	2 (50.0)	15 (93.3)	6 (100.0)
Surgical skin incision							
Vertical midline	16 (76.2)	37 (61.7)	1 (25.0)	9 (37.5)	2 (50.0)	8 (53.3)	3 (50.0)
Pfannenstiel	5 (23.8)	17 (28.3)	1 (25.0)	9 (37.5)	1 (25.0)	6 (40.0)	3 (50.0)
Maylard	0 (0.0)	2 (3.3)	1 (25.0)	1 (4.2)	1 (25.0)	0 (0.0)	0 (0.0)
Cherney	0 (0.0)	1 (1.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Other	0 (0.0)	2 (3.3)	1 (25.0)	4 (16.7)	0 (0.0)	1 (6.7)	0 (0.0)
Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Uterine incision							
Low transverse	3 (14.3)	15 (25.0)	2 (50.0)	10 (41.7)	2 (50.0)	7 (46.7)	1 (16.7)
High transverse	2 (9.5)	7 (11.7)	1 (25.0)	2 (8.3)	0 (0.0)	4 (26.7)	2 (33.3)
Classical	7 (33.3)	9 (15.0)	1 (25.0)	2 (8.3)	0 (0.0)	1 (6.7)	1 (16.7)
Fundal	2 (9.5)	19 (31.7)	0 (0.0)	4 (16.7)	0 (0.0)	1 (6.7)	1 (16.7)
Other	7 (33.3)	10 (16.7)	0 (0.0)	5 (20.8)	2 (50.0)	2 (13.3)	1 (16.7)
Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (4.2)	0 (0.0)	0 (0.0)

Data presented as n (%)