

Original Article

Pilot Study on the Current Management of Children with COVID-19 In Hospitals in Bangladesh; Findings and Implications

Nadia Nusrat¹, Mainul Haque², Kona Chowdhury³, Nihad Adnan⁴, Afzalunnessa Binte Lutfor⁵, Enamul Karim⁶, Mahmuda Hassan⁷, Arif Rabbany⁸, Dilara Begum⁹, Mohammad Nazmul Hasan¹⁰, Nazmus Sihan¹¹, Sifat Uz Zaman⁴, Salequl Islam⁴, Natalie Schellack¹², Marshall Gowere¹², Amanj Kurdi^{13a, b, c}, Brian Godman^{14a, b, c}

Abstract

Objective: The focus on COVID-19 in children in low- and middle-income countries including Bangladesh has been on addressing key issues including poor vaccination rates as well as mental health issues, domestic violence and child labour. However, the focus on optimally managing children in hospitals is changing with new variants and concerns with the development of hyperinflammatory syndromes. There are also concerns with the overuse of antimicrobials to treat patients with COVID-19 in hospitals enhancing resistance rates. The Bangladesh Paediatric Association have developed guidelines to improve patient care building on national guidance. Consequently, there is a need to document the current management of children with COVID-19 in Bangladesh and use the findings for future guidance. **Methods:** Rapid analysis of the management of children with COVID-19 among eight private and public hospitals in Bangladesh with varying numbers of in-patient beds using purposely developed case report forms (CRFs). The CRFs were piloted before full roll-out. **Results:** Overall low numbers of children in hospital with COVID-19 (4.3% of in-patient beds). The majority were male (59.6%) and aged 5 years or under (63.5%). Reasons for admission included respiratory distress/ breathing difficulties with 94.2% of COVID-19 cases confirmed. All children were prescribed antibiotics empirically, typically those on the Watch list of antibiotics and administered parenterally, with only a small minority switched to oral therapy before discharge. There was appreciable prescribing of Vitamins (C and D) and zinc and encouragingly limited prescribing of other antimicrobials (antivirals, antimalarials and antiparasitic medicines). Length of stay was typically 5 to 10 days. **Conclusion:** Encouraging to see low hospitalisation rates and limited use of antimicrobials apart from antibiotics. Concerns with high empiric use of antibiotics and limited switching to oral formulations can be addressed by instigating antimicrobial stewardship programmes. We will be monitoring this.

Keywords: Antimicrobials, Bangladesh, Children, COVID-19, Guidelines, Hospitals, Vitamins

Bangladesh Journal of Medical Science, Special Issue on COVID-19, 2021. Page : 188-198
DOI: <https://doi.org/10.3329/bjms.v20i5.55615>

1. Department of Paediatrics, Delta Medical College and Hospital, 26/2, Principal Abul Kashem Road, Mirpur-1, Dhaka-1216, Bangladesh. Orcid ID: <https://orcid.org/0000-0003-0133-0303>
2. Unit of Pharmacology, Faculty of Medicine and Defence Health, Universiti Pertahanan Nasional Malaysia (National Defence University of Malaysia), Kem Sungai, Besi, 57000 Kuala Lumpur, Malaysia. Orcid ID: <https://orcid.org/0000-0002-6124-7993>
3. Department of Paediatrics, Gonoshasthaya Samaj Vittik Medical College and Hospital, Dhaka, Bangladesh. Orcid ID: <https://orcid.org/0000-0002-3836-1691>
4. Department of Microbiology, Jahangirnagar University, Savar, Dhaka-1342, Bangladesh. Orcid ID: <https://orcid.org/0000-0002-4999-4793> (NA), <https://orcid.org/0000-0001-6061-2488> (SUZ) and <https://orcid.org/0000-0001-6131-4132> (SI)
5. Department of Microbiology, Ad-Din Women's Medical College, 2 Boro Mogbazar, Dhaka, Bangladesh. Orcid ID: <https://orcid.org/0000-0002-0176-8746>
6. Department of Paediatrics, US-Bangla Medical College, Kornogop, Tarabo, Rupgonj, Narayangonj, Dhaka, Bangladesh.
7. Department of Paediatrics, Ad-din Women's Medical College, 2 Boro Mogbazar, Dhaka, Bangladesh. Orcid ID: <https://orcid.org/0000-0001-9935-3460>
8. Department of Paediatrics, Mymensingh Medical College Hospital, Dhaka-Mymensingh Road, Mymensingh Sadar, Mymensingh, Bangladesh. Orcid ID: <https://orcid.org/0000-0001-5037-6346>
9. Department of Paediatrics, Dhaka Medical College Hospital, 100 Ramna Central Shaheed Minar Area, Bakshi Bazar, Dhaka, Bangladesh. Orcid ID: <https://orcid.org/0000-0002-8165-4203>

Correspondence: Brian Godman, Department of Pharmacoepidemiology, Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, UK, Email: brian.godman@strath.ac.uk

1. Introduction

COVID-19 was first identified in Wuhan, China, in December 2019^{1,2}, and towards the end of August 2021, there were already 213.7 million recorded cases worldwide with just under 4.5 million recorded deaths giving a case fatality ratio (CFR) of 2.09%³. This includes Bangladesh where 1.48 million cases had been recorded towards the end of August 2021 with over 25 thousand deaths giving a CFR of 1.73%³.

Since COVID-19 is principally spread through physical contact and airborne transmission^{4,5}, the focus in the early stages of the pandemic was its prevention especially with an appreciable number of patients being asymptomatic^{1,5}. Effective and cost-effective strategies included quarantining and lockdown measures incorporating the closure of borders as well as social distancing^{4,6-8}. However, varying degrees of implementation resulted in appreciable differences in prevalence and mortality rates among different African and Asian countries as well as with Western European countries and the United States of America⁹⁻¹¹.

Various re-purposed medicines were also suggested to reduce morbidity and mortality in patients with COVID-19 including hydroxychloroquine, lopinavir/ritonavir and remdesivir. However, these three had little impact on patient outcomes when objectively trialled in an appreciable number of patients, with potential issues with hydroxychloroquine increasing morbidity, mortality and costs^{4,12-17}. There is also ongoing controversy surrounding the potential use of ivermectin¹⁸⁻²¹. Concerns generally with the level of misinformation regarding potential approaches to prevention and treatment resulted in respected bodies including the British Medical Journal, European Centre for Disease Prevention and Control (ECDC) and the World Health Organisation (WHO) providing guidance from March 2020 onwards²²⁻²⁶. As a result, encouraging the implementation of evidence-based

approaches^{1,27}. This was helped in Bangladesh by the Ministry of Health and Family Welfare making national guidelines available from Spring 2020 onwards²⁸. This included guidelines for managing patients in hospital depending on their severity, which were subsequently built upon as more data became available. In addition, guidance on how to run a COVID-19 Hospital in Bangladesh²⁹.

Similarly, separate guidelines have been developed for managing paediatric patients with COVID-19 in Bangladesh³⁰. This is in addition to international guidelines developed to improve the management of children with COVID-19 in paediatric intensive care units (PICUs) in both high- and low-income countries³¹. This is important as adherence to trusted guidelines are seen to enhance the quality of care, with adherence increasingly seen as a robust marker of the quality of care to improve outcomes including patients in hospital with infectious diseases³²⁻³⁶, with up-to-date and easy-to-use guidelines facilitating their use³⁷.

Other developments in Bangladesh to improve current and future management of patients with COVID-19 include assessing antibody kinetics in these patients as well as performing genomic sequencing of different SARS-CoV-2 strains detected among COVID-19 patients³⁸⁻⁴⁰.

However, we are aware that children have a lower risk of infection with COVID-19 and milder clinical manifestations when they become infected compared with adults⁴¹⁻⁴⁶. Typically, fever, cough, respiratory infections, nausea and diarrhoea are the most frequent clinical characteristics of children with COVID-19, with males typically more prone to COVID-19 than females^{41,44-48}. In addition, an appreciable number of children are asymptomatic^{44,48}. Overall, approximately 6% to 10% of infected children with COVID-19 experience severe disease as compared to as high as 26% of adults who progress to ICUs^{44,45,49}.

10. Department Paediatric Surgery, Cumilla Medical College Hospital, Cumilla, Bangladesh.

11. Department of Paediatrics, Cumilla Medical College Hospital, Cumilla, Bangladesh.

12. Department of Pharmacology, University of Pretoria, Pretoria, South Africa. Orcid ID: (NS); <https://orcid.org/0000-0003-2154-3023> (MG)

13. ^aStrathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, United Kingdom; ^b Division of Public Health Pharmacy and Management, School of Pharmacy, Sefako Makgatho Health Sciences University, Pretoria, South Africa; ^c Department of Pharmacology, College of Pharmacy, Hawler Medical University, Erbil, Iraq. Orcid ID: <https://orcid.org/0000-0001-5036-1988>

14. ^a Division of Public Health Pharmacy and Management, School of Pharmacy, Sefako Makgatho Health Sciences University South Africa; ^b Department of Pharmacoepidemiology, Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, UK, and ^c School of Pharmaceutical Sciences, Universiti Sains Malaysia, Penang, Malaysia. Orcid ID: <https://orcid.org/0000-0001-6539-6972>

The lower susceptibility of children to COVID-19 infection may be due to elevated antibodies compared with adults along with other factors⁵⁰.

Consequently, the focus in children in low- and middle-income countries (LMICs) including Bangladesh has typically been addressing key concerns including poor vaccination rates as a result of lockdown and other activities⁵¹⁻⁵³. In Bangladesh, this includes organising immunisation outreach services, tracking children who had missed their vaccinations as well as increasing home visits by health workers^{53,54}. In addition, addressing some of the mental health challenges arising from lockdown measures including increasing domestic violence and child labour⁵⁵⁻⁶⁰.

However, the focus on the optimal management of children with COVID-19 in hospitals is increasing especially with new variants. Whilst a lower percentage of children with severe symptoms are admitted to intensive care units (ICUs) in LMICs versus high-income countries, deaths among hospitalised children are higher⁴⁵. This may be because of manpower issues. For instance, Bangladesh has the second lowest doctor-to-patient ratio in South Asia making managing patients with COVID-19 in hospitals challenging⁶¹. Consequently, monitoring current management approaches as well as providing guidance should help improve the future care of these patients.

Having said this, we are aware of the high use of antimicrobials in hospitals including intensive care units in treating patients with COVID-19 despite only a limited number of patients have concomitant fungal or bacterial infections⁶²⁻⁶⁹. This includes hospitals in Bangladesh^{70,71}. As a result, potentially increasing rates of antimicrobial resistance (AMR) with its associated increase in morbidity, mortality and costs^{66,72-77}. The instigation of antimicrobial stewardship programmes (ASPs) should help to address key concerns as these programmes have reduced inappropriate antibiotic utilisation among a number of LMICs^{72,73,78,79}, with educational and other initiatives also reducing inappropriate antibiotic prescribing to prevent surgical site infections among LMICs^{72,80}.

We see though that children diagnosed with COVID-19 do develop Kawasaki Disease (KD)-like symptoms/ hyperinflammatory syndromes^{41,81-83}, which is reflected by higher C-reactive protein and erythrocyte sedimentation rates and children experiencing leucopenia with marked lymphopenia, hypoalbuminemia, and hyponatremia along with gastrointestinal changes, abnormal electrocardiogram (ECG) and respiratory changes⁸¹, which can result in the need for intensive care⁸¹. This is important as we know from research with adults with COVID-19 in ICUs in Bangladesh that over 50% died whilst in ICU⁸⁴.




Figure 1: Rationale for The Pilot Study

As a result of growing concerns with the number of children with COVID-19 in Bangladesh especially those with symptoms, coupled with concerns with the overuse of antimicrobials, we believed it was important to document the current situation among paediatric patients hospitalized in Bangladesh as a result of COVID-19⁶¹ (Figure 1).

The findings can be utilised to improve future management building on the earlier study of Hussain *et al.* (2020) and the recent guidelines^{30,53}. This was the objective of this pilot study. We believed this was important given the current higher mortality rates in children with COVID-19 in LMICs versus high-income countries⁴⁵.

2. Methodology

A rapid analysis was undertaken among eight public and private hospitals in Bangladesh with an appreciable range of in-patient paediatric beds treating children with COVID-19. Case report forms were purposely developed for this study based on the published literature as well as the experience of the co-authors^{30,44,45,53,85}. These were piloted before roll-out between Mid July 2021 and Mid August 2021.

Key data sets based on the past ten days was included (10 days was chosen due to potentially low numbers of children with COVID-19, especially in general hospitals):

- Number of paediatric patients typically being treated in the hospital
- % of children with COVID-19 in the hospital
- Patient's ages in years (3 bands: 0 – 5 years; 6 – 10 years; 11 – 18 years) and Gender
- Principal reasons for admittance (3 principal reasons – drop-down menu of breathing difficulties/respiratory distress; prolonged fever; cough; diarrhoea; and feeding difficulty/vomiting)
- Comorbidities (mention top 3)
- % of these patients admitted to PICU and the reasons – taken from severe respiratory distress/ low O₂ saturation; shock; coagulation disorders/ thromboembolic manifestations; and extensive lung involvement in high-resolution CT scan (HRCT)
- % of prescribed antimicrobials (which ones – ATC Grouping) and % empiric
- Route of administration, when prescribing re-assessed and total length of prescription

- Whether any antivirals (e.g., remdesivir), antimalarials (e.g., hydroxychloroquine), anti-inflammatories (e.g., dexamethasone) or supplements/ immune boosters such as Vitamins C or D or zinc administered
- Recovery – mortality, morbidity or recovery
- Possible costs

We also assessed antibiotic prescribing against the WHO Access, Watch, or Reserve (AWaRe) list, which is increasingly being used to assess the quality of antimicrobial prescribing^{86,87}. In addition, where documented, against key guidance contained with the recent guidance from the Bangladesh Paediatric Association (Box 1)³⁰.

Box 1: Key recommendations from Bangladesh Paediatric Association on the management of children with COVID-19 in hospital [Adapted from³⁰]

- Ideally PCR test to confirm the diagnosis and subsequent management on a designated COVID-19 ward or treatment area.
- Comorbid conditions can include bronchial asthma, chronic kidney or liver disease and rheumatological conditions including Kawasaki disease.
- General treatment includes supportive care, electrolyte balancing and providing oxygen when necessary.
- Children transferred to ICU if experiencing severe/critical symptoms, respiratory failure requiring mechanical ventilation, shock and organ failure. Treatment includes vasoactive drugs if required as well as balanced/ buffered crystalloids.
- Medicines:
 - Antivirals – potentially reserved for children with severe acute respiratory syndrome. If remdesivir – then ideally part of ongoing clinical trials. Conflicting evidence regarding lopinavir/ ritonavir.
 - Antibiotics – specific cases only avoiding excessive use. 1st line – ampicillin plus gentamicin; 2nd line ceftriaxone.
 - Corticosteroids – not used routinely. Low dose dexamethasone may be beneficial in children with severe disease.
 - Vitamins – may be beneficial.
- For suspected Kawasaki Disease/ MIS-C – typically admit to PICU with supportive care including antibiotics for suspected infections as well as steroids and other immune modifying therapies.

The co-authors orchestrated ethical approval for their hospitals and others involved in the study. The reference numbers were: ibh/mirpur/2020/01.

Date: 31/08/2021; GSVMC/2021/346, Date: August 22, 2021; BBEC, JU/M 2021/COVID-19/7, Date: 18/7/2021 and AVMC/IRB-21 July 2021/028, Date: August 10, 2021.

3. Results

Overall, there were low numbers of children with COVID-19 among the eight hospitals taking part

in the pilot study at 4.3% of paediatric in patients (Table 1). The majority were male (59.6%) and aged between 0 to 5 years (63.5%). The principal reasons for admission for actual or suspected COVID-19 included respiratory distress/ breathing difficulties, prolonged fever, cough and feeding difficulties. Overall, 94.2% of cases were confirmed.

Table 1: Key Characteristics of The Children Among the Eight Hospitals

Hospital	Date Survey Conducted	Total Patients Childrens' Wards	Total with COVID-19	% with COVID-19	Number confirmed (tests) and %	Male	Female	0 to 5 years	6 to 10 years	11 to 18 years
Hosp 1	11.08.2021	65	3	4.6%	3 (100%)	2	1	3	0	0
Hosp 2	31.08.2021	125	5	4.0%	5 (100%)	4	1	2	3	0
Hosp 3	30.07.2021	19	2	10.5%	1 (one suspected)	1	1	1	1	0
Hosp 4	31.07.2021	210	26	12.4%	26 (100%)	15	11	13	6	7
Hosp 5	16.07.2021	32	2	6.3%	0 (two suspected)	1	1	2	0	0
Hosp 6	25.07.2021	18	1	5.6%	1 (100%)	1	0	0	1	0
Hosp 7	17.07.2021	654	9	1.4%	9 (100%)	5	4	9	0	0
Hosp 8	10.08.2021	85	4	4.7%	4 (100%)	2	2	3	1	0
Total		1208	52	4.3%	49 (94.2%) -3 suspected	31	21	33	12	7

Only a minority of those with COVID-19 were admitted to PICU (17.3%), with the principal reasons being severe respiratory distress/ low O₂ saturation and shock.

All of children with COVID-19 received antibiotics, which was typically empiric and principally parenterally (Table 2), for typically 5 to 10 days, with children generally reviewed after 3 to 5 days. However, only a small minority of children were switched to oral therapy before discharge (2%). There was also appreciable prescribing of Vitamins (C and D) and zinc with limited prescribing of antimalarials, antivirals and antiparasitic medicines (Table 2). Steroids were also administered to children in three of the hospitals.

Typically, the antibiotics prescribed were on the Watch as opposed to the Reserve list of antibiotics (Table 2). Encouragingly, mortality rates were low (under 2% of children) with the vast majority of children (88.5%) making a full recovery from COVID-19. Where measured, 4 out of the 5 hospitals surveyed stated that they adhered to the Bangladesh Paediatric Guidelines when managing their patients. Having said this, there was appreciable empiric use of antibiotics in all hospitals (Table 2). In addition, where antibiotics were administered there was

limited use of ampicillin although greater use of aminoglycosides including gentamicin (Table 2).

Length of stay in hospitals was typically 5 to 10 days (range 5 to 15 days), with total costs when available ranging between 3,000 and 20,000 BDT.

4. Discussion

We believe this is the first study conducted in Bangladesh among a number of both private and public hospitals regarding the management of hospitalised children with COVID-19. This builds on the earlier study of Hussain *et al.* (2020) who found a higher mortality rate (13.3%) among children in the early stages of the pandemic⁵³. This may reflect improved management as more knowledge is gained regarding the aetiology of COVID-19 and optimal treatment approaches. In addition, only a low number of paediatric patients in hospital had COVID-19 (Table 1). This is important when looking to increase services such as elective paediatric surgery which was put on hold in the early stages of the pandemic⁸⁸.

Table 2: Management of Children With COVID-19 Among the Eight Hospitals

Hospital	Number and % Rx antimicrobials	Empiric or CST	Antibiotic Rx	Antiviral	Anti-parasitic	Anti-inflammatory (steroid)	Supplements, etc
Hosp 1	3 (100%)	All empiric	Ceftriaxone	No	No	No	Vit C, D, Zinc
Hosp 2	5 (100%)	All empiric	Aminoglycosides, carbapenems, cephalosporins	No	No	No	No
Hosp 3	2 (100%)	All empiric	Ceftriaxone, ciprofloxacin	No	No	No	Vit C, D, Zinc
Hosp 4	100%	All empiric	Ceftriaxone, ceftazidim, meropenem, vancomycin, flucloxacillin, amikacin	No	No	Steroids	Vit C, D, Zinc
Hosp 5	2 (100%)	All empiric	Ampicillin, aminoglycoside, cephalosporin	No	No	No	No
Hosp 6	1 (100%)	All empiric	Ceftriaxone	No	No	No	Vit C, D, Zinc
Hosp 7	9 (100%)	All empiric	Aminoglycosides, carbapenem, macrolides	No	No	Dexamethasone	Vit C, D, Zinc
Hosp 8	4 (100%)	All empiric	Aminoglycosides, carbapenems, cephalosporins	No	Ivermectin	Steroids	Vit C, D, Zinc

Overall, it was encouraging to see low hospitalisation rates for children with COVID-19 (Table 1) reflecting other recent publications^{44,46,48}. Alongside this, similar to other published studies, fever and cough as well as respiratory diseases were the most frequent clinical characteristics of children admitted with COVID-19, with again boys accounting for more cases of COVID-19 in hospital than girls^{41,44-48}.

Encouragingly there was limited use of antimalarials, antivirals and anti-parasitic medicines (Table 2) given concerns with their lack of effectiveness in treating patients with COVID-19 despite the hype^{12-15,20,89}. In addition, the high use of vitamins and zinc is in line with current national guidelines in Bangladesh (Box 1). However, there were concerns with the high empiric use of antibiotics (Table 2) despite, as mentioned, only a limited number of patients across ages have concomitant fungal or bacterial infections⁶²⁻⁶⁹. In addition, concerns with overuse are documented in the current national guidelines (Box 1). This is similar to the high use of antibiotics among adults admitted to hospital with COVID-19 in Bangladesh^{70,71}. This is despite clinicians in most of the surveyed hospitals stating they follow the current national guidelines.

Potential ways forward to improve antimicrobial use include the instigation of ASPs, which have been successful across countries to enhance the appropriate use of antibiotics in hospitals including LMICs^{72,73,78,79,90}. Future activities should also include encouraging prescribing where pertinent according to local antibiograms to improve empiric use along with the instigation of ASPs^{91,92}. These combined activities should also reduce the prescribing of ‘Watch antibiotics in favour of increased prescribing of Access antibiotics’^{93,94}. In addition, encourage earlier switching from parental to oral antibiotics to hasten earlier discharge from hospital and conserve resources⁹⁵⁻⁹⁷. We will be following these activities in the future.

Limitations

We are aware of a number of limitations with this study. This included the fact that this was only a pilot study. In addition, the study involved a retrospective review of notes without the ability to check with physicians regarding the rationale for their treatment approaches and all cases of COVID-19 could not be confirmed by rt-PCR. Despite these limitations, we believe the findings are robust providing direction for the future.

5. Conclusions

It was encouraging to see low hospitalisation rates for children with COVID-19 among the pilot hospitals in Bangladesh, which mirrors findings in other countries. It was also encouraging to see limited use of antimicrobials apart from antibiotics given the hype that has surrounded some of these. However, there are concerns with the high empiric use of antibiotics and limited switching to oral formulations. This needs to be addressed to reduce

the potential for increasing antimicrobial resistance and costs. This can be addressed by instigating or enhancing antimicrobial stewardship programmes in hospitals. We will be monitoring this in the future.

Conflicts of interest

The authors declare they have no conflicts of interest

Funding

There was no funding for this project – it was self funded by the authors.

References

- Godman B. Combating COVID-19: Lessons learnt particularly among developing countries and the implications. *Bangladesh Journal of Medical Science, Special Issue on Covid19*. **2020**; S103-8. DOI: <https://doi.org/10.3329/bjms.v19i0.48413>.
- Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. **2020**;323(13):1239-42, doi: 10.1001/jama.2020.2648
- World Health Organization. WHO Coronavirus (COVID-19) Dashboard. 26 August 2021. Available at URL: <https://covid19.who.int/>. (26 August, 2021)
- Haque M, Gowere M, Nusrat N, Chowdhury K, Godman B. The response to COVID 19 across countries and the implications for future pandemics. *Bangladesh Journal of Medical Science*. **2021**; 20 (Special Issue on Covid 19): S7-14.
- Ayouni I, Maatoug J, Dhouib W, Zammit N, Fredj SB, Ghammam R, et al. Effective public health measures to mitigate the spread of COVID-19: a systematic review. *BMC Public Health*. **2021**;21(1):1015. DOI: 10.1186/s12889-021-11111-1
- Courtemanche C, Garuccio J, Le A, Pinkston J, Yelowitz A. Strong Social Distancing Measures In The United States Reduced The COVID-19 Growth Rate. *Health Aff (Millwood)*. **2020**;39(7):1237-46. doi: 10.1377/hlthaff.2020.00608
- Rezapour A, Souresrafil A, Peighambari MM, Heidarali M, Tashakori-Miyanroudi M. Economic evaluation of programs against COVID-19: A systematic review. *International journal of surgery*. **2021**;85:10-8. doi: 10.1016/j.ijssu.2020.11.015
- Mendez-Brito A, El Bcheraoui C, Pozo-Martin F. Systematic review of empirical studies comparing the effectiveness of non-pharmaceutical interventions against COVID-19. *J Infect*. **2021** (EPrint). doi: 10.1016/j.jinf.2021.06.018
- Godman B, Haque M, Islam S, Iqbal S, Urmi UL, Kamal ZM, et al. Rapid Assessment of Price Instability and Paucity of Medicines and Protection for COVID-19 Across Asia: Findings and Public Health Implications for the Future. *Front in Public Health*. **2020**;8(744). doi: 10.3389/fpubh.2020.585832
- Ogunleye OO, Basu D, Mueller D, Sneddon J, Seaton RA, Yinka-Ogunleye AF, et al. Response to the Novel Corona Virus (COVID-19) Pandemic Across Africa: Successes, Challenges, and Implications for the Future. *Front Pharmacol*. **2020**;11(1205). doi: 10.3389/fphar.2020.01205
- Sefah I, Ogunleye O, Essah D, Opanga S, Rizvi N, Wamaitha A, et al. Rapid assessment of the potential paucity and price increases for suggested medicines and protection equipment for COVID-19 across developing countries with a particular focus on Africa and the implications. *Front Pharmacol*. **2021**;11(2055). doi: 10.3389/fphar.2020.588106
- World Health Organization. WHO discontinues hydroxychloroquine and lopinavir/ritonavir treatment arms for COVID-19. 4 July 2020. Available at URL: <https://www.who.int/news-room/detail/04-07-2020-who-discontinues-hydroxychloroquine-and-lopinavir-ritonavir-treatment-arms-for-covid-19>. (26 August, 2021)
- RECOVERY Collaborative Group. Lopinavir-ritonavir in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet*. **2020**;396(10259):1345-52. DOI: 10.1016/S0140-6736(20)32013-4
- Horby P, Mafham M, Linsell L, Bell JL, Staplin N, Emberson JR, et al. Effect of Hydroxychloroquine in Hospitalized Patients with Covid-19. *N Engl J Med*. **2020**;383(21):2030-40. doi: 10.1056/NEJMoa2022926
- Charan J, Kaur RJ, Bhardwaj P, Haque M, Sharma P, Misra S, et al. Rapid review of suspected adverse drug events

- due to remdesivir in the WHO database; findings and implications. *Expert Rev Clin Pharmacol*. **2021**;14(1):95-103. doi: 10.1080/17512433.2021.1856655
16. Abena PM, Declodt EH, Bottieau E, Suleman F, Adejumo P, Sam-Agudu NA, et al. Chloroquine and Hydroxychloroquine for the Prevention or Treatment of COVID-19 in Africa: Caution for Inappropriate Off-label Use in Healthcare Settings. *Am J Trop Med Hyg*. **2020**;102(6):1184-8. doi: 10.4269/ajtmh.20-0290
 17. Haque M, Abubakar A, Ogunleye O, Sani I, Sefah I, Kurdi A, et al. Changes in availability, utilization, and prices of medicines and protection equipment for COVID-19 in an Urban population of Northern Nigeria. *J Res Pharm Pract*. **2021**;10(1):17-22. doi: 10.4103/jrpp.JRPP_20_92
 18. Ahmed S, Karim MM, Ross AG, Hossain MS, Clemens JD, Sumiya MK, et al. A five-day course of ivermectin for the treatment of COVID-19 may reduce the duration of illness. *Int J Infect Dis*. **2021**;103:214-6. DOI: 10.1016/j.ijid.2020.11.191
 19. Wadvalla BA. Covid-19: Ivermectin's politicisation is a warning sign for doctors turning to orphan treatments. *BMJ*. **2021**;373:n747. DOI: 10.1136/bmj.n747
 20. Garegnani LI, Madrid E, Meza N. Misleading clinical evidence and systematic reviews on ivermectin for COVID-19. *BMJ Evid Based Med*. **2021**:bmjebm-2021-111678. DOI: 10.1136/bmjebm-2021-111678
 21. Mahmud R, Rahman MM, Alam I, Ahmed KGU, Kabir A, Sayeed S, et al. Ivermectin in combination with doxycycline for treating COVID-19 symptoms: a randomized trial. *J Int Med Res*. **2021**;49(5):3000605211013550. DOI: 10.1177/03000605211013550
 22. World Health Organisation. Operational considerations for case management of COVID-19 in health facility and community: interim guidance, 19 March **2020**. Available at URL: <https://apps.who.int/iris/handle/10665/331492>. (26 August, 2021)
 23. World Health Organisation. Coronavirus disease (COVID-19) advice for the public: Myth busters. **2020**. Available at URL: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public/myth-busters>. (26 August, 2021)
 24. WHO. Countering misinformation about COVID-19. **2020**. Available at URL: <https://www.who.int/news-room/feature-stories/detail/countering-misinformation-about-covid-19>. (26 August, 2021)
 25. BMJ Best Practice - Coronavirus disease 2019 (COVID-19). 30 July **2020**. Available at URL: <https://bestpractice.bmj.com/topics/en-gb/3000168/pdf/3000168/Coronavirus%20disease%202019%20%28COVID-19%29.pdf>. (26 August, 2021)
 26. ECDC. Vaccines and treatment of COVID-19. **2020**. Available at URL: <https://www.ecdc.europa.eu/en/covid-19/latest-evidence/vaccines-and-treatment>. (26 August, 2021)
 27. Raimi MO, Mcfubara KG, Abisoye OS, Ifeanyi-chukwu Ezekwe C, Henry Sawyerr O, Raimi GA-anyam. Responding to the Call through Translating Science into Impact: Building an Evidence-Based Approaches to Effectively Curb Public Health Emergencies [Covid-19 Crisis]. *GJEID*. **2021**;1(1). Available from: <https://www.scipublications.com/journal/index.php/gjeid/article/view/72>. (26 August, 2021)
 28. Ministry of Health and Family Welfare. Government of the People's Republic of Bangladesh. National Guidelines on Clinical Management of Coronavirus Disease 2019 (COVID-19). May **2020**. Available at URL: http://file.portal.gov.bd/files/cs.sylhet.gov.bd/notices/245c5244_369b_4305_958d_1f70d99d6681/4c021caf6cf2915278aa3598e74f5c09.pdf. (26 August, 2021)
 29. Disease Control Division, Directorate General of Health Services, Ministry of Health & Family Welfare, Government of Bangladesh. How to Run a COVID-19 Hospital in Bangladesh. **2020**. Available at URL: <https://corona.gov.bd/storage/press-releases/April2020/g0MbdFSz5rwPzA8qhYFN.pdf>. (26 August, 2021)
 30. Bangladesh Paediatric Association (BPA). Management Guideline for Paediatric COVID-19. November **2020**. Available at URL: https://www.bpabd.org/asset/uploads/file/COVID_BPA_2nd%20ED_3.pdf. (26 August, 2021)
 31. Kache S, Chisti MJ, Gumbo F, Mupere E, Zhi X, Nallasamy K, et al. COVID-19 PICU guidelines: for high- and limited-resource settings. *Pediatr Res*. **2020**;88(5):705-16. DOI: 10.1038/s41390-020-1053-9
 32. Nakwatumbah S, Kibuule D, Godman B, Haakuria V, Kalemeera F, Baker A, et al. Compliance to guidelines for the prescribing of antibiotics in acute infections at Namibia's national referral hospital: a pilot study and the implications. *Expert Rev Anti Infect Ther*. **2017**;15(7):713-21. DOI: 10.1080/14787210.2017.1320220
 33. Versporten A, Zarb P, Caniaux I, Gros MF, Drapier N, Miller M, et al. Antimicrobial consumption and resistance in adult hospital inpatients in 53 countries: results of an internet-based global point prevalence survey. *The Lancet Global health*. **2018**;6(6):e619-e29. DOI: 10.1016/S2214-109X(18)30186-4
 34. Campbell S, Meyer JC, Godman B. Why compliance to national prescribing guidelines is important especially across sub-Saharan Africa and suggestions for the future. *J Biomed Pharm Sci* 4 (**2021**): 316. Available at URL: <https://www.hilarispublisher.com/open-access/why-compliance-to-national-prescribing-guidelines-is-important-especially-across-subsaharan-africa-and-suggestions-for-t.pdf>. (26 August, 2021)
 35. Almatar M, Peterson GM, Thompson A, McKenzie D, Anderson T, Zaidi ST. Clinical Pathway and Monthly Feedback Improve Adherence to Antibiotic

- Guideline Recommendations for Community-Acquired Pneumonia. *PloS one*. **2016**;11(7):e0159467. DOI: 10.1371/journal.pone.0159467
36. Niaz Q, Godman B, Masseur A, Campbell S, Kurdi A, Kagoya HR, et al. Validity of World Health Organisation prescribing indicators in Namibia's primary healthcare: findings and implications. *International journal for quality in health care*. **2019**;31(5):338-45. DOI: 10.1093/intqhc/mzy172
 37. Niaz Q, Godman B, Campbell S, Kibuule D. Compliance to prescribing guidelines among public health care facilities in Namibia; findings and implications. *Int J Clin Pharm*. **2020**;42(4):1227-36. DOI: 10.1007/s11096-020-01056-7
 38. Adnan N, Khondoker MU, Rahman MS, Ahmed MF, Sharmin S, Sharif N, et al. Coding-Complete Genome Sequences and Mutation Profiles of Nine SARS-CoV-2 Strains Detected from COVID-19 Patients in Bangladesh. *Microbiol Resour Announc*. **2021**;10(10). DOI: 10.1128/MRA.00124-21.
 39. Jamiruddin MR, Haq MA, Tomizawa K, Kobatake E, Mie M, Ahmed S, et al. Longitudinal Antibody Dynamics Against Structural Proteins of SARS-CoV-2 in Three COVID-19 Patients Shows Concurrent Development of IgA, IgM, and IgG. *J Inflamm Res*. **2021**;14:2497-506. DOI: 10.2147/JIR.S313188
 40. Matin MA, Nahid AA, Sarkar B, Ullah MA, Araf Y, Adnan N, et al. Genome Sequences of Two Novel Coronavirus (SARS-CoV-2) Isolates from Dhaka, Bangladesh. *Microbiol Resour Announc*. **2021**;10(27):e0051121. DOI: 10.1128/MRA.00511-21
 41. Mantovani A, Rinaldi E, Zusi C, Beatrice G, Saccomani MD, Dalbeni A. Coronavirus disease 2019 (COVID-19) in children and/or adolescents: a meta-analysis. *Pediatr Res*. **2021**;89(4):733-7. DOI: 10.1038/s41390-020-1015-2
 42. Kanthimathinathan HK, Dhesi A, Hartshorn S, Ali SH, Kirk J, Nagakumar P, et al. COVID-19: A UK Children's Hospital Experience. *Hosp Pediatr*. **2020**;10(9):802-5. DOI: 10.1542/hpeds.2020-000208
 43. Mehta NS, Mytton OT, Mullins EWS, Fowler TA, Falconer CL, Murphy OB, et al. SARS-CoV-2 (COVID-19): What Do We Know About Children? A Systematic Review. *Clin Infect Dis*. **2020**;71(9):2469-79. DOI: 10.1093/cid/ciaa556
 44. Bhuiyan MU, Stiboy E, Hassan MZ, Chan M, Islam MS, Haider N, et al. Epidemiology of COVID-19 infection in young children under five years: A systematic review and meta-analysis. *Vaccine*. **2021**;39(4):667-77. DOI: 10.1016/j.vaccine.2020.11.078
 45. Irfan O, Muttalib F, Tang K, Jiang L, Lassi ZS, Bhutta Z. Clinical characteristics, treatment and outcomes of paediatric COVID-19: a systematic review and meta-analysis. *Arch Dis Child*. **2021**;106(5):440-8. DOI: 10.1136/archdischild-2020-321385
 46. Cui X, Zhao Z, Zhang T, Guo W, Guo W, Zheng J, et al. A systematic review and meta-analysis of children with coronavirus disease 2019 (COVID-19). *J Med Virol*. **2021**;93(2):1057-69. DOI: 10.1002/jmv.26398
 47. Islam MA, Kundu S, Alam SS, Hossain T, Kamal MA, Hassan R. Prevalence and characteristics of fever in adult and paediatric patients with coronavirus disease 2019 (COVID-19): A systematic review and meta-analysis of 17515 patients. *PloS One*. **2021**;16(4):e0249788. DOI: 10.1371/journal.pone.0249788
 48. Mansourian M, Ghandi Y, Habibi D, Mehrabi S. COVID-19 infection in children: A systematic review and meta-analysis of clinical features and laboratory findings. *Arch Pediatr*. **2021**;28(3):242-8. DOI: 10.1016/j.arcped.2020.12.008
 49. Singh T, Heston SM, Langel SN, Blasi M, Hurst JH, Fouda GG, et al. Lessons From COVID-19 in Children: Key Hypotheses to Guide Preventative and Therapeutic Strategies. *Clin Infect Dis*. **2020**;71(8):2006-13. doi: 10.1093/cid/ciaa547
 50. Schellack N, Coetzee M, Schellack G, Gijzelaar M, Hassim Z, Milne M, et al. COVID-19: guidelines for pharmacists in South Africa. *S Afr J Infect Dis*. **2020**;35(1):206. DOI: 10.4102/sajid.v35i1.206
 51. Abbas K, Procter SR, van Zandvoort K, Clark A, Funk S, Mengistu T, et al. Routine childhood immunisation during the COVID-19 pandemic in Africa: a benefit-risk analysis of health benefits versus excess risk of SARS-CoV-2 infection. *Lancet Glob Health*. **2020**;8(10):e1264-e72. DOI: 10.1016/S2214-109X(20)30308-9
 52. Gaythorpe KA, Abbas K, Huber J, Karachaliou A, Thakkar N, Woodruff K, et al. Impact of COVID-19-related disruptions to measles, meningococcal A, and yellow fever vaccination in 10 countries. *eLife*. **2021**;10. DOI: 10.7554/eLife.67023
 53. Hussain M, Al Mamun MA. COVID-19 in children in Bangladesh: Situation analysis. *Asia Pac J Paediatr Child Health*. **2020**; 3: 59-65.
 54. Rana S, Shah R, Ahmed S, Mothabbir G. Post-disruption catch-up of child immunisation and health-care services in Bangladesh. *The Lancet Infectious diseases*. **2021**;21(7):913. DOI: 10.1016/S1473-3099(21)00148-1
 55. Yeasmin S, Banik R, Hossain S, Hossain MN, Mahumud R, Salma N, et al. Impact of COVID-19 pandemic on the mental health of children in Bangladesh: A cross-sectional study. *Child Youth Serv Rev*. **2020**;117:105277. DOI: 10.1016/j.chilgyouth.2020.105277
 56. Mallik CI, Radwan RB. Impact of lockdown due to COVID-19 pandemic in changes of prevalence of predictive psychiatric disorders among children and adolescents in Bangladesh. *Asian J Psychiatr*. **2021**;56:102554. DOI: 10.1016/j.ajp.2021.102554
 57. Ahad MA, Parry YK, Willis E. Spillover Trends of Child Labor During the Coronavirus Crisis- an Unnoticed

- Wake-Up Call. *Front Public Health*. **2020**;8:488. DOI: 10.3389/fpubh.2020.00488
58. Rashid Soron T, Ashiq MAR, Al-Hakeem M, Chowdhury ZF, Uddin Ahmed H. Domestic Violence and Mental Health During COVID-19 in Bangladesh. *JMIR Form Res*. **2021** (EPrint). DOI: 10.2196/24624
 59. Sifat RI. Impact of the COVID-19 pandemic on domestic violence in Bangladesh. *Asian J Psychiatr*: **2020**;53:102393. DOI: 10.1016/j.ajp.2020.102393
 60. Opanasenko A, Lugova H, Mon AA, Ivanko O. Mental Health Impact of Gender-Based Violence Amid COVID-19 Pandemic: A Review. *Bangladesh Journal of Medical Science, Special Issue on COVID-19*. **2021**: S17 - S25. DOI: <https://doi.org/10.3329/bjms.v20i5.55396>.
 61. Ali MR, Hasan MA, Rahman MS, Billah M, Karmakar S, Shimu AS, et al. Clinical manifestations and socio-demographic status of COVID-19 patients during the second-wave of pandemic: A Bangladeshi experience. *J Infect Public Health*. **2021**. EPrint. DOI: 10.1016/j.jiph.2021.06.011
 62. Ul Mustafa Z, Salman M, Aldeyab M, Kow CS, Hasan SS. Antimicrobial consumption among hospitalized patients with COVID-19 in Pakistan. *SN Compr Clin Med*. **2021**:1-5. DOI: 10.1007/s42399-021-00966-5
 63. Saini V, Jain C, Singh NP, Alsulimani A, Gupta C, Dar SA, et al. Paradigm Shift in Antimicrobial Resistance Pattern of Bacterial Isolates during the COVID-19 Pandemic. *Antibiotics*. **2021**;10(8):954. DOI: 10.3390/antibiotics10080954
 64. Langford BJ, So M, Raybardhan S, Leung V, Soucy JR, Westwood D, et al. Antibiotic prescribing in patients with COVID-19: rapid review and meta-analysis. *Clin Microbiol Infect*. **2021**;27(4):520-31. DOI: 10.1016/j.cmi.2020.12.018
 65. Langford BJ, So M, Raybardhan S, Leung V, Westwood D, MacFadden DR, et al. Bacterial co-infection and secondary infection in patients with COVID-19: a living rapid review and meta-analysis. *Clin Microbiol Infect*. **2020**;26(12):1622-9. DOI: 10.1016/j.cmi.2020.07.016
 66. Adebisi YA, Alaran AJ, Okereke M, Oke GI, Amos OA, Olaoye OC, et al. COVID-19 and Antimicrobial Resistance: A Review. *Infectious diseases*. **2021**;14:11786337211033870. DOI: 10.1177/11786337211033870
 67. Cheng LS, Chau SK, Tso EY, Tsang SW, Li IY, Wong BK, et al. Bacterial co-infections and antibiotic prescribing practice in adults with COVID-19: experience from a single hospital cluster. *Ther Adv Infect Dis*. **2020**;7:2049936120978095. DOI: 10.1177/2049936120978095
 68. Abu-Rub LI, Abdelrahman HA, Johar A-RA, Alhussain HA, Hadi HA, Eltai NO. Antibiotics Prescribing in Intensive Care Settings during the COVID-19 Era: A Systematic Review. *Antibiotics*. **2021**;10(8):935. DOI: 10.3390/antibiotics10080935
 69. Grau S, Hernández S, Echeverría-Esnal D, Almendral A, Ferrer R, Limón E, et al. Antimicrobial Consumption among 66 Acute Care Hospitals in Catalonia: Impact of the COVID-19 Pandemic. *Antibiotics*. **2021**;10(8):943. DOI: 10.3390/antibiotics10080943
 70. Molla MMA, Yeasmin M, Islam MK, Sharif MM, Amin MR, Nafisa T, et al. Antibiotic Prescribing Patterns at COVID-19 Dedicated Wards in Bangladesh: Findings from a Single Center Study. *Infect Prev Pract*. **2021**:100134. DOI: 10.1016/j.infpip.2021.100134
 71. Mah EMS, Hassan MZ, Biswas M, Rahman F, Akhtar Z, Das P, et al. Use of Antimicrobials among Suspected COVID-19 Patients at Selected Hospitals, Bangladesh: Findings from the First Wave of COVID-19 Pandemic. *Antibiotics*. **2021**;10(6). DOI: 10.3390/antibiotics10060738
 72. Godman B, Egwuenu A, Haque M, Malande OO, Schellack N, Kumar S, et al. Strategies to Improve Antimicrobial Utilization with a Special Focus on Developing Countries. *Life*. **2021**;11(6):528. doi: 10.3390/life11060528
 73. Haque M, Godman B. Potential Strategies to Improve Antimicrobial Utilisation in Hospitals in Bangladesh Building on Experiences Across Developing Countries. *Bangladesh Journal of Medical Science*. **2021**; 19 (3): 469-77. DOI: <https://doi.org/10.3329/bjms.v20i3.52787>.
 74. Hsu J. How covid-19 is accelerating the threat of antimicrobial resistance. *BMJ*. **2020**;369:m1983. DOI: 10.1136/bmj.m1983
 75. Founou RC, Founou LL, Essack SY. Clinical and economic impact of antibiotic resistance in developing countries: A systematic review and meta-analysis. *PLoS One*. **2017**;12(12):e0189621. DOI: 10.1371/journal.pone.0189621
 76. Hofer U. The cost of antimicrobial resistance. *Nat Rev Microbiol*. **2019**;17(1):3. doi: 10.1038/s41579-018-0125-x
 77. Majumder MAA, Rahman S, Cohall D, Bharatha A, Singh K, Haque M, et al. Antimicrobial Stewardship: Fighting Antimicrobial Resistance and Protecting Global Public Health. *Infect Drug Resist*. **2020**;13:4713-38. DOI: 10.2147/IDR.S290835
 78. Nathwani D, Varghese D, Stephens J, Ansari W, Martin S, Charbonneau C. Value of hospital antimicrobial stewardship programs [ASPs]: a systematic review. *Antimicrob Resist Infect Control*. **2019**;8:35. doi: 10.1186/s13756-019-0471-0
 79. Akpan MR, Isemin NU, Udoh AE, Ashiru-Oredope D. Implementation of antimicrobial stewardship programmes in African countries: a systematic literature review. *J Glob Antimicrob Resist*. **2020**;22:317-24. doi: 10.1016/j.jgar.2020.03.009
 80. Mwita JC, Ogunleye OO, Olalekan A, Kalungia AC,

- Kurdi A, Saleem Z, et al. Key Issues Surrounding Appropriate Antibiotic Use for Prevention of Surgical Site Infections in Low- and Middle-Income Countries: A Narrative Review and the Implications. *Int J Gen Med.* 2021;14:515-30. DOI: 10.2147/IJGM.S253216
81. Mardi P, Esmaeili M, Irvani P, Abdar ME, Pourrostami K, Qorbani M. Characteristics of Children With Kawasaki Disease-Like Signs in COVID-19 Pandemic: A Systematic Review. *Front Pediatr.* 2021;9:625377. DOI: 10.3389/fped.2021.625377
82. Yasuhara J, Watanabe K, Takagi H, Sumitomo N, Kuno T. COVID-19 and multisystem inflammatory syndrome in children: A systematic review and meta-analysis. *Pediatr Pulmonol.* 2021;56(5):837-48. DOI: 10.1002/ppul.25245
83. Hobbs CV, Khaitan A, Kirmse BM, Borkowsky W. COVID-19 in Children: A Review and Parallels to Other Hyperinflammatory Syndromes. *Front Pediatr.* 2020;8:593455. DOI: 10.3389/fped.2020.593455
84. Saha A, Ahsan MM, Quader TU, Shohan MUS, Naher S, Dutta P, et al. Characteristics, management and outcomes of critically ill COVID-19 patients admitted to ICU in hospitals in Bangladesh: a retrospective study. *J Prev Med Hyg.* 2021;62(1):E33-e45. doi: 10.15167/2421-4248/jpmh2021.62.1.1838
85. Panfili FM, Roversi M, D'Argenio P, Rossi P, Cappa M, Fintini D. Possible role of vitamin D in Covid-19 infection in pediatric population. *J Endocrinol Invest.* 2021;44(1):27-35. DOI: 10.1007/s40618-020-01327-0
86. Hsia Y, Sharland M, Jackson C, Wong ICK, Magrini N, Bielicki JA. Consumption of oral antibiotic formulations for young children according to the WHO Access, Watch, Reserve (AWaRe) antibiotic groups: an analysis of sales data from 70 middle-income and high-income countries. *Lancet Infect Dis.* 2019;19(1):67-75. DOI: 10.1016/S1473-3099(18)30547-4
87. Skosana PP, Schellack N, Godman B, Kurdi A, Bennie M, Kruger D, et al. A point prevalence survey of antimicrobial utilisation patterns and quality indices amongst hospitals in South Africa; findings and implications. *Expert Rev Anti Infect Ther.* 2021:1-13. DOI: 10.1080/14787210.2021
88. Hannan J, Parveen MK, Hasan S. Pediatric surgical services in Bangladesh during the COVID 19 pandemic: How they are affected and how to overcome the backlog, keeping healthcare professionals safe. 2020. *Med Rxiv Preprint.* doi: <https://doi.org/10.1101/2020.08.16.20169979>.
89. Dyer O. Covid-19: Remdesivir has little or no impact on survival, WHO trial shows. *BMJ.* 2020;371:m4057. doi: 10.1136/bmj.m4057
90. van den Bergh D, Messina AP, Goff DA, van Jaarsveld A, Coetzee R, de Wet Y, et al. A pharmacist-led prospective antibiotic stewardship intervention improves compliance to community-acquired pneumonia guidelines in 39 public and private hospitals across South Africa. *Int J Antimicrob Agents.* 2020;56(6):106189. DOI: 10.1016/j.ijantimicag.2020.106189
91. Morales A, Campos M, Juarez JM, Canovas-Segura B, Palacios F, Marin R. A decision support system for antibiotic prescription based on local cumulative antibiograms. *J Biomed Inform.* 2018;84:114-22. DOI: 10.1016/j.jbi.2018.07.003
92. Klinker KP, Hidayat LK, DeRyke CA, DePestel DD, Motyl M, Bauer KA. Antimicrobial stewardship and antibiograms: importance of moving beyond traditional antibiograms. *Ther Adv Infect Dis.* 2021;8:20499361211011373. DOI: 10.1177/20499361211011373.
93. Hsia Y, Lee BR, Versporten A, Yang Y, Bielicki J, Jackson C, et al. Use of the WHO Access, Watch, and Reserve classification to define patterns of hospital antibiotic use (AWaRe): an analysis of paediatric survey data from 56 countries. *Lancet Glob Health.* 2019;7(7):e861-e71. DOI: 10.1016/S2214-109X(19)30071-3
94. Sharland M, Pulcini C, Harbarth S, Zeng M, Gandra S, Mathur S, et al. Classifying antibiotics in the WHO Essential Medicines List for optimal use-be AWaRe. *Lancet Infect Dis.* 2018;18(1):18-20. DOI: 10.1016/S1473-3099(17)30724-7
95. Broom J, Broom A, Adams K, Plage S. What prevents the intravenous to oral antibiotic switch? A qualitative study of hospital doctors' accounts of what influences their clinical practice. *J Antimicrob Chemother.* 2016;71(8):2295-9. DOI: 10.1093/jac/dkw129
96. Jones M, Huttner B, Madaras-Kelly K, Nechodom K, Nielson C, Bidwell Goetz M, et al. Parenteral to oral conversion of fluoroquinolones: low-hanging fruit for antimicrobial stewardship programs? *Infect Control Hosp Epidemiol.* 2012;33(4):362-7. DOI: 10.1086/664767
97. Cunha BA. Beyond IV to PO switch therapy: Oral antimicrobial therapy. *Infectious Disease News.* 2015. Available at URL: <https://www.healio.com/news/infectious-disease/20151014/beyond-iv-to-po-switch-therapy-oral-antimicrobial-therapy> (26 August, 2021)