

Genetic networks that govern sexual reproduction in the Pezizomycotina

Andi M. Wilson,^a P. Markus Wilken,^a Michael J. Wingfield,^a Brenda D. Wingfield^{a#}

^aForestry & Agricultural Biotechnology Institute (FABI), Department of Biochemistry, Genetics & Microbiology, University of Pretoria, Pretoria, Gauteng, South Africa

Running head: Networks governing sexual reproduction in fungi

#Address correspondence to Brenda D. Wingfield, Brenda.wingfield@up.ac.za.

SUMMARY	1
INTRODUCTION	2
MATING IN FILAMENTOUS FUNGI	3
THE <i>MAT</i> GENES	7
The <i>MAT1</i> Locus	7
The Mating-Type Genes	8
The Primary Mating-Type Genes	9
The Secondary Mating-Type Genes	10
The Functions of the Mating-Type Genes	12
FEMALE AND MALE FERTILITY	17
RECOGNITION OF A SUITABLE MATING PARTNER	19
The Mating Pheromones	20
Processing of the a-factor	24
Processing of the α -factor	26
The Cognate Receptors	28
Signal Transduction Pathways	29
PROTOASCOMATAL DEVELOPMENT AND ASCOMATAL MATURATION	32
The STRIPAK Complex	33
The PACC Pathway	35
The COP9 Signalosome	37
ASCOSPORE PRODUCTION & GERMINATION	38
Ascosporegenesis	38

Germination	41
LOOKING TO THE FUTURE	42
CONCLUDING REMARKS	45
ACKNOWLEDGEMENTS	45
REFERENCES	46
AUTHOR BIOGRAPHIES	76

SUMMARY

Sexual development in filamentous fungi is a complex process that relies on the precise control of and interaction between a variety of genetic networks and pathways. The mating-type (*MAT*) genes are the master regulators of this process and typically act as transcription factors, which control the expression of genes involved all stages of the sexual cycle. In many fungi, the sexual cycle typically begins when the mating pheromones of one mating type are recognized by a compatible partner, after which physical interaction and fertilization occurs. Subsequently, highly specialized sexual structures are formed, within which the sexual spores develop after rounds of meiosis and mitosis. These spores are then released and germinate, forming new individuals that initiate new cycles of growth. This review provides an overview of the known genetic networks and pathways that are involved in each major stage of the sexual cycle in filamentous ascomycete fungi.

KEYWORDS

Sexual reproduction

Pezizomycotina

MAT genes

Pheromones

Fertility

INTRODUCTION

Sexual reproduction is an essential component of the life cycle in most Eukaryotes. This is particularly true for metazoans, where sexual reproduction is usually the only process that allows for the production of offspring (1). Fungi are no exception and have maintained complex and active sexual cycles, in addition to being capable of asexual reproduction (2). Combined with the energy intensive nature of sexual development, this suggests that regular cycles of sexual development are important to the long-term survival of fungal species, resulting in benefits beyond simply the production of genetically-unique offspring (3).

Sexual reproduction has been described in many species across the Phylum, Ascomycota (4, 5). Although the members of this group show immense diversity in morphology and ecology, their sexual cycles remain highly conserved. In most species, sexual reproduction is centrally controlled by two sets of major regulators (6); the mating-type (*MAT*) genes and the pheromone response pathway (7). While the sexual cycles of only a subset of species have been thoroughly investigated, their conserved nature, even between distantly related taxa, suggests that these processes are likely to be similar in most fungal lineages. Despite these similarities, there are some differences with regards to the functions of the genes involved and the role of external processes in the sexual cycles of fungi.

Ascomycete fungi reside in three sub-phyla: Saccharomycotina, Taphrinomycotina and Pezizomycotina (8). The first two groups accommodate yeasts and dimorphic species,

which have been extensively studied, particularly with regards to their sexual cycles (9, 10). In contrast, the Pezizomycotina is comprised entirely of filamentous fungi. Representing the largest sub-phylum and encompassing a wide diversity of important fungi, the Pezizomycotina accommodates both detrimental as well as beneficial fungi. Examples include the human pathogenic opportunist *Aspergillus fumigatus* (11) and the dermatophyte *Trichophyton rubrum* (12), as well as plant pathogens such as the notorious chestnut blight fungus, *Cryphonectria parasitica* (13) and the causal agent of pine pitch canker, *Fusarium circinatum* (14). A number of ecologically important species have also been described, including those that are important for wood and litter decay and well as those that form associations with other organisms, including lichens (15, 16).

Importantly, the sub-phylum Pezizomycotina also accommodates a number of important model species, including *Neurospora crassa*. Research on this species has significantly contributed to our understanding of many fundamental genetic processes and molecular mechanisms (as reviewed in 17). The “one gene, one enzyme” hypothesis that gave birth to molecular biology was developed using nutritional mutants of *N. crassa* in 1941. Since then, this species has been used as a model to study the molecular components of circadian rhythms, the Repeat-Induced Point (RIP) mutation genome defence mechanism, RNA interference (RNAi) and epigenetics. Together with other model Pezizomycotina such as *Aspergillus nidulans*, fungi have provided insights into a variety of important Eukaryotic processes including photosensing (18), secondary metabolism (19, 20), cellular synthesis pathways (21), asexual reproduction (22, 23) and sexual

development (24–27). In this review, we consider the genes and genetic networks that govern sexual reproduction in a wide variety of species from the Pezizomycotina, considering research done in both model and non-model fungi.

MATING IN FILAMENTOUS FUNGI

Mating in ascomycete fungi is relatively well-conserved, particularly with regards to the genetic mechanisms that govern sexual development (28). In both filamentous fungi and the yeasts, sexual reproduction is regulated by the mating-type (*MAT*) genes, mating pheromones and various signal transduction pathways. In fact, much of our understanding of sexual reproduction in filamentous fungi is based on similar pathways and processes that were first identified and studied in the yeasts (28). For example, the *MAT* genes and the pheromone genes were first described in *Saccharomyces cerevisiae* (29–31). Later, similar genes were identified in the genomes of many filamentous fungi (32–37) and, in some cases, these genes can fulfill their roles when heterologously expressed in *S. cerevisiae* (38, 39). This alone illustrates the phenomenal level of conservation in the regulatory pathways of sexual development across diverse ascomycete fungi.

Despite the regulatory and genetic conservation of sexual reproduction in the yeasts and the filamentous fungi, there are notable differences, particularly with regards to the sexual tissues that are formed during the sexual cycle. In *S. cerevisiae*, compatible mating partners typically fuse and undergo karyogamy to form a diploid cell. Under conducive environmental conditions, this is followed by meiosis and the production of

sexual progeny cells within a simple ascus as reviewed by Haber (40). In contrast, filamentous fungi form highly complex and multicellular sexual structures, including the ascomata (Fig. 1). Fungi residing in different orders produce ascomata of different shapes and sizes, but all of these structures contain asci in which the sexual ascospores are produced [reviewed by (41)].

While ascomatal structures are morphologically diverse across the Pezizomycotina, the intermediate steps that result in the production of these tissues are relatively well-conserved (Fig. 1). In many fungi, particularly self-sterile fungi, sexual reproduction begins with the fertilization of a female structure (ascogonium) by male structures, in the form of conidia or spermatia (42). Fertilization initiates the production of highly specialized cells and tissues, such as the protoascomata that mature into ascomata which bear asci. And within the asci, ascospores are produced via numerous rounds of meiosis and mitosis. The culmination of sexual development occurs when the ascospores are released into the environment to germinate.

The initiation of sexual reproduction, ie: the step(s) before fertilization, differs from species to species. The requirements for initiation can be used to describe the four major sexual strategies found in the the Pezizomycotina, ie: heterothallism, primary homothallism, secondary homothallism, and unisexuality (Fig. 2). Individuals that exhibit heterothallic behavior are self-sterile, requiring a partner to engage in sexual reproduction (Fig. 1, Step 1). Each of the mating partners carries a unique version of the *MAT* locus, and these must be different for sexual reproduction to occur. Each partner

thus has a unique sexual identity, referred to as mating type, and this is conferred by the type and distribution of the *MAT* genes (Fig. 2).

It is worth noting that while the sexual and mating identities in heterothallic fungi can be described using mating type, these fungi also independently produce cells and structures which can be classified as male or female. Thus, in addition to being of the opposite mating type, compatible partners must also produce complementary male and female structures (43, 44). These two identities are not linked and many heterothallic fungi are hermaphroditic, producing both male cells and female structures. Mating type is specifically determined by the genes present at the *MAT* locus as discussed in detail below. In contrast, male and female structures are produced independently of the *MAT* information present. Importantly, a successful sexual interaction only requires the recognition of one set of male/female cells. Thus, male cells from an individual of the MAT1-1 type must interact with the female structures from an individual of the MAT1-2 type or *vice versa* (43, 44). Mating type and male/female identity [primarily](#) ~~only~~ interact upon the expression of the mating pheromones, a process that is discussed in detail later in this review.

It is somewhat more difficult to distinguish between the different forms of homothallism at the physiological level because in all cases, a single isolate is capable of independent sexual development and is considered self-fertile (Figure 2). These can, however, be differentiated genetically by studying the distribution of the *MAT* genes [Figure 2, (45)]. In primary homothallism, all of the necessary *MAT* genes are present and expressed by

a single individual and within a single nucleus (46). Secondary homothallism describes two different strategies; pseudohomothallism and mating type switching. In the former, the entire *MAT* complement is expressed by a single individual, but the *MAT* information is distributed between two independent nuclei that produce a heterokaryotic mycelium (47). Species capable of mating type switching harbor two versions of the mating information. One of these versions has a copy of all the *MAT* genes, but can be altered to produce the second variant, which has a different gene complement (48, 49). Each of these two variants has a different mating specificity, and it is the expression of the *MAT* genes that drives compatibility between the two, thereby resulting in self-fertility in a single isolate (50). Unisexuality, describes a system where a self-fertile individual retains and expresses only mating information from a single mating type (51).

The complex physiological processes underpinning sexual reproduction are stringently controlled by a huge number of genes. The primary regulators are encoded by the *MAT* genes, which regulate the expression of hundreds of sex-related genes (38, 52, 53), such as those involved in mating partner recognition. Additionally, many non-*MAT* regulated genes, including those involved in male and female fertility, ascomatal maturation and ascosporegenesis, are essential for the completion of the sexual cycle. The following sections provide details of the roles that each of these genes play.

THE *MAT* GENES

The *MAT1* Locus

The mating-type locus contains the *MAT* genes and has a primary role in the sexual process. This locus was formally described as *MAT1* based on the assumption that all ascomycetes would have a single copy of this region (54). Two versions of the *MAT1* locus are present in the diallelic mating system of heterothallic fungi, and this is recognized by naming these *MAT1-1* and *MAT1-2*, with the use of capital letters acknowledging the lack of dominance between the two “alleles” [Fig. 1, (54)]. The non-allelic nature of these versions was formally acknowledged in 1990, when the term idiomorph was suggested by Professor John Wyatt and introduced by Metzberg and Glass (55) to describe these two versions of the *MAT1* locus. Although the two different idiomorphs can easily be recognized in most heterothallic species, the term does not extend to their homothallic counterparts. Most homothallic species have a single *MAT1* locus that is defined based on the presence of homologous mating-type genes (56), although in some cases more than one, often unlinked, *MAT* locus can be present within a single genome (32, 57, 58).

The Mating-Type Genes

The precise definition of what constitutes a *MAT* gene is centered around their position in one of the two mating-type idiomorphs (56). This definition is broadly accepted, but an alternative description holds that a true mating-type gene is involved in establishing the mating type identity of the nuclei after fertilization, and that its protein product would

localize only to the nucleus (59). Although such a definition may be considered the golden standard, it is impractical because many mating-type genes have not been the subject of functional studies. There is also convincing experimental and computational evidence that at least one of the known mating-type proteins, MAT1-1-2, localizes only to the cytoplasm and not the nucleus (60, 61). Furthermore, some homologous *MAT* genes have different functions in different species, precluding the application of function as an indication of mating identity. In the absence of a more suitable option, a system that defines mating-type genes as those present at the mating-type locus remains the most intuitive.

The locus-based definition for mating-type genes has been subject to valid criticism. Named *MAT* genes are widely considered to be the master regulators of sexual reproduction, responsible for conferring sexual identity and orchestrating processes linked to the sexual cycle. These roles can be validated only through functional studies (56, 62), which are lacking for most *MAT* genes. This implies that newly described *MAT* genes are mating-type genes in name only. This disparity is now being addressed, with new functional studies of diverse *MAT* genes being published regularly (63–65). Another criticism of a locus-based *MAT* gene definition is that the dynamic nature of this region can result in the inclusion of genes with no known role in mating into the *MAT1* locus (66, 67). This contradicts the positional-based definition of a mating-type gene, a shortcoming highlighted in a recent revision of the *MAT* gene nomenclature system (56). Assigning a novel, *MAT*-specific name to a gene within the *MAT1* locus, only when no known homolog exists, can resolve this issue (56). This approach has been

successfully applied to many novel *MAT* genes (68–71). In cases where homologs of known genes have been observed, these are significantly diverged to the point of representing *MAT*-specific versions of these genes (67, 72). Future studies must consider whether these genes have gained novel, *MAT*-specific functions.

The Primary Mating-Type Genes

Two *MAT* genes are almost universally present in all studied Pezizomycotina Classes (Table 1). *MAT1-1-1* was named as the first gene (*MAT1-1-1*) from the *MAT1-1* (***MAT1-1-1***) idiomorph (73, 74), and its presence defines that idiomorph. Similarly, the *MAT1-2-1* gene typifies the *MAT1-2* idiomorph as the first described gene from this idiomorph (73, 75). The protein products of both these genes have conserved domains involved in DNA binding (76). Additionally, a conserved intron is present in the *MAT1-2-1* HMG-box domain positioned across a serine codon, a characteristic that is unique to mating-type genes that have this domain (56, 77, 78). The core roles of *MAT1-1-1* and *MAT1-2-1* in the sexual process, as well as their wide distribution across the Pezizomycotina, has led to these genes being known as the core, or primary, mating-type genes (56, 62).

The Secondary Mating-Type Genes

Apart from *MAT1-1-1* and *MAT1-2-1*, 11 additional *MAT1-1*-specific and 14 additional *MAT1-2*-specific genes have been described in the Pezizomycotina (56, 68, 70, 79, 80), and are all considered secondary *MAT* genes (Table 1). Although a formal definition for secondary mating-type genes does not exist, these genes are present within the confines of the mating-type locus, have no known homologs outside of the mating-type

genes, and they tend to show a lineage-specific distribution (56). *MAT1-1-2* and *MAT1-1-3* have been identified from species in the Sordariomycetes and Leotiomycetes (56), while homologs of *MAT1-1-7* have been reported among the Lecanoromycetes and Eurotiomycetes (81), making these three genes the most widely spread secondary *MAT* genes. In a small number of Sordariomycetes, the *MAT1-1-2* gene is unusually present in the *MAT1-2* idiomorph (66, 82, 83), although this does not impact the sexual cycle in these fungi. Most other *MAT* genes have been identified in only a single Order, Family or Genus (56).

Since the last overview and revision of all known mating-type genes (56), four secondary *MAT1-1* genes have either had names assigned or amended. A homolog of the known *MAT1-1-7* gene was recently identified in several families of the Lecanoromycetes (81). Through positional conservation and phylogenetic analysis, it was shown that this gene is also homologous to the *MAT1-1-9* gene described from members of the Eurotiomycetes (81), leaving the *MAT1-1-9* gene name vacant (Table 1). *MAT1-1-10* and *MAT1-1-11* have been described from the *MAT* loci of *Morchella* species [Pezizomycetes, (68)]. Another putative *MAT1-1-10* gene was described in the Dothideomycete genus *Teratosphaeria* (69, 84) and appears also to be present in the *MAT1* locus of other members of the Mycosphaerellaceae (69). There is no apparent homology between the two *MAT1-1-10* genes, and they follow a lineage specific distribution.

Four novel genes have been identified from the *MAT1-2* mating-type idiomorph. Two of these have been named *MAT1-2-12*; one in *Teratosphaeria* species (69, 84) and the other in *Calonectria* species (70). There is no homology between the two *MAT1-2-12* genes or their encoded proteins (B. Wingfield, unpublished). Following the guidelines of Wilken *et al.* (56), the *Teratosphaeria* *MAT1-1-10* (69, 84) should be renamed *MAT1-1-12*, while the *Calonectria* *MAT1-2-12* (70) should be known as *MAT1-2-13*. A novel *MAT* gene was also described in species of the *Lecanoromycetes* and has been named *MAT1-2-14* (81), while another gene identified in *Colletotrichum* species was added as *MAT1-2-15* (80). None of these genes have any conserved domains that might suggest their possible functions.

The complexity in naming novel mating-type genes together with the need to constantly reevaluate already established *MAT* gene names, highlights the need for a dynamic, curated database of mating-type gene names and homologs. A web-based *MAT* gene server to assist in mating-type gene identification is currently being developed (Wingfield *et al.*, in preparation) and will go some way to address these challenges.

The Functions of the Mating-Type Genes

Functional analyses remain the only true test for the role of a gene in the biology of a fungus, but such studies are lacking for most of the *MAT* genes. It is not surprising that the majority of available functional studies have targeted the core *MAT1-1-1* and *MAT1-2-1* genes, largely due to their wide distribution and central role in the sexual process (25, 52, 53, 60, 85–87). More recently, some studies have emerged that target the

secondary *MAT* genes, illustrating the complex interactions that underlie sexual reproduction (63, 64, 85, 86, 88).

The history of functional studies on *MAT* genes has closely followed technological advances. Early studies relied on the transformation of cloned DNA into a complementary strain (33, 89). Later, mutations were induced in target genes by employing techniques such as the RIP mechanism that silences duplicate sequence products (25). The commercialization of molecular techniques relying on PCR amplification has strongly advanced new technologies used in functional studies. These include transcription profiling during sexual development (53, 87), comparisons of whole expression profiles between wild type and mutant strains (53), antisense RNA approaches to disrupt gene function (87) and microarray experiments (52). Genome-based technologies have further expanded the available toolbox, with techniques such as CHIP-Seq analysis (90) and transcriptomics (91) becoming more prominent. The advent of novel, easy-to-use gene editing technologies including the CRISPR-Cas9 system promises further progress, and has already led to the functional analysis of some mating-type genes (60, 64).

Functional studies on the *MAT1-1-1* and *MAT1-2-1* genes and their core domains are too numerous to cover in detail here. They have, however, provided an impression of conserved functions intermingled with great adaptability. Generally, both primary mating-type genes rarely impact vegetative growth directly (25, 52, 87, 88, 92), although they are necessary for successful completion of the sexual cycle in both homothallic

and heterothallic fungi (25, 38, 52, 60, 75, 87, 92, 93). *MAT* genes control the sexual process by conferring sexual identity to individual isolates (34, 75), mediating the formation of male and female reproductive tissues (88, 93, 94), regulating the mating pheromone and receptor system (35, 52, 53, 88), and by controlling ascomatal and ascospore development (34, 38, 88, 95). Although these functions appear to be well conserved, even among distantly related species, it is worth noting that these effects may not be universal. For example, in the smut fungus *Villosiclava virens* that causes false smut in rice, deletion of the *MAT1-1-1* gene impacts significantly on vegetative growth rate and conidial structure (60). Whether this species is an exception or provides evidence of as yet undiscovered complexity in the functions of the mating genes, remains to be discovered.

Among the secondary *MAT* genes, the *MAT1-1-2* and *MAT1-1-3* genes have been the subject of most of the functional studies. Such studies have shown that these genes are generally not essential for sexual reproduction, but rather function to fine-tune the sexual process. Early work on the model species *N. crassa* showed that both *MAT1-1-2* and *MAT1-1-3* influence the expression of sex-related genes, with mutants showing reduced fertility (25, 43). In contrast, in *Fusarium graminearum*, both genes are important for male and female fertility, as well as ascospore formation (96). Once again, there are exceptions to this generalization, such as the *Sordaria macrospora* *MAT1-1-2* gene, which is essential for completion of the sexual cycle (85). In some cases, this gene is also involved in functions attributed to the core *MAT* genes, including nuclear recognition, control of the pheromone response pathway, and ascomatal development

(43, 59, 85). Deletion of the *S. macrospora* *MAT1-1-3* gene shows no observable phenotype (85), although its deletion of the homologous gene in some homothallic *Fusarium* species abolishes self-fertility (97) and leads to significantly reduced fertility in *Podospora anserina* (43).

An interesting finding that has emerged from studies on *MAT1-1-1*, *MAT1-1-2*, *MAT1-1-3* and *MAT1-2-1* genes is that many non-mating functions are also under the control of these genes. These include general metabolic processes and energy production (52, 53), stress responses (60), the production of secondary metabolites (90), virulence (60), vegetative growth (64) and other diverse biological processes. This should not be surprising as all of these genes encode proteins that are characterized by conserved DNA binding domains that are associated with transcription factors (56, 98). Therefore, it was plausible that these proteins would be involved in regulating a complex network of gene expression to accomplish the sexual process (99). Additionally, sexual reproduction is an involved process that relies on specialized tissue differentiation in response to environmental and genetic cues (62). Unfortunately, many functional studies only evaluate the effect of *MAT* gene disruption on the sexual process, and do not screen for pleiotropic effects. As new functional studies on the *MAT* genes are conducted, the effect of the *MAT* genes on non-mating functions will likely become more clear.

Apart from *MAT1-1-2* and *MAT1-1-3*, only five additional secondary *MAT* genes have been functionally characterized. Deletion of the *MAT1-1-5* gene in *Sclerotinia*

sclerotiorum reduces the production of spermatia and completely blocks ascomatal development, halting the sexual process (88). The *MAT1-2-10* gene in the same species is involved in regulation of the cytoskeleton, with deletion mutants showing delayed ascomatal development, and when mature, produce aberrant numbers of ascospores (88). In the closely related species *Botrytis cinerea*, the roles of *MAT1-1-5* and *MAT1-2-10* are different, with both proteins jointly regulating the transition to karyogamy (92). The *MAT1-2-9* gene in *F. graminearum* appears not to be essential for sexual reproduction, but its transcription patterns do mirror those of the primary *MAT* genes (97). The *MAT1-2-4* gene in *A. fumigatus* is another essential regulator of sexual reproduction, but deletion of this gene did not allow for the elucidation of a specific function in sexual reproduction (63). A gene knock-out study of the *Huntia omanensis* *MAT1-2-7* gene showed that this gene is essential for maturation of the ascomata, a function shared with other secondary mating-type genes (64).

The mating-type genes are themselves subject to genetic regulation, mostly as a response to environmental cues. Studies on the model species *P. anserina* have revealed a network of HMG-box motif genes that regulate sexual development, some through direct transcriptional control of the mating-type genes and their downstream targets (99). Two genes, *ste11* and *hmg8* were identified as directly controlling *MAT* gene expression. A third gene, *pro1*, connects environmental cues, such as nutrient starvation, directly to *MAT* gene expression by activating *hmg8* (100). In the heterothallic species *Trichoderma reesei*, the effect of light on the sexual process is mediated by the *env1* and *vel1* genes through direct transcriptional regulation of *MAT1-*

2-1 (101, 102). Many other environmental response genes have been linked either directly or indirectly to *MAT* gene expression, and these have been extensively reviewed by Wilson *et al* (62). Although these studies have provided some clarity, a thorough understanding of the genes that control *MAT* gene expression is still needed to fully appreciate the complexities of sexual reproduction in the Pezizomycotina.

Despite great progress having been made in recent times to identify and characterize mating-type genes, substantial gaps remain in our knowledge of this subject. For example, *MAT* genes have not been formally described in many Classes of fungi, and even in the Classes with known mating type-genes, some Orders lack representation (Fig. 3). It has been shown that variations in mating-type strategy can be present among even members of the same genus (58, 66), and this is often linked to interesting variations in the mating type genes or *MAT1* locus structure (71, 82, 83). Examples exist of gene conversions [e.g. *MAT1-1-1* to *MAT1-2-7*, (103)] and locus expansions (66, 67) that have helped to shape the *MAT1* region, but much remains unknown regarding the evolution and maintenance of this important region. Future studies on *MAT* gene function as well as the description of the *MAT1* loci in underrepresented groups (Fig. 3) will contribute to addressing current knowledge gaps.

FEMALE AND MALE FERTILITY

Regardless of their mating type, most heterothallic fungi are capable of producing both male and female structures. Conidia or other spermitizing cells act as the male cells, which act by fertilizing the female structures known as ascogonia (Fig. 1, (104)). Given

that these structures are produced independently of one another, there are genes whose functions are exclusively linked to either male or female fertility. As a direct consequence, certain isolates can be female fertile and male sterile, or *vice versa* (105). Isolates that are considered male sterile are typically unable to produce spermitizing agents or are incapable of fertilizing the ascogonia of the female partner. In contrast, isolates considered female sterile are either unable to form ascogonia or protoascomata or produce protoascomata that are unable to mature.

Some of the most important genes that influence male and female fertility are the mating pheromones and their cognate receptors, respectively (37, 106, 107). In general, the pheromones are expressed by the male cells and are recognized by their receptors, which are present on the cells of the female ascogonia (107). Understandably, deletion of the pheromone factors or receptors results in either a decrease or complete loss of fertility and the entire sexual pathway. Given the complexity and importance of this pathway in sexual development, a later section of this review is devoted entirely to the pheromone response pathway.

Very few genes have been associated with male fertility in the filamentous fungi. Instead, many of the genes that have been shown to contribute to the early steps of the sexual cycle predominantly regulate female fertility (108–111). Thus, apart from certain *MAT* genes (96) and the pheromone genes (109, 112), only the *mcm1* gene in *Magnaporthe oryzae* has thus far been described as being essential to male fertility (113). Encoding a MADS box transcription factor, this gene is also required for the

production of microconidia, which likely act as the spermitizing agents in this fungus. It is thus not surprising that $\Delta mcm1$ mutants are male sterile (113).

In *P. anserina*, four genes have been shown to contribute significantly to female fertility; *nox1* is important for the production of maternal tissues (114), *mpk1* is important for the development of the underlying mycelium that surrounds and nourishes the growing ascomata (115), and the two IDC genes, *idc2* and *idc3*, are involved in the communication between these two tissue types (116). It is thought that *nox1*, which encodes an NADPH oxidase, is responsible for providing the developing protoascomata with fresh nutrients via ROS-mediated signaling, and its deletion results in very small protoascomata and delayed development (114). Similarly, deletion of either of the IDC genes results in isolates that can differentiate into both male and female structures, but exhibit delayed protoascomatal maturation after fertilization. In these isolates, some ascomata will develop, but only a few ascospores are released (116).

In *N. crassa*, three genes coding for proteins involved in the calcium ion signaling pathway are important for female fertility (117); a phospholipase (*plc1*), a secretory phospholipase (*sp/A2*) and a calcium-hydrogen ion exchanger (*cpe1*). Two double mutants, $\Delta plc1/\Delta sp/A2$ and $\Delta plc1/\Delta cpe1$, are entirely female sterile and produce no ascomata. It is thought that these phenotypes are the result of a significant decrease in the expression of the two mating pheromones as well as a female/male fertility gene (*fmf-1*) in the double mutants (117). A fourth gene, *fig1*, is involved in a calcium ion uptake system and is essential for female fertility in MAT1-2 isolates of *N. crassa* (118),

illustrating not only that calcium uptake and signaling are important for fertility, but that the role of a protein can change based on the mating type of the isolate.

RECOGNITION OF A SUITABLE MATING PARTNER

The sexual cycle, particularly in heterothallic species, begins with the identification and recognition of an appropriate mating partner (28). This process is facilitated in many species by the pheromone response pathway (7). Partners of opposite mating type express one of the two mating pheromones, which are secreted into the environment by male cells, and are subsequently recognized by a suitable opposite mating type partner via pheromone receptors (Fig. 1). Upon recognition of these diffusible peptides, specialized hyphae known as trichogynes grow from the female partner towards the male cell. Once the male and female cells interact physically, fertilization takes place, and the sexual cycle begins. At a molecular level, the pheromone factor/receptor recognition activates a heterotrimeric G-protein, thereby initiating the signal transduction pathway, which culminates in the transcription of genes related to mating (Fig. 4).

The Mating Pheromones

The two most commonly utilized fungal mating pheromones are the α - and **a**-factors (7, 119). The genes encoding these small proteins are transcriptionally regulated by the MAT1-1-1 and MAT1-2-1 proteins, respectively. Consequently, the pheromone genes are expressed in a mating type-dependant manner in most heterothallic species, allowing for recognition between individuals of opposite mating type (35, 91, 120, 121).

Genes encoding the two pheromones have been described in species with different sexual strategies. These genes have been found in diverse heterothallic species such as *N. crassa* (35), *H. omanensis* (91), *C. parasitica* (120), and *Magnaporthe grisea* (121). Similarly, the homothallic species *F. graminearum* (122), *A. nidulans* (36), and *S. macrospora* (123, 124) all harbor genes encoding mating pheromones. Even species exhibiting the more unique strategies such as unisexuality [*Huntia moniliformis* (69) and *Neurospora africana* (125)] and pseudohomothallism [*P. anserina* (37)] express mating pheromones.

Interestingly, there are groups of fungi where only a single mating pheromone has been identified. This is despite a wealth of genomic and transcriptomic data and a significant effort to identify these genes. In *A. nidulans* (36) and other Eurotiomycete fungi (125) for example, the gene encoding a protein with similarity to the **a**-factor pheromone has remained elusive, while the **α**-factor pheromone has been identified. Similarly, species in the Pezizomycetes, including *Tuber melanosporum* (126), also harbor only the **α**-factor encoding gene. Whether a second factor exists, with or without similarity to the **a**-factor, remains to be discovered. These genes, being highly variable and typically very short, are notoriously difficult to identify. Their identification consequently requires gene sequences from very close relatives to be used as BLAST queries against the genomes of newly investigated species. Alternatively, when such sequences are not available, it is possible to make use of the microsynteny that exists between more distantly related species (125). In this way, pheromone-flanking genes in species where the pheromones are known can be used to identify the locus harboring the pheromone, after which the

pheromone genes can be manually annotated. Additionally, the conserved domains of the pheromones, as detailed below, can also be used to search through and filter protein-coding genes in these genomes. This makes it possible to identify proteins that share structural similarities to the mating pheromones.

Given their important role in partner recognition, it is not surprising that the pheromone genes are essential for sexual reproduction in many model heterothallic species (Table 2). In *N. crassa* (106) and *C. parasitica* (112, 127), both pheromone types are indispensable for male fertility, with deletion mutants unable to complete the sexual cycle. In *N. crassa*, the importance of each pheromone in male fertility is mating type dependent (106). Thus, **a**-factor deletion inhibits sexual development in a MAT1-2 strain, while α -factor deletion precludes sexual development in a MAT1-1 strain. Additionally, the **a**-factor pheromone also plays an important role in female fertility in *N. crassa* (109), a role that is not restricted to either mating type. Similarly, the α -factor is important for male fertility (112) in *C. parasitica*, while the **a**-factor pheromone is essential for female fertility (120).

The mating pheromones play an important role in the sexual development of the homothallic species, *F. graminearum* and *S. macrospora* [Table 2, (122, 128)]. While not essential for this pathway, gene deletion mutants in both species display decreased fertility, producing significantly fewer mature ascomata than wild type isolates. This phenotype is observed in *F. graminearum* when the α -factor gene or both the α - and **a**-factor genes are deleted (122). In contrast, only the deletion of both pheromone genes

induces this phenotype in *S. macrospora* (128). In both species, deletion of only the **a**-factor pheromone results in fully self-fertile isolates (122, 128), suggesting that this gene is not important in homothallic mating.

The strict association of the α -factor with the MAT1-1-1 protein and the **a**-factor with the MAT1-2-1 protein has some interesting implications and may play an important role in establishing the sexual strategies observed in various fungal species. This association is typically strongest in heterothallic species, where the pheromones are used to communicate the mating type of an individual to potential mating partners, thereby facilitating partner recognition (35, 91, 120, 121). This association is less stringent in homothallic species, with both the primary and secondary MAT proteins regulating pheromone expression. In *S. macrospora*, MAT1-1-1 positively regulates the expression of both pheromones, MAT1-1-2 negatively regulates **a**-factor expression while MAT1-2-1 positively regulates **a**-factor expression (52, 85). Similarly, in the homothallic *F. graminearum*, the MAT1-1-1, MAT1-1-2, MAT1-1-3 and MAT1-2-1 proteins all negatively regulate the expression of both pheromone factors to varying degrees (96). The lack of a requirement to attract and recognize partners of opposite mating type has evidently changed the MAT-dependent regulation of the pheromones, even while these mating factors remain important for sexual development in these species.

Pheromone expression and regulation is particularly noteworthy in species with the more unique mating strategies. Both pheromones are expressed in the two unisexual species, *H. moniliformis* and *N. africana*, despite the absence of MAT1-1-associated

genes in *H. moniliformis* and *MAT1-2*-associated genes in *N. africana* (91, 125). Similarly, species of the genus *Colletotrichum* are known to exhibit various mating strategies, including mating type switching (129) and unbalanced heterothallism, a type of self-sterility in which suitable mating partners have complementary mutations in genes involved in sexual reproduction (130). This is despite the fact that these species harbor only genes associated with the *MAT1-2* idiomorph (80). These species have also been shown to have undergone concerted gene loss events associated with both pheromone factors, with some species harboring only one of the two pheromones and some species harboring neither (80). This suggests that the pheromones in these species may be regulated in a *MAT*-independent manner that may, at least partly, explain their divergence from the typical mating strategies.

Fungal mating pheromones are produced as larger pre-proteins that are subsequently processed into smaller, mature pheromone factors that are functionally active [Fig. 5, (7)]. Much of our understanding regarding the processing of mating pheromones in fungi arises from research conducted on *S. cerevisiae* (7). However, many of the processing proteins found in this model yeast have homologs in filamentous fungi and are thought to play similar roles in pheromone processing.

Processing of the α -factor.

The defining feature of the α -factor pre-protein is a terminal CaaX domain, a region comprised of a cysteine residue followed by two aliphatic residues and an amino acid of any type [Fig. 5A, (131)]. This highly conserved domain is post-translationally modified

561 by a variety of other proteins before the mature pheromone factor is secreted [reviewed in (7)]. The first post-translational modification occurs when RAM1 and RAM2 proteins farnesylate the cysteine residue of the CaaX domain. The three C-terminal residues are then removed by RCE1 or STE24, where after STE14 adds a carboxymethyl group to the now-terminal C residue. The final step is the proteolysis of the N-terminal region by STE24 and AXL1. This produces a mature factor which is subsequently secreted by the STE6 transport protein.

The **a**-factor pheromone processing pathway has not been elucidated in any filamentous fungi, but many of these fungi contain genes encoding homologs of the *S. cerevisiae* processing proteins. Almost all of these genes have been identified in the genomes of species as diverse as *A. fumigatus* (132), *A. cristatus* (133), *A. niger* (134), *M. oryzae* (135), *N. crassa* (109), *Penicillium marneffeii* (141), *P. anserina* (53), *Verticillium dahliae* (136). An interesting exception is *axl1*, which has only been described in *N. crassa* (109). The fact that most of these processing proteins as well as the **a**-factor pheromone have all been maintained in the Pezizomycotina strongly implies that there may be some conservation in the entire processing pathway.

Unfortunately, very little research has been conducted on the function of the pheromone processing genes in the Pezizomycotina. Consequently, it remains unclear whether they play the same roles in pheromone processing as they do in *S. cerevisiae*. Experimental evidence does suggest that the two RAM proteins play a far more general role in the biology of these fungi than simply being involved in pheromone processing. The

deletion of *ram1* and *ram2* is lethal in *P. anserina* (53) and *M. oryzae* (135), respectively; while *ram1* disruption in *M. oryzae* affects processes as diverse as growth, stress resistance, and virulence (135). This is perhaps not surprising because the RAM proteins are post-translational modifiers that add farnesyl groups to proteins in order to ensure their appropriate cellular localization (137). This form of modification affects a large number of proteins, particularly those involved in signal transduction, and consequently affects a variety of biological processes not limited to mating (135).

Manipulation of the *A. cristatus* *MAT1-1-1* and *MAT1-2-1* genes provides further experimental evidence to support the importance of the α -factor pheromone processing proteins in the sexual development of filamentous fungi. The *rce1* and *ste14* genes, both involved in some of the final processing steps, are significantly down regulated in *MAT1-1-1* and *MAT1-2-1* mutants, suggesting that these genes are (in)directly under the control of these two master regulators (133). Similarly, under conditions of high osmolarity, where sexual reproduction is precluded, *rce1* is down regulated in *A. cristatus* (138). While such results do not confirm a function for these proteins in pheromone processing, they certainly imply a role for these genes during sexual development.

Processing of the α -factor

The α -factor pre-protein harbors a hydrophobic N-terminal secretion signal, a pro-region and repeating units of the mature pheromone sequence [Fig. 5B, reviewed in (7)]. The first step in processing is the removal of the secretion signal by a signal peptidase in the

endoplasmic reticulum. This is followed by proteolytic cleavage and removal of the pro-region by the KEX2 protein. Once the N-terminus has been removed, the pre-protein is left containing repeats of a mature α -factor. Release of the mature pheromone peptides relies on the presence of two dipeptide motifs that flank these repeats. The first, either the dipeptide KR or RR, is a KEX2-like peptidase recognition site. KEX2 activity releases the numerous immature pheromone peptides, which are then subjected to terminal amino acid removal by KEX1. The final step in the processing of this pheromone is due to STE13-like aminopeptidase activity, which recognizes the second dipeptide motif, either XA or XP. Finally, STE13 activity releases the mature pheromone, which is subsequently packaged into vesicles and secreted via the classical secretion pathway.

Genes encoding the KEX1, KEX2, and STE13 processing proteins have been found in the genomes of various filamentous fungi, including *S. macrospora* (39, 139), various *Trichoderma* species (140), *A. fumigatus* (132), *A. nidulans* (32), *N. crassa* (35), *P. marneffeii* (141), *H. omanensis* (91) and *H. moniliformis* (91). Similarly, the processing sites for these proteins are present in the α -factor pheromone pre-proteins of these species, suggesting conservation in the processing of these proteins across yeast and filamentous fungi. This has been experimentally proven in *S. macrospora*, where the α -factor pre-protein in this species can be properly processed, secreted, and is functional in *S. cerevisiae* MAT α cells (39).

Gene deletion and silencing studies in *N. crassa* and *C. parasitica* have further supported the role of these processing proteins in the sexual development of filamentous fungi. Fertility in *N. crassa* is influenced by KEX1, where deletion of the *kex1* homolog, *scp*, results in greatly reduced male fertility (35, 142). Unsurprisingly, this effect is only apparent in the α -factor-expressing MAT1-1 cells, while MAT1-2 cells are unaffected by the gene's deletion. Similarly, *kex2*-silencing in *C. parasitica* also results in reduced male fertility and an inability to produce ascospore-bearing ascomata (143). Lastly, a genome-wide gene expression study in *N. crassa* also showed that the expression patterns of *ste13* closely mimic those of the α -factor pheromone, both increasing during ascomatal development (144). This hints at a role for this protein in the sexual cycle.

The Cognate Receptors

Each of the mating pheromones is recognized by a specific cognate receptor. Both receptors are seven-transmembrane domain proteins, which are comprised of an extracellular recognition domain, seven membrane spanning domains and an intracellular domain that is linked to a G-protein [Fig. 4, reviewed in (145, 146)]. The α -factor pheromone is recognized by the PRE1 protein, which has similarity to the STE3 protein from *S. cerevisiae* (107). Similarly, the PRE2 protein, which is similar to the *S. cerevisiae* STE2 protein, is the α -factor's cognate receptor (38).

As with the pheromone factors themselves, the pheromone receptor genes have been identified and functionally characterized in many filamentous ascomycete fungi (24, 53,

107, 122, 128). While these genes are found in both heterothallic and homothallic fungi, their relative importance in sexual development differs from species to species (Table 2). In some species, both of the receptor genes are essential for sexual development, while in others, these genes are dispensable although their deletion may result in decreased fertility (53, 128, 147, 148).

An interesting characteristic of the two pheromone receptor proteins is that they often have mating type-dependent roles in heterothallic species. In these species, the PRE1 protein plays an important role in female fertility in MAT1-1 isolates, while female fertility in the MAT1-2 isolates depends on the PRE2 protein. This is understandable given that MAT1-1 isolates express the α -factor and need to recognize the a -factor via the expression of *pre1*, while the converse is true of the MAT1-2 isolates. This mating type-dependent effect has been observed in *P. anserina* (53), *N. crassa* (24, 107) and *T. reesei* (147), where $\Delta pre1$ MAT1-1 and $\Delta pre2$ MAT1-2 isolates are female sterile-producing protoascomata that are incapable of being fertilized and thus do not mature into ascospore-bearing ascomata. Importantly, $\Delta pre1$ MAT1-2 and $\Delta pre2$ MAT1-1 isolates interact normally with wild type partners and produce fully mature ascomata and viable ascospores.

The pheromone receptors are important, but typically not essential, for the sexual cycle in a number of homothallic species (Table 2). In both *S. macrospora* (128) and *A. nidulans* (148), the deletion of both *pre1* and *pre2* results in the complete loss of the sexual cycle by precluding the production of ascomata. However, deletion of a single

receptor produces no detectable effect in *S. macrospora* (128) but does result in reduced fertility in *A. nidulans*, with mutants producing smaller ascomata and fewer ascospores (148). In *F. graminearum*, *pre1* is dispensable for sexual development, with $\Delta pre1$ isolates sporulating normally (122). In contrast, single $\Delta pre2$ mutants and double $\Delta pre1/\Delta pre2$ mutants exhibit sub-normal fertility levels (122).

Signal Transduction Pathways

As G-protein coupled receptors (GPCR), the pheromone receptors are linked to a G-protein via their terminal intracellular domains [Fig. 4, (149)]. Prior to pheromone recognition, the G-protein exists as a trimer of three protein subunits- $G\alpha$, $G\beta$ and $G\gamma$. Once the pheromone has physically bound to the receptor, the G-protein is activated, resulting in dissociation of the subunits into a $G\alpha$ monomer and a $G\beta$ - $G\gamma$ dimer, each of which is capable of initiating a downstream signal transduction cascade (149).

Given their close association with the pheromone receptors, it is not surprising that all three subunits play an important role in female fertility (Table 2). Two genes, *gna1* (150) and *gna2* (151), encode $G\alpha$ -subunits that have been functionally characterized with respect to their roles in mating in the heterothallic *N. crassa*. Deletion of *gna1* results in mutant isolates that produce unusually shaped protoascomata (150) and are incapable of growth towards the spermatia (84). There is consequently little or no maturation of these structures into mature ascomata. Interestingly, *gna2* deletion mutants do not produce particularly notable sexual defects, but $\Delta gna1/\Delta gna2$ mutants are more severely affected than single *gna1* deletion mutants (151). This fact illustrates that while

the two α -subunits may have overlapping functions, the GNA1 protein can compensate for GNA2, but not *vice versa*. Deletion of the G α -subunit gene in *S. macrospora*, results in a similar phenotype, although, while protoascomatal production is delayed and reduced in $\Delta gsa1$ mutants, mature ascomata are produced after sufficient incubation (152). This could reflect a change in the importance of the pheromone response pathway in the homothallic *S. macrospora* compared to the heterothallic *N. crassa* (Table 2).

The *N. crassa* β - and γ -subunits, encoded by *gnb1* and *gng1*, respectively, are essential for the maturation of normal ascomata. Single and double mutants of the *gnb1* and *gng1* genes are all female sterile, producing immature protoascomata that do not develop further, even when adequately fertilized (111, 153). On the rare occasion where mature ascomata are produced, they remain small and are unable to eject the ascospores (111). It is perhaps not surprising that deletion of the G α -, G β - and G γ -subunits produce similar phenotypes, and in fact, the expression of the three subunit genes are controlled by one another. For example, *gnb1* deletion results in the complete down-regulation of *gna1* and *gng1* expression (111, 153); while *gng1* deletion results in significant down regulation of *gnb1* (153).

After dissociation, the G α and G β -G γ subunits often act by activating MAP kinase signal transduction pathways that are capable of eliciting a variety of effects- including the initiation of sexual reproduction via the activation or repression of relevant transcription factors [Fig. 4, reviewed in (154)]. In general, the first component of this kind of

transduction pathway is a MAP kinase kinase kinase (MAPKKK), followed by a MAPKK, a terminal MAPK and finally, the targeting protein, which is often a transcription factor. In *N. crassa*, three of these pathways are involved in sexual development and are named after their terminal kinases, ie. MAK1 (110), MAK2 (155) and OS2 (156, 157) (Fig. 4). Intriguingly, while all three pathways have been implicated in protoascomatal development, only the MAK2 pathway is directly linked to the mating pheromone receptors [Fig. 4, (158)]. This suggests that the cross talk between these three pathways is essential for sexual development in *N. crassa*. Thus the deletion of even a single gene in one of these pathways results in female sterility [Table 2, (110)]. The MAK1 (159) and MAK2 (160) pathways are also important for sexual reproduction in *S. macrospora*, where single gene deletions also preclude sexual development (Table 2).

There are some important differences between heterothallic and homothallic fungi relating to the pheromone response pathway and various MAPK cascades. To illustrate these differences, the mutant phenotypes of genes involved in the heterothallic *N. crassa* and homothallic *S. macrospora* are shown in Table 2. What is evident is that the proteins that play a role during the initiation of the pheromone response pathway, ie. the pheromone factors, their receptors and the coupled G-proteins, are less important for sexual development in *S. macrospora* than they are in *N. crassa*. In fact, single gene deletions in *S. macrospora* either result in no sexual defects or simply delay sexual development. In contrast, similar deletions in *N. crassa* drastically affect male and/or female fertility. When the MAPK cascades are considered, however, the effects of single gene deletions in the heterothallic and homothallic species are comparable. This

suggests that while the process of initiating sexual recognition is different in heterothallic and homothallic fungi, the downstream processes are the same.

PROTOASCOMATAL DEVELOPMENT AND ASCOMATAL MATURATION

After partner recognition occurs, male cells are capable of fertilizing the female ascogonia, which will begin to develop into a protoascoma [reviewed in (41)]. It is within this immature sexual structure that the various stages of nucleus migration, karyogamy, mitosis and meiosis will commence. During this time, the protoascoma develops into a mature ascoma containing asci and the sexual ascospores. Three protein complexes and/or pathways (the STRIPAK complex, the PACC pathway and the COP9 signalosome) have been specifically linked to ascomatal development in the filamentous fungi (Fig. 6).

The STRIPAK Complex

One of the most important protein complexes that controls the development of the ascoma is the striatin-interacting phosphatases and kinases (STRIPAK) complex [Fig. 6, (161, 162)]. In fungi, this complex is made up of a protein phosphatase (PP2AA), a striatin (PRO11), a striatin-interacting protein (PRO22), a phocein (MOB3), kinases (KIN3, KIN24), a membrane anchoring protein (PRO45) and many other proteins (162). This complex is highly conserved across the Eukaryotes and is thought to be involved in various signal transduction pathways important for the differentiation of multicellular structures (161).

The first fungal STRIPAK subunits were identified in *N. crassa* (163) and some research has been conducted to elucidate the roles that these proteins play in this species (163–165). However, this complex, its functions, and protein-protein interactions are much better understood in *S. macrospora* [reviewed in (162)]. The *S. macrospora* STRIPAK complex appears to be involved in diverse biological processes due to its primary function being the (de)phosphorylation of a variety of target proteins (162). Many of these targets are known to play important roles in sexual development (166). For example, the α -factor pheromone processing proteins KEX1 and KEX2 are both (de)phosphorylation targets of the STRIPAK complex in *S. macrospora*.

The importance of the STRIPAK complex and its subunits in sexual development has been experimentally shown in *S. macrospora*. Deletion of the genes encoding STRIPAK proteins commonly results in similar phenotypes- particularly during the development of the immature ascomata. In $\Delta pro11$ (167), $\Delta pro22$ (161), $\Delta PP2Ac1$ (168), $\Delta kin24$ (169), $\Delta pro45$ (170) mutants, protoascomata are produced normally, but do not become pigmented over time and do not develop into mature sexual structures. In strains where *mob3* (171) or *pro11* (161) are deleted, an even more extreme phenotype is apparent, with the complete absence of any sexual tissues. Furthermore, the deletion of *PP2Ac2*, the catalytic subunit of the phosphatase, is lethal (168).

The precise manner in which the STRIPAK complex regulates ascomatal development is not fully understood. However, *CLA4*, a STRIPAK target might provide some insight into this pathway (166). This protein is a member of the p21-activated kinase family and

is important for sexual reproduction in *F. graminearum*, *N. crassa* and *M. grisea* (172–174). In *S. macrospora* specifically, $\Delta cla4$ strains produce deformed ascomata and significantly fewer asci and ascospores than wild type strains (166). The role that CLA4 plays in sexual development in *S. macrospora* is heavily dependent on a phosphorylation site within its catalytic subunit, a site that is directly regulated by STRIPAK. In fact, when this site was mutated so that it could not be dephosphorylated by STRIPAK, the production of asci and ascospores was greatly reduced, producing a phenotype comparable to $\Delta cla4$ (166). Further investigation into the other STRIPAK target proteins is likely to identify additional developmental pathways regulated by this complex.

The PACC Pathway

A second pathway that appears to be important in the development of the protoascomata in filamentous fungi, particularly in *N. crassa*, is the PACC pathway. This name reflects the final step of the pathway that involves the activation of the PACC transcription factor (175). This pathway has been implicated in pH response, the regulation of xylanase, cellulolytic and endoglucanase activities, and metabolism of glycogen and trehalose in *N. crassa* (176).

The PACC pathway has been particularly well-characterized in *S. cerevisiae* and *A. nidulans*, where it is activated when there are significant pH changes in the environment [Fig. 5, (177–179)]. The PALH protein acts as the pH sensor and is physically bound to the PALF protein, which is phosphorylated upon sensor stimulation. This results in

PALH being released and joining a complex of proteins including PALA, PALB, PALC and PACC. This in turn stimulates the activation of PACC via PALB-mediated cleavage. The activated PACC transcription factor is then translocated into the nucleus, where it brings about transcriptional change (175).

The genes that encode the PACC pathway proteins in *A. nidulans* do not appear to play an important role in sexual development. In contrast, genes encoding the six proteins of this pathway have been identified in the *N. crassa* genome and all are essential for the production of protoascomata (175). Deletion of any of the six genes that encode PACC pathway proteins result in mutants which, when grown on synthetic nitrogen-limited, crossing media, are unable to produce ascogonia or protoascomata. This phenotype was rescued when mutants were grown on medium containing the carbohydrates typically required for sexual development in *N. crassa* (175). Given that the pH-dependent PACC pathway is known to regulate the accumulation and metabolism of glycogen and trehalose in *N. crassa* (180), it is possible that the production of the highly complex protoascomata, which is an energetically intensive process, relies on the metabolism of these energy-rich carbohydrates, and hence the requirement of a functional PACC pathway.

The PACC pathway is also essential in *N. crassa* for steps downstream of protoascomatal production (175). In single gene mutants where protoascomatal formation had taken place, fertilization resulted in melanization and growth of the protoascomata into mature ascomata. However, the maturing ascomata did not reach

full size, did not harbor mature asci and no ascospores were produced, a phenotype that could not be rescued under different environmental conditions. This might be explained by the fact that the *S. cerevisiae* Rim1p, a *pacC* homolog, is important for meiosis (181, 182). Given that most of this pathway is conserved in *N. crassa* and *S. cerevisiae* (175), it is likely that the PACC pathway is also important for meiosis in *N. crassa*. Overall, this indicates that the PACC pathway is important in the initial as well as the later stages of sexual development.

The COP9 Signalosome

The COP9 signalosome (CSN) is another evolutionarily conserved complex that regulates sexual development in fungi. The complex regulates this pathway via its roles in response to oxidative stress, the regulation of redox status and cell wall rearrangement (183). The complex is made up of a variety of proteins, such as those involved in CSN assembly [eg. PCI-domain proteins, (184, 185)] and those with enzymatic activities [eg. metalloproteases, (184, 186)]. Together, these proteins act in post-translational processes, such as protein ubiquitylation and subsequent degradation (186). Via these important enzymatic activities, it is thought that the CSN complex plays an important role as a regulator of multicellularity (184, 187). While the specific manner in which this form of cellular differentiation remains to be understood, it appears that ubiquitin-dependent protein degradation is important for the development of complex, multicellular structures with specific spatial requirements (184).

While the CSN can be made up of as few as six or seven proteins, as is the case in *Schizosaccharomyces pombe* and *N. crassa* (186), the most common conformation of this Eukaryotic complex is exemplified by *A. nidulans* [Fig. 6, (184)]. The *A. nidulans* CSN is comprised of eight proteins, which are encoded by the genes named *cnsA* to *csnH*. Four of the eight genes have been functionally characterized with respect to their role in sexual development, including *csnA* (184), *csnB* (184), *csnD* (185) and *csnE* (185). Given the importance of this complex in the development of multicellular structures, it is perhaps not surprising that all four of these genes are important for the production of protoascomata. Furthermore, their deletion results in mutant isolates capable of forming only the initial structures (primordia) associated with sexual development, but these initials do not mature into protoascomata or any other mature sexual structures. Remarkably, even a single gene deletion results in this extreme phenotype, strongly suggesting that the functioning of this complex relies on all of the eight proteins (184, 185).

ASCOSPORE PRODUCTION & GERMINATION

The final stages of sexual reproduction include the production of the ascospores, their release into the environment and subsequent germination.

Ascosporegenesis

The production of ascospores relies on multiple rounds of cell division. Meiotic divisions within the ascus result in four haploid nuclei that are typically arranged in a linear conformation, each of which undergo a round of mitosis to produce the final eight

ascospores (115). Further ascospore development takes place to ensure the eight spores are arranged, partitioned and delineated correctly in the ascus.

There are various genes in *A. nidulans* and *N. crassa* that, when deleted, produce empty asci. This phenotype indicates that the appropriate sexual tissues are present, but that meiosis cannot be initiated within the mature asci. Four of these genes are *tubB* (188) and *grrA* (189) in *A. nidulans* and *sad1* (190) and *sad2* (191) in *N. crassa*. The two *A. nidulans* genes have very different functions; the *tubB* gene is involved in microtubule assembly (188), while *grrA* plays an important role in protein degradation (189). In contrast, both of the *N. crassa* genes are important for meiotic silencing (190, 191) and importantly, SAD2 ensures the proper localization and thus functioning of SAD1 (191). Despite difference in their precise functions, each of these genes enables the commencement of the meiotic cycle within the sexual structures.

The *ski8* gene in *S. macrospora* is essential for many different stages of ascosporegenesis, with functions as early as ascomatal production and ascus emergence and as late as meiosis (192). Importantly, it is thought that the SKI8 protein is particularly important for the production of double strand breaks in the DNA during meiosis, as $\Delta ski8$ mutants do not produce chiasmata, and they undergo meiosis aberrantly with delayed karyogamy. These mutants typically contain many asci in which unfused nuclei are common and mature ascospores have not been produced (192).

The *S. macrospora* STE12, a transcription factor that interacts directly with the MAT1-1-1 protein, is important for ascus and ascospore formation (193). When the *ste12* gene is deleted, mutants produce ascomata that appear normal but with very few asci, and those that are produced have very thin and fragile cell walls that burst easily. Similarly, the number of reproductive structures, including hooked cells and meiosis-specific cell stages, are significantly reduced in mutants compared to wild type isolates (193).

Several genes that encode proteins involved in autophagy in *S. macrospora* are important for production of asci and ascospores (194–196). Two such examples are *nbr1* and *lon2*, which encode an autophagy cargo receptor and a Lon protease, respectively (194, 195). Given that autophagy results in the reconstitution of cellular components and may thereby act as a mechanism of nutrient provision, particularly during periods of cellular development, it is not surprising that this pathway directly affects the production of highly specialized sexual tissues. The deletion of *nbr1* resulted in the production of immature, malformed asci which have significantly fewer ascospores than wild type asci (195). Similarly, $\Delta lon2$ mutants produced fewer ascomata, most of which contained asci housing only immature ascospores (194). Both mutants were more sensitive to stressors as well- with the $\Delta nbr1$ mutants not producing fruiting bodies under temperature and nutritional stress and $\Delta lon2$ mutants not producing fruiting bodies during starvation (194, 195). These results strongly suggest that the autophagous pathway is important in the environmental stress response, while maintaining the nutritional status necessary for sexual development.

In *N. crassa*, the *asd1* gene partially regulates ascosporegenesis once meiosis and mitosis have already taken place (197). This gene encodes a rhamnogalacturonase, which is thought to play a role in the regulation of sexual development, via the production of various polysaccharides. In Δ *asd1* mutants, the asci form normally but despite going through meiosis as usual, the spores do not delineate, and the haploid nuclei remain unpackaged (197). The *sec22* gene plays a similar role in *S. macrospora* where it is involved in ascospore maturation (198). In wild type isolates, the eight spores present in the asci develop at a uniform rate before being ejected together. In Δ *sec22* mutants, however, these spores mature at different rates and some are ejected well before maturation (198).

Germination

Ascospore germination has been studied in detail in *S. macrospora* and a number of genes that directly influence germination rate have been described. In addition to the role played by the autophagous pathway in the production of asci as described above, autophagy is also important for ascospore germination in *S. macrospora* (196). Two genes, *atg4* and *atg8*, encode autophagy-related proteins, and both directly affect the germination rate of ascospores in this species (196). Deletion of either *atg4* or *atg8* results in a germination rate reduction of about 50%. Since germination involves the growth of a mycelium from a spore, it is reasonable to speculate that autophagy may play an important role in providing nutrients to the germ tube. Given this role, it is not surprising that these two genes are also involved in the development of the maturing ascomata as well (196).

A third gene involved in ascospore germination in *S. macrospora* is *sec22* (198). Interestingly this gene, which encodes a membrane-associated protein involved in membrane fusion, may also be involved in autophagy (199), further supporting the role of this process in germination. In $\Delta sec22$ mutants, a mixture of immature and mature spores are released from the asci (198). The immature spores never germinate, and the mature spores germinate at a slower rate than the wild type (198).

As with *atg4* and *atg8*, *sec22* appears to have multiple roles in sexual development, ranging from spore production, pigmentation and maturation to ascospore germination post release (196, 198). Interestingly, deletion of *sec22* resulted in the deregulation of genes involved in numerous developmental pathways- including those putatively involved in vesicle transport (198). Given the large number of diverse molecular processes that rely on this cellular transport pathway, it is not surprising that the *sec22* gene is involved at numerous stages in the sexual cycle.

Another two genes in *S. macrospora* that influence germination are important for G-protein coupled signal transduction (152). These genes, *gsa3* and *sac1*, encode the $G\alpha$ -subunit and the adenylyl cyclase protein, respectively. Upon activation of the GPCR, the $G\alpha$ -subunit activates the adenylyl cyclase, thereby initiating an appropriate transduction cascade, which, in this case, would result in the formation of the germ tube. Deletion of the *gsa3* gene results in normal ascomata with ascospore-bearing asci, but only approximately 5% of these ascospores germinate (152). Similarly, $\Delta sac1$ mutants also

exhibit a significant decrease in germination but also produce much smaller ascomata during sexual development (152).

LOOKING TO THE FUTURE

Sexual reproduction in fungi has been studied for decades and the progress that has been made is nothing short of remarkable. From the original descriptions of different fungal fruiting bodies and mating behaviors, to the identification of the diffusible mating pheromones and the elucidation of the *MAT* genes, our understanding of this pathway has grown considerably, stretching from physiological observations to the characterization of the underlying genetic mechanisms that govern the process.

Studies on the pathways that govern sexual reproduction relied heavily on the functional characterization of single genes and their direct effect on the sexual capabilities of model fungi. This approach has proven very useful to shape our understanding of these process, but is plagued by some significant drawbacks. The first of these is that such a complex pathway is necessarily governed by extensive, multifaceted gene and protein networks that need to be characterized with respect to, and within the context of, one another. Additionally, a significant level of redundancy often exists when it comes to the function of these genes, with two or three proteins having overlapping functions and thus compensating for one another. This essentially masks the effects and importance of certain genes when they are knocked out. Lastly, it is often not possible to fully characterize the function of a gene if it is essential for numerous steps in a particular pathway. For example, if a gene is essential for both fertilization and the production of

asci, its role in ascus production will be masked because the block in fertilization will preclude any further sexual development. Thus, single gene knockouts do not always accurately allow for the characterization of the functions of these genes and cannot be used to describe all forms of inter-connected complexity. The advent of more powerful tools such as CRISPR-Cas9 and NGS promises to change this. Apart from the ability to produce single gene knockouts, CRISPR can efficiently produce multigene knockouts in a single organism, allowing complex pathways to be better dissected. With transformation protocols becoming more prevalent across the Pezizomycotina, functional studies in underrepresented taxa will become an exciting source of new information in future.

Currently, extensive data is available linking the functions of individual genes and genetic pathways to various steps of the sexual cycle in model species such as *N. crassa*, *A. nidulans*, *S. macrospora* and the other species discussed above (Fig. 3). However, a significant step forward in our understanding of sexual development in the Pezizomycotina has been the broadening of the field to include non-model species. Over the past few years, the sexual cycles of non-model species like *V. virens* (60, 65) and *H. omanensis* (64) have also been subjected to genetic inquiry. This has enabled the comparison of gene function of homologs from different species, which in turn enables further dissection of the evolutionary conservation of these processes. This is particularly important if we seek to continue using fungi as multicellular representatives of the Eukarya and extrapolating the conclusions drawn from fungi to other Eukaryotes.

A number of gaps exist in the literature (Fig. 3), paving the way for future investigations and a better understanding of sexual development in the Pezizomycotina. Some key genes and pathways remain to be elucidated- including the identification of the **a**-like pheromone in the Eurotiomycetes and related fungi, as well as the targets of the evolutionarily conserved protein complexes, STRIPAK and CSN. Additionally, much work remains to be done on how various environmental cues regulate sexual development. For example, the effect of light on the sexual cycles of *N. crassa* and *A. nidulans* is very well understood both in terms of the environmental stimuli and the genetic and physiological responses. However, the importance of other environmental factors such as pH, temperature and nutrient availability are less well understood in filamentous species, particularly with regards to the genes and proteins involved in regulating these pathways.

The role of post-transcriptional and post-translational modification clearly requires further investigation. It is known, for example, that RNA editing is important for ascosporegenesis and ascus maturation in both *F. graminearum* (200) and *N. crassa* (201). Similarly, post-translational modifications are known to be essential for the functioning of both pheromone factors. However, the extent to which these types of modifications are necessary and essential for the sexual cycle is not yet known. Whether such modifications occur in all of the Pezizomycotina also remains a topic open for debate.

CONCLUDING REMARKS

Fungi provide fascinating and effective models for studies focused on elucidating a process as complicated as sexual reproduction. Fungi exhibit numerous distinct and, in some cases, unusual sexual strategies and there are a growing number of model species available for study and comparison. The fungi also include an unusually wide diversity of morphologies, sexual structures and preferred environments providing a wide array of opportunities to study sexual development in depth.

New and emerging technologies are driving opportunities to better understand sexual reproduction in the fungi. For example, genome editing, whole genome sequencing and comparative transcriptomics are becoming increasingly accessible and available for such investigations. These are likely to reveal a substantially improved understanding of sexual development in a diverse array of model and non-model fungal species. And the results will make it possible to extrapolate emerging knowledge beyond the fungal Kingdom to the other Eukaryotes.

ACKNOWLEDGEMENTS

This project was supported by the University of Pretoria, the Department of Science and Innovation (DSI)/National Research Foundation (NRF) Centre of Excellence in Plant Health Biotechnology (CPHB). The project was further supported by the DSI/NRF SARChI chair in Fungal Genomics (Grant number: 98353) held by Prof. Brenda D. Wingfield. The grant holders acknowledge that the opinions, findings and conclusions or recommendations expressed are theirs and that the funding bodies accept no liability in

this regard. The authors declare no conflict of interest. We would also like to thank Glenda Brits of the Department of Education Innovation at the University of Pretoria for producing the figures used in this review.

REFERENCES

1. Speijer D, Lukeš J, Eliáš M. 2015. Sex is a ubiquitous, ancient, and inherent attribute of Eukaryotic life. *Proc Natl Acad Sci* 112:8827–8834.
2. Nieuwenhuis BPS, James TY. 2016. The frequency of sex in fungi. *Philos Trans R Soc B Biol Sci* 371.
3. Nieuwenhuis BPS, Aanen DK. 2012. Sexual selection in fungi. *J Evol Biol* 25:2397–2411.
4. Bennett RJ, Turgeon BG. 2016. Fungal sex: The Ascomycota. *Microbiol Spectr* 4:1–28.
5. Wallen RM, Perlin MH. 2018. An overview of the function and maintenance of sexual reproduction in dikaryotic fungi. *Front Microbiol* 9:1–24.
6. Butler G. 2007. The evolution of *MAT*: The Ascomycetes, p. 3–18. *In* Heitman, J, Kronstad, JW, Taylor, J, Casselton, LA (eds.), *Sex in Fungi: Molecular Determination and Evolutionary Implications*. ASM Press, Washington, DC.
7. Jones SK, Bennett RJ. 2011. Fungal mating pheromones: Choreographing the dating game. *Fungal Genet Biol* 48:668–676.
8. Lumbsch HT, Huhndorf SM. 2010. Myconet. *Fieldiana Life Earth Sci* 14:1–64.
9. Usher J. 2019. The mechanisms of mating in pathogenic fungi- A plastic trait. *Genes* 10:16–18.

10. Sun S, Lin X, Coelho MA, Heitman J. 2020. Evolution of yeast mating systems: All roads lead to selfing. *Curr Biol* 29:1–6.
11. Latgé JP, Chamilos G. 2019. *Aspergillus fumigatus* and Aspergillosis in 2019. *Clin Microbiol Rev* 33:1–75.
12. Nenoff P, Krüger C, Ginter-Hanselmayer G, Tietz HJ. 2014. Mycology - An update. Part 1: Dermatomycoses: Causative agents, epidemiology and pathogenesis. *JDDG - J Ger Soc Dermatology* 12:188–210.
13. Rigling D, Prospero S. 2018. *Cryphonectria parasitica*, the causal agent of chestnut blight: Invasion history, population biology and disease control. *Mol Plant Pathol* 19:7–20.
14. Wingfield MJ, Hammerbacher A, Ganley RJ, Steenkamp ET, Gordon TR, Wingfield BD, Coutinho TA. 2008. Pitch canker caused by *Fusarium circinatum* - A growing threat to pine plantations and forests worldwide. *Australas Plant Pathol* 37:319–334.
15. Gueidan C, Hill D, Miadlikowska J, Lutzoni FL. 2015. Pezizomycotina: Lecanoromycetes, p. 89–120. *In* McLaughlin, D, Spatafora, J (eds.), *Systematics and Evolution*, 2nd ed. Springer, Berlin.
16. Spatafora JW, Sung G-H, Johnson D, Hesse C, O'Rourke B, Serdani M, Spotts R, Lutzoni F, Hofstetter V, Miadlikowska J, Reeb V, Gueidan C, Fraker E, Lumbsch T, Lücking R, Schmitt I, Hosaka K, Aptroot A, Roux C, Miller AN, Geiser DM, Hafellner J, Hestmark G, Arnold AE, Büdel B, Rauhut A, Hewitt D, Untereiner WA, Cole MS, Scheidegger C, Schultz M, Sipman H, Schoch CL. 2006. A five-gene phylogeny of Pezizomycotina. *Mycologia* 98:1018–1028.

17. Roche CM, Loros JJ, McCluskey K, Glass N. 2014. *Neurospora crassa*: Looking back and looking forward at a model microbe. *Am J Bot* 101(12):2022-2035.
18. Bayram Ö, Braus GH, Fischer R, Rodriguez-Romero J. 2010. Spotlight on *Aspergillus nidulans* photosensory systems. *Fungal Genet Biol* 47:900–908.
19. Bayram Ö, Braus GH. 2012. Coordination of secondary metabolism and development in fungi: The velvet family of regulatory proteins. *FEMS Microbiol Rev* 36:1–24.
20. Bayram ÖS, Dettmann A, Karahoda B, Moloney NM, Ormsby T, McGowan J, Cea-Sánchez S, Miralles-Durán A, Brancini GTP, Luque EM, Fitzpatrick DA, Cánovas D, Corrochano LM, Doyle S, Selker EU, Seiler S, Bayram Ö. 2019. Control of development, secondary metabolism and light-dependent carotenoid biosynthesis by the velvet complex of *Neurospora crassa*. *Genetics* 212:691–710.
21. Patel PK, Free SJ. 2019. The genetics and biochemistry of cell wall structure and synthesis in *Neurospora crassa*, a model filamentous fungus. *Front Microbiol* 10:1–18.
22. Alkhayyat F, Chang Kim S, Yu JH. 2015. Genetic control of asexual development in *Aspergillus fumigatus*. *Adv Appl Microbiol* 90:93–107.
23. Ruger-Herreros C, Corrochano LM. 2020. Conidiation in *Neurospora crassa*: Vegetative reproduction by a model fungus. *Int Microbiol* 23:97–105.
24. Kim H, Wright SJ, Park G, Ouyang S, Krystofova S, Borkovich KA. 2012. Roles for receptors, pheromones, G-proteins, and mating-type genes during sexual reproduction in *Neurospora crassa*. *Genetics* 190:1389–1404.
25. Ferreira AVB, An Z, Metzzenberg RL, Glass NL. 1998. Characterization of *mat A-2*,

- mat A-3* and *ΔmatA* mating-type mutants of *Neurospora crassa*. *Genetics* 148:1069–1079.
26. Dyer PS, O’Gorman CM. 2012. Sexual development and cryptic sexuality in fungi: Insights from *Aspergillus* species. *FEMS Microbiol Rev* 36:165–192.
 27. Lim JY, Kang EH, Park YH, Kook JH, Park HM. 2020. Survival factor *SvfA* plays multiple roles in differentiation and is essential for completion of sexual development in *Aspergillus nidulans*. *Sci Rep* 10:1–13.
 28. Debuchy R, Berteaux-Lecellier V, Silar P. 2010. Mating systems and sexual morphogenesis in Ascomycetes, p. 501–535. *In* Borkovich, KA, Ebbole, DJ (eds.), *Cellular and molecular biology of filamentous fungi*. ASM Press, Washington D.C.
 29. Astell CR, Ahlstrom-Jonasson L, Smith M, Tatchell K, Nasmyth KA, Hall BD. 1981. The sequence of the DNAs coding for the mating-type loci of *Saccharomyces cerevisiae*. *Cell* 27:15–23.
 30. Brake A, Brenner C, Najarian R, Laybourn P, Merryweather J. 1985. Structure of genes encoding precursors of the yeast peptide mating pheromone α -factor, p. 103–108. *In* Gething, M (ed.), *Protein transport and secretion*. Cold Spring Harbor Laboratory Press, New York.
 31. Kurjan J, Herskowitz I. 1982. Structure of a yeast pheromone gene (*MF α*): A putative α -factor precursor contains four tandem copies of mature α -factor. *Cell* 30:933–943.
 32. Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman JR, Batzoglou S, Lee SI, Baştürkmen M, Spevak CC, Clutterbuck J, Kapitonov V, Jurka J, Scazzocchio C, Farman M, Butler J, Purcell S, Harris S, Braus GH, Draht O, Busch S, D’Enfert C,

- Bouchier C, Goldman GH, Bell-Pedersen D, Griffiths-Jones S, Doonan JH, Yu J, Vienken K, Pain A, Freitag M, Selker EU, Archer DB, Peñalva MÁ, Oakley BR, Momany M, Tanaka T, Kumagai T, Asai K, Machida M, Nierman WC, Denning DW, Caddick M, Hynes M, Paoletti M, Fischer R, Miller B, Dyer P, Sachs MS, Osmani SA, Birren BW. 2005. Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* 438:1105–1115.
33. Glass NL, Vollmer SJ, Staben C, Grotelueschen J, Metzzenberg RL, Yanofsky C. 1988. DNAs of the two mating-type alleles of *Neurospora crassa* are highly dissimilar. *Science* 241:570–573.
34. Debuchy R, Coppin E. 1992. The mating types of *Podospora anserina*: Functional analysis and sequence of the fertilization domains. *Mol Gen Genet* 233:113–121.
35. Bobrowicz P, Pawlak R, Correa A, Bell-Pedersen D, Ebbole DJ. 2002. The *Neurospora crassa* pheromone precursor genes are regulated by the mating-type locus and the circadian clock. *Mol Microbiol* 45:795–804.
36. Dyer PS, Paoletti M, Archer DB. 2003. Genomics reveals sexual secrets of *Aspergillus*. *Microbiology* 149:2301–2303.
37. Coppin E, de Renty C, Debuchy R. 2004. The function of the coding sequences for the putative pheromone precursors in *Podospora anserina* is restricted to fertilization. *Eukaryot Cell* 4:407–420.
38. Böhm J, Hoff B, O’Gorman CM, Wolfers S, Klix V, Binger D, Zadra I, Kürnsteiner H, Pöggeler S, Dyer PS, Kück U. 2013. Sexual reproduction and mating type-mediated strain development in the penicillin-producing fungus *Penicillium chrysogenum*. *Proc Natl Acad Sci* 110:1476–1481.

39. Mayrhofer S, Pöggeler S. 2005. Functional characterization of an α -factor-like *Sordaria macrospora* peptide pheromone and analysis of its interaction with its cognate receptor in *Saccharomyces cerevisiae*. *Eukaryot Cell* 4:661–672.
40. Haber JE. 2012. Mating-type genes and *MAT* switching in *Saccharomyces cerevisiae*. *Genetics* 191:33–64.
41. Pöggeler S, Nowrousian M, Kück U. 2018. Fruiting-body development in Ascomycetes, p. 1–56. *In* Anke, T, Schüffler, A (eds.), *Physiology and Genetics*, 2nd ed. Springer International Publishing, Basel, Switzerland.
42. Coppin E, Debuchy R, Arnaise S, Picard M. 1997. Mating types and sexual development in filamentous Ascomycetes. *Microbiol Mol Biol Rev* 61:411–428.
43. Arnaise S, Zickler D, Le Bilot S, Poisier C, Debuchy R. 2001. Mutations in mating-type genes of the heterothallic fungus *Podospora anserina* lead to self-fertility. *Genetics* 159:545–556.
44. Shiu PKT, Glass NL. 2000. Cell and nuclear recognition mechanisms mediated by mating type in filamentous Ascomycetes. *Curr Opin Microbiol* 3:183–188.
45. Wilson AM, Markus Wilken P, Van Der Nest MA, Steenkamp ET, Wingfield MJ, Wingfield BD. 2015. Homothallism: An umbrella term for describing diverse sexual behaviours. *IMA Fungus* 6:207–214.
46. Lin X, Heitman J. 2007. Mechanisms of homothallism in fungi and transitions between heterothallism and homothallism, p. 35–57. *In* Heitman, J, Kronstad, JW, Taylor, JW, Casselton, LA (eds.), *Sex in Fungi*. ASM Press, Washington D.C.
47. Merino ST, Nelson MA, Jacobson DJ, Natvig DO. 1996. Pseudohomothallism and evolution of the mating-type chromosome in *Neurospora tetrasperma*. *Genetics*

- 143:789–799.
48. Chitrampalam P, Inderbitzin P, Maruthachalam K, Wu BM, Subbarao K V. 2013. The *Sclerotinia sclerotiorum* mating-type locus (*MAT*) contains a 3.6-kb region that is inverted in every meiotic generation. PLoS One 8:e56895.
 49. Wilken PM, Steenkamp ET, Wingfield MJ, De Beer ZW, Wingfield BD. 2014. DNA loss at the *Ceratocystis fimbriata* mating locus results in self-sterility. PLoS One 9:e92180.
 50. Perkins D. 1987. Mating type switching in filamentous Ascomycetes. Genetics 115:215–216.
 51. Roach KC, Feretzaki M, Sun S, Heitman J. 2014. Unisexual reproduction, p. 255–305. In Friedmann, T, Dunlap, JC, Goodwin, SF (eds.), Advances in Genetics, 85th ed. Academic Press, Waltham.
 52. Pöggeler S, Nowrousian M, Ringelberg C, Loros JJ, Dunlap JC, Kück U. 2006. Microarray and real-time PCR analyses reveal mating type-dependent gene expression in a homothallic fungus. Mol Genet Genomics 275:492–503.
 53. Bidard F, Benkhali J, Coppin E, Imbeaud S, Grognet P, Delacroix H, Debuchy R. 2011. Genome-wide gene expression profiling of fertilization competent mycelium in opposite mating types in the heterothallic fungus *Podospora anserina*. PLoS One 6:e21476.
 54. Yoder OC, Valent B, Chumley FG. 1986. Genetic nomenclature and practice for plant pathogenic fungi. Phytopathology 76:383–385.
 55. Metzberg RL, Louise Glass N. 1990. Mating type and mating strategies in *Neurospora*. BioEssays 12:53–59.

56. Wilken PM, Steenkamp ET, Wingfield MJ, de Beer ZW, Wingfield BD. 2017. Which *MAT* gene? Pezizomycotina (Ascomycota) mating-type gene nomenclature reconsidered. *Fungal Biol Rev* 31:199–211.
57. Petters-Vandresen DAL, Rossi BJ, Groenewald J, Crous P, Machado MA, Stukenbrock E, Glienke C. 2020. Mating-type locus rearrangements and shifts in thallism states in *Citrus*-associated *Phyllosticta* species. *Fungal Genet Biol* 144:103444.
58. Nagel JH, Wingfield MJ, Slippers B. 2018. Evolution of the mating types and mating strategies in prominent genera in the Botryosphaeriaceae. *Fungal Genet Biol* 114:24–33.
59. Arnaise S, Debuchy R, Picard M. 1997. What is a bona fide mating-type gene? Internuclear complementation of *mat* mutants in *Podospora anserina*. *Mol Gen Genet* 256:169–178.
60. Yong M, Yu J, Pan X, Yu M, Cao H, Song T, Qi Z, Du Y, Zhang R, Yin X, Liu W, Liu Y. 2020. Two mating-type genes *MAT1-1-1* and *MAT1-1-2* with significant functions in conidiation, stress response, sexual development, and pathogenicity of rice false smut fungus *Villosiclava virens*. *Curr Genet* 66:989–1002.
61. Debuchy R, Turgeon B. 2006. Mating-type structure, evolution, and function in Eucaryotes, p. 293–323. In Kües, U, Fischer, R (eds.), *The Mycota: Growth, Differentiation and Sexuality*, 1st ed. Springer, Berlin.
62. Wilson A, Wilken P, van der Nest M, Wingfield M, Wingfield B. 2019. It's all in the genes: The regulatory pathways of sexual reproduction in filamentous ascomycetes. *Genes* 10:330.

63. Yu Y, Amich J, Will C, Eagle CE, Dyer PS, Krappmann S. 2017. The novel *Aspergillus fumigatus* *MAT1-2-4* mating-type gene is required for mating and cleistothecia formation. *Fungal Genet Biol* 108:1–12.
64. Wilson AM, Wilken PM, Nest MA Van Der, Wing MJ, Wing BD. 2020. The novel *Huntia omanensis* mating gene, *MAT1-2-7*, is essential for ascomatal maturation. *Fungal Genet Biol* 137:103335.
65. Yong M, Yu J, Pan X, Yu M, Cao H, Qi Z, Du Y, Zhang R, Song T, Yin X, Chen Z, Liu W, Liu Y. 2020. *MAT1-1-3*, a mating-type gene in the *Villosiclava virens*, is required for fruiting bodies and sclerotia formation, asexual development and pathogenicity. *Front Microbiol* 11:1–18.
66. Kanzi AM, Steenkamp ET, Van der Merwe NA, Wingfield BD. 2019. The mating system of the *Eucalyptus* canker pathogen *Chrysosporthe austroafricana* and closely related species. *Fungal Genet Biol* 123:41–52.
67. Mandel MA, Barker BM, Kroken S, Rounsley SD, Orbach MJ. 2007. Genomic and population analyses of the mating-type loci in *Coccidioides* species reveal evidence for sexual reproduction and gene acquisition. *Eukaryot Cell* 6:1189–1199.
68. Chai H, Chen W, Zhang X, Su K, Zhao Y. 2019. Structural variation and phylogenetic analysis of the mating-type locus in the genus *Morchella*. *Mycologia* 111:551–562.
69. Aylward J, Havenga M, Dreyer LL, Roets F, Wingfield BD, Wingfield MJ. 2020. Genomic characterization of mating-type loci and mating type distribution in two apparently asexual plantation tree pathogens. *Plant Pathol* 69:28–37.

70. Li JQ, Wingfield BD, Wingfield MJ, Barnes I, Fourie A, Crous PW, Chen SF. 2020. Mating genes in *Calonectria* and evidence for a heterothallic ancestral state. *Persoonia* 45:163–176.
71. Wilson AM, Godlonton T, van der Nest MA, Wilken PM, Wingfield MJ, Wingfield BD. 2015. Unisexual reproduction in *Huntia moniliformis*. *Fungal Genet Biol* 80:1–9.
72. Kanematsu S, Adachi Y, Ito T. 2007. Mating-type loci of heterothallic *Diaporthe* spp.: Homologous genes are present in opposite mating types. *Curr Genet* 52:11–22.
73. Turgeon BG, Yoder OC. 2000. Proposed nomenclature for mating type genes of filamentous ascomycetes. *Fungal Genet Biol* 31:1–5.
74. Glass NL, Grotelueschen J, Metzenberg RL. 1990. *Neurospora crassa* A mating-type region. *Proc Natl Acad Sci* 87:4912–4916.
75. Staben C, Yanofsky C. 1990. *Neurospora crassa* a mating-type region. *Genetics* 87:4917–4921.
76. Štros M, Launholt D, Grasser KD. 2007. The HMG-box: A versatile protein domain occurring in a wide variety of DNA-binding proteins. *Cell Mol Life Sci* 64:2590–2606.
77. Arie T, Christiansen SK, Yoder OC, Turgeon BG. 1997. Efficient cloning of ascomycete mating type genes by PCR amplification of the conserved *MAT* HMG box. *Fungal Genet Biol* 21:118–130.
78. Arie T, Kaneko I, Yoshida T, Noguchi M, Nomura Y, Yamaguchi I. 2007. Mating-type genes from asexual phytopathogenic ascomycetes *Fusarium oxysporum* and

- Alternaria alternata*. Mol Plant-Microbe Interact 13:1330–1339.
79. Aylward J. 2017. Comparative genomics of *Knoxdaviesia* species in the Core Cape Subregion. Stellenbosch University.
 80. Wilson AM, Lelwala R V., Taylor PWJ, Wingfield MJ, Wingfield BD. 2021. Unique patterns of mating pheromone presence and absence could result in the ambiguous sexual behaviours of *Colletotrichum* species. G3 Genes, Genomes, Genet DOI: 10.1093/g3journal/jkab187.
 81. Ament-Velásquez SL, Tuovinen V, Bergström L, Spribille T, Vanderpool D, Nascimbene J, Yamamoto Y, Thor G, Johannesson H. 2021. The plot thickens: Haploid and triploid-Like thalli, hybridization, and biased mating type ratios in *Letharia*. Front Fungal Biol 2:656386.
 82. Krämer D, Lane FA, Steenkamp ET, Wingfield BD, Wilken PM. 2021. Unidirectional mating-type switching confers self-fertility to *Thielaviopsis cerberus*, the only homothallic species in the genus. Fungal Biol 125:427–434.
 83. Wilken PM, Steenkamp ET, van der Nest MA, Wingfield MJ, de Beer ZW, Wingfield BD. 2018. Unexpected placement of the *MAT1-1-2* gene in the *MAT1-2* idiomorph of *Thielaviopsis*. Fungal Genet Biol 113:32–41.
 84. Havenga M, Wingfield BD, Wingfield MJ, Roets F, Dreyer LL, Tatham CT, Duong TA, Wilken PM, Chen SF, Aylward J. 2020. Mating strategy and mating type distribution in six global populations of the *Eucalyptus* foliar pathogen *Teratosphaeria destructans*. Fungal Genet Biol 137:103350.
 85. Klix V, Nowrousian M, Ringelberg C, Loros JJ, Dunlap JC, Pöggeler S. 2010. Functional characterization of *MAT1-1*-specific mating-type genes in the

- homothallic ascomycete *Sordaria macrospora* provides new insights into essential and non-essential sexual regulators. *Eukaryot Cell* 9:894–905.
86. Debuchy R, Arnais S, Lecellier G. 1993. The *mat-* allele of *Podospora anserina* contains three regulatory genes required for the development of fertilized female organs. *Mol Gen Genet* 241:667–673.
 87. Paoletti M, Seymour FA, Alcocer MJC, Kaur N, Calvo AM, Archer DB, Dyer PS. 2007. Mating type and the genetic basis of self-fertility in the model fungus *Aspergillus nidulans*. *Curr Biol* 17:1384–1389.
 88. Doughan B, Rollins JA. 2016. Characterization of *MAT* gene functions in the life cycle of *Sclerotinia sclerotiorum* reveals a lineage-specific *MAT* gene functioning in apothecium morphogenesis. *Fungal Biol* 120:1105–1117.
 89. Pöggeler S, Risch S, Kück U, Osiewacz HD. 1997. Mating-type genes from the homothallic fungus *Sordaria macrospora* are functionally expressed in a heterothallic ascomycete. *Genetics* 147:567–580.
 90. Becker K, Beer C, Freitag M, Kück U. 2015. Genome-wide identification of target genes of a mating-type α -domain transcription factor reveals functions beyond sexual development. *Mol Microbiol* 96:1002–1022.
 91. Wilson AM, van der Nest MA, Wilken PM, Wingfield MJ, Wingfield BD. 2018. Pheromone expression reveals putative mechanism of unisexuality in a saprobic ascomycete fungus. *PLoS One* 13:e0192517.
 92. Rodenburg SYA, Terhem RB, Veloso J, Stassen JHM, van Kan JAL. 2018. Functional analysis of mating-type genes and transcriptome analysis during fruiting body development of *Botrytis cinerea*. *MBio* 9:1–19.

93. Lee J, Lee T, Lee YW, Yun SH, Turgeon BG. 2003. Shifting fungal reproductive mode by manipulation of mating-type genes: Obligatory heterothallism of *Gibberella zeae*. *Mol Microbiol* 50:145–152.
94. Böhm J, Dahlmann TA, Gümüşer H, Kück U. 2015. A MAT1-2 wild-type strain from *Penicillium chrysogenum*: Functional mating-type locus characterization, genome sequencing and mating with an industrial penicillin-producing strain. *Mol Microbiol* 95:859–874.
95. Kim W, Cavinder B, Proctor RH, O'Donnell K, Townsend JP, Trail F. 2019. Comparative genomics and transcriptomics during sexual development gives insight into the life history of the cosmopolitan fungus *Fusarium neocosmosporiellum*. *Front Microbiol* 10:1–15.
96. Zheng Q, Hou R, Juanyu, Zhang, Ma J, Wu Z, Wang G, Wang C, Xu JR. 2013. The *MAT* locus genes play different roles in sexual reproduction and pathogenesis in *Fusarium graminearum*. *PLoS One* 8:e66980.
97. Kim HK, Cho EJ, Lee S, Lee YS, Yun SH. 2012. Functional analyses of individual mating-type transcripts at *MAT* loci in *Fusarium graminearum* and *Fusarium asiaticum*. *FEMS Microbiol Lett* 337:89–96.
98. Dyer PS, Inderbitzin P, Debuchy R. 2016. Mating-type structure, function, regulation and evolution in the Pezizomycotina, p. 351–385. *In* Wendland, J (ed.), *Growth, Differentiation and Sexuality*, 3rd ed. Springer International Publishing, Switzerland.
99. Ait Benkhali J, Coppin E, Brun S, Peraza-Reyes L, Martin T, Dixelius C, Lazar N, van Tilbeurgh H, Debuchy R. 2013. A network of HMG-box transcription factors

- regulates sexual cycle in the fungus *Podospora anserina*. PLoS Genet 9:e1003642.
100. Gautier V, Tong L, Nguyen T-S, Debuchy R, Silar P. 2018. PaPro1 and IDC4, two genes controlling stationary phase, sexual development and cell degeneration in *Podospora anserina*. J Fungi 4:1–28.
 101. Bazafkan H, Dattenböck C, Böhmendorfer S, Tisch D, Stappler E, Schmoll M. 2015. Mating type-dependent partner sensing as mediated by VEL1 in *Trichoderma reesei*. Mol Microbiol 96:1103–1118.
 102. Seibel C, Tisch D, Kubicek CP, Schmoll M. 2012. ENVOY is a major determinant in regulation of sexual development in *Hypocrea jecorina* (*Trichoderma reesei*). Eukaryot Cell 11:885–895.
 103. Aylward J, Steenkamp ET, Dreyer LL, Roets F, Wingfield MJ, Wingfield BD. 2016. Genetic basis for high population diversity in *Protea*-associated *Knoxdaviesia*. Fungal Genet Biol 96:47–57.
 104. Nauta MJ, Hoekstra RF. 1992. Evolution of reproductive systems in filamentous ascomycetes. I. Evolution of mating types. Heredity 68:405–410.
 105. Leslie JF, Klein KK. 1996. Female fertility and mating type effects on effective population size and evolution in filamentous fungi. Genetics 144:557–567.
 106. Kim H, Borkovich KA. 2006. Pheromones are essential for male fertility and sufficient to direct chemotropic polarized growth of trichogynes during mating in *Neurospora crassa*. Eukaryot Cell 5:544–554.
 107. Kim H, Borkovich KA. 2004. A pheromone receptor gene, *pre-1*, is essential for mating type-specific directional growth and fusion of trichogynes and female

- fertility in *Neurospora crassa*. Mol Microbiol 52:1781–1798.
108. Cano-Domínguez N, Álvarez-Delfín K, Hansberg W, Aguirre J. 2008. NADPH oxidases NOX-1 and NOX-2 require the regulatory subunit NOR-1 to control cell differentiation and growth in *Neurospora crassa*. Eukaryot Cell 7:1352–1361.
 109. Kim H, Metzenberg RL, Nelson MA. 2002. Multiple functions of *mfa-1*, a putative pheromone precursor gene of *Neurospora crassa*. Eukaryot Cell 1:987–999.
 110. Park G, Pan S, Borkovich KA. 2008. Mitogen-activated protein kinase cascade required for regulation of development and secondary metabolism in *Neurospora crassa*. Eukaryot Cell 7:2113–2122.
 111. Yang Q, Poole SI, Borkovich KA. 2002. A G-protein β subunit required for sexual and vegetative development and maintenance of normal $G\alpha$ protein levels in *Neurospora crassa*. Eukaryot Cell 1:378–390.
 112. Turina M, Prodi A, Van Alfen NK. 2003. Role of the *Mf1-1* pheromone precursor gene of the filamentous ascomycete *Cryphonectria parasitica*. Fungal Genet Biol 40:242–251.
 113. Zhou X, Liu W, Wang C, Xu Q, Wang Y, Ding S, Xu JR. 2011. A MADS-box transcription factor MoMcm1 is required for male fertility, microconidium production and virulence in *Magnaporthe oryzae*. Mol Microbiol 80:33–53.
 114. Malagnac F, Lalucque H, Lepère G, Silar P. 2004. Two NADPH oxidase isoforms are required for sexual reproduction and ascospore germination in the filamentous fungus *Podospora anserina*. Fungal Genet Biol 41:982–997.
 115. Silar P. 2014. Simple genetic tools to study fruiting body development in fungi. Open Mycol J 8:148–155.

116. Lalucque H, Malagnac F, Green K, Gautier V, Grognet P, Chan Ho Tong L, Scott B, Silar P. 2017. *IDC2* and *IDC3*, two genes involved in cell non-autonomous signaling of fruiting body development in the model fungus *Podospora anserina*. *Dev Biol* 421:126–138.
117. Barman A, Tamuli R. 2017. The pleiotropic vegetative and sexual development phenotypes of *Neurospora crassa* arise from double mutants of the calcium signaling genes *plc-1*, *splA2*, and *cpe-1*. *Curr Genet* 63:861–875.
118. Cavinder B, Trail F. 2012. Role of Fig1, a component of the low-affinity calcium uptake system, in growth and sexual development of filamentous fungi. *Eukaryot Cell* 11:978–988.
119. Bolker M, Kahmann R. 1993. Sexual pheromones and mating responses in fungi. *Plant Cell* 5:1461–1469.
120. Zhang L, Baasiri RA, Van Alfen NK. 1998. Viral repression of fungal pheromone precursor gene expression. *Mol Cell Biol* 18:953–959.
121. Shen WC, Bobrowicz P, Ebole DJ. 1999. Isolation of pheromone precursor genes of *Magnaporthe grisea*. *Fungal Genet Biol* 27:253–263.
122. Lee J, Leslie JF, Bowden RL. 2008. Expression and function of sex pheromones and receptors in the homothallic ascomycete *Gibberella zeae*. *Eukaryot Cell* 7:1211–1221.
123. Pöggeler S. 2000. Two pheromone precursor genes are transcriptionally expressed in the homothallic ascomycete *Sordaria macrospora*. *Curr Genet* 37:403–411.
124. Pöggeler S, Kück U. 2001. Identification of transcriptionally expressed pheromone

- receptor genes in filamentous ascomycetes. *Gene* 280:9–17.
125. Wilson AM, Gabriel R, Singer SW, Schuerg T, Wilken PM, van der Nest MA, Wingfield MJ, Wingfield BD. 2021. Doing it alone: Unisexual reproduction in filamentous ascomycete fungi. *Fungal Biol Rev* 35:1–13.
 126. Martin F, Kohler A, Murat C, Balestrini R, Coutinho PM, Jaillon O, Montanini B, Morin E, Noel B, Percudani R, Porcel B, Rubini A, Amicucci A, Amselem J, Anthouard V, Arcioni S, Artiguenave F, Aury JM, Ballario P, Bolchi A, Brenna A, Brun A, Buée M, Cantarel B, Chevalier G, Couloux A, Da Silva C, Denoeud F, Duplessis S, Ghignone S, Hilselberger B, Iotti M, Marçais B, Mello A, Miranda M, Pacioni G, Quesneville H, Riccioni C, Ruotolo R, Splivallo R, Stocchi V, Tisserant E, Viscomi AR, Zambonelli A, Zampieri E, Henrissat B, Lebrun MH, Paolocci F, Bonfante P, Ottonello S, Wincker P. 2010. Périgord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* 464:1033–1038.
 127. Zhang L, Churchill AC, Kazmierczak P, Kim DH, Van Alfen NK. 1993. Hypovirulence-associated traits induced by a mycovirus of *Cryphonectria parasitica* are mimicked by targeted inactivation of a host gene. *Mol Cell Biol* 13:7782–92.
 128. Mayrhofer S, Weber JM, Pöggeler S. 2006. Pheromones and pheromone receptors are required for proper sexual development in the homothallic ascomycete *Sordaria macrospora*. *Genetics* 172:1521–1533.
 129. Liang X, Yao L, Kong Y, Li B, Hao X, Lin Y, Cao M, Dong Q, Zhang R, Rollins JA, Sun G. 2021. Molecular dissection of perithecial mating line development in *Colletotrichum fructicola*, a species with a nontypical mating system featured by

- plus-to-minus switch and plus-minus mediated sexual enhancement. *Appl Environ Microbiol* 87:e00474-21.
130. Wheeler H. 1954. Genetics and evolution of heterothallism in *Glomerella*. *Phytopathology* 44:342–345.
 131. Anderegg RJ, Betzll R, Carr SA, Crabb JW, Duntzell W, Chern WJB. 1988. Structure of *Saccharomyces cerevisiae* mating hormone a-factor. *J Biol Chem* 263:18236–18240.
 132. Pöggeler S. 2002. Genomic evidence for mating abilities in the asexual pathogen *Aspergillus fumigatus*. *Curr Genet* 42:153–160.
 133. Ge YY, Yu FM, Yang ZJ, Tan YM, Xiang T, Chen X, Yu YX, Chen L, Liu ZY. 2019. Genetic basis and function of mating-type genes in *Aspergillus cristatus*. *Mycosphere* 10:622–633.
 134. Pál K, Van Diepeningen AD, Varga J, Debets AJM, Hoekstra RF. 2008. Sexual genes in the asexual filamentous fungus *Aspergillus niger* and related *Aspergilli*. *Aspergillus Genomic Era* 107–131.
 135. Aboelfotoh Hendy A, Xing J, Chen X, Chen XL. 2019. The farnesyltransferase β -subunit RAM1 regulates localization of RAS proteins and appressorium-mediated infection in *Magnaporthe oryzae*. *Mol Plant Pathol* 20:1264–1278.
 136. Short DPG, Gurung S, Hu X, Inderbitzin P, Subbarao K V. 2014. Maintenance of sex-related genes and the co-occurrence of both mating types in *Verticillium dahliae*. *PLoS One* 9:e112145.
 137. Maurer-Stroh S, Washietl S, Eisenhaber F. 2003. Protein prenyltransferases. *Genome Biol* 4:212.

138. Ge Y, Yu F, Tan Y, Zhang X, Liu Z. 2017. Comparative transcriptome sequence analysis of sporulation-related genes of *Aspergillus cristatus* in response to low and high osmolarity. *Curr Microbiol* 74:806–814.
139. Nowrousian M, Stajich JE, Chu M, Engh I, Espagne E, Halliday K, Kamerewerd J, Kempken F, Knab B, Kuo HC, Osiewacz HD, Pöggeler S, Read ND, Seiler S, Smith KM, Zickler D, Kück U, Freitag M. 2010. *De novo* assembly of a 40 Mb eukaryotic genome from short sequence reads: *Sordaria macrospora*, a model organism for fungal morphogenesis. *PLoS Genet* 6:e1000891.
140. Hernández-Chávez MJ, González-Hernández RJ, Trujillo-Esquivel JE, Hernández-Cervantes A, Mora-Montes HM. 2014. The secretory pathway in the filamentous fungus *Trichoderma*, p. 115–121. *In* Gupta, V, Schmoll, M, Herrera-Estrella, A, Upadhyay, R, Druzhinina, I, Tuohy, M (eds.), *Biotechnology and biology of Trichoderma*. Elsevier.
141. Woo PCY, Chong KTK, Tse H, Cai JJ, Lau CCY, Zhou AC, Lau SKP, Yuen K yung. 2006. Genomic and experimental evidence for a potential sexual cycle in the pathogenic thermal dimorphic fungus *Penicillium marneffeii*. *FEBS Lett* 580:3409–3416.
142. Kato E. 1998. Characterization of a serine carboxypeptidase in *Neurospora crassa*, homologous to the KEX1 gene of *S. cerevisiae*. University of Hawaii, Honolulu.
143. Jacob-Wilk D, Turina M, Kazmierczak P, Van Alfen NK. 2009. Silencing of *Kex2* significantly diminishes the virulence of *Cryphonectria parasitica*. *Mol Plant-Microbe Interact* 22:211–221.

144. Wang Z, Lopez-giraldez F, Lehr N, Common R, Trail F, Townsend JP. 2014. Global gene expression and focused knockout analysis reveals genes associated with fungal fruiting body development in *Neurospora crassa*. *Eukaryot Cell* 13:154–169.
145. Xue C, Hsueh YP, Heitman J. 2008. Magnificent seven: Roles of G-protein-coupled receptors in extracellular sensing in fungi. *FEMS Microbiol Rev* 32:1010–1032.
146. Fischer MS, Glass NL. 2019. Communicate and fuse: How filamentous fungi establish and maintain an interconnected mycelial network. *Front Microbiol* 10:1–20.
147. Seibel C, Tisch D, Kubicek CP, Schmoll M. 2012. The role of pheromone receptors for communication and mating in *Hypocrea jecorina* (*Trichoderma reesei*). *Fungal Genet Biol* 49:814–824.
148. Seo JA, Han KH, Yu JH. 2004. The *gprA* and *gprB* genes encode putative G-protein-coupled receptors required for self-fertilization in *Aspergillus nidulans*. *Mol Microbiol* 53:1611–1623.
149. Brown NA, Schrevens S, Van Dijck P, Goldman GH. 2018. Fungal G-protein-coupled receptors: Mediators of pathogenesis and targets for disease control. *Nat Microbiol* 3:402–414.
150. Ivey FD, Hodge PN, Turner GE, Borkovich KA. 2013. The Gai homologue *gna-1* controls multiple differentiation pathways in *Neurospora crassa*. *Mol Biol Cell* 7:1283–1297.
151. Baasiri RA, Lu X, Rowley PS, Turner GE, Borkovich KA. 1997. Overlapping

- functions for two G protein α subunits in *Neurospora crassa*. *Genetics* 147:137–145.
152. Kamerewerd J, Jansson M, Nowrousian M, Pöggeler S, Kück U. 2008. Three α -subunits of heterotrimeric G-proteins and an adenylyl cyclase have distinct roles in fruiting body development in the homothallic fungus *Sordaria macrospora*. *Genetics* 180:191–206.
153. Krystofova S, Borkovich KA. 2005. The heterotrimeric G-protein subunits GNG-1 and GNB-1 form a G $\beta\gamma$ dimer required for normal female fertility, asexual development, and G α protein levels in *Neurospora crassa*. *Eukaryot Cell* 4:365–378.
154. Xu JR. 2000. MAP kinases in fungal pathogens. *Fungal Genet Biol* 31:137–152.
155. Li D, Bobrowicz P, Wilkinson HH, Ebole DJ. 2005. A mitogen-activated protein kinase pathway essential for mating and contributing to vegetative growth in *Neurospora crassa*. *Genetics* 170:1091–1104.
156. Jones C, Greer-Phillips S, Borkovich K. 2007. The response regulator RRG-1 functions upstream of a mitogen-activated protein kinase pathway impacting asexual development, female fertility, osmotic stress, and fungicide resistance in *Neurospora crassa*. *Mol Biol Cell* 18:2123–2136.
157. Fujimura M, Ochiai N, Oshima M, Motoyama T, Ichiishi A, Usami R, Horikoshi K, Yamaguchi I. 2003. Putative homologs of SSK22 MAPKK kinase and PBS2 MAPK kinase of *Saccharomyces cerevisiae* encoded by *os-4* and *os-5* genes for osmotic sensitivity and fungicide resistance in *Neurospora crassa*. *Biosci Biotechnol Biochem* 67:186–191.

158. Frawley D, Bayram Ö. 2020. The pheromone response module, a mitogen-activated protein kinase pathway implicated in the regulation of fungal development, secondary metabolism and pathogenicity. *Fungal Genet Biol* 144:103469.
159. Teichert I, Steffens EK, Schnaß N, Fränzel B, Krisp C, Wolters DA, Kück U. 2014. PRO40 is a scaffold protein of the cell wall integrity pathway, linking the MAP kinase module to the upstream activator protein kinase C. *PLoS Genet* 10:e1004582.
160. Schmidt S, Märker R, Ramšak B, Beier-Rosberger AM, Teichert I, Kück U. 2020. Crosstalk between pheromone signaling and NADPH oxidase complexes coordinates fungal developmental processes. *Front Microbiol* 11.
161. Bloemendal S, Bernhards Y, Bartho K, Dettmann A, Voigt O, Teichert I, Seiler S, Wolters DA, Pöggeler S, Kück U. 2012. A homologue of the human STRIPAK complex controls sexual development in fungi. *Mol Microbiol* 84:310–323.
162. Kück U, Beier AM, Teichert I. 2016. The composition and function of the striatin-interacting phosphatases and kinases (STRIPAK) complex in fungi. *Fungal Genet Biol* 90:31–38.
163. Xiang Q, Rasmussen C, Louise Glass N. 2002. The *ham-2* locus, encoding a putative transmembrane protein, is required for hyphal fusion in *Neurospora crassa*. *Genetics* 160:169–180.
164. Simonin AR, Rasmussen CG, Yang M, Glass NL. 2010. Genes encoding a striatin-like protein (*ham-3*) and a forkhead associated protein (*ham-4*) are required for hyphal fusion in *Neurospora crassa*. *Fungal Genet Biol* 47:855–868.

165. Fu C, Iyer P, Herkal A, Abdullah J, Stout A, Free SJ. 2011. Identification and characterization of genes required for cell-to-cell fusion in *Neurospora crassa*. *Eukaryot Cell* 10:1100–1109.
166. Märker R, Blank-Landeshammer B, Beier-Rosberger A, Sickmann A, Kück U. 2020. Phosphoproteomic analysis of STRIPAK mutants identifies a conserved serine phosphorylation site in PAK kinase CLA4 to be important in fungal sexual development and polarized growth. *Mol Microbiol* 113:1053–1069.
167. Pöggeler S, Kück U. 2004. A WD40 repeat protein regulates fungal cell differentiation and can be replaced functionally by the mammalian homologue striatin. *Eukaryot Cell* 3:232–240.
168. Beier A, Teichert I, Krisp C, Wolters DA, Kück U. 2016. Catalytic subunit 1 of protein phosphatase 2A is a subunit of the STRIPAK complex and governs fungal sexual development. *MBio* 7:1–11.
169. Frey S, Reschka EJ, Pöggeler S. 2015. Germinal center kinases SmKIN3 and SmKIN24 are associated with the *Sordaria macrospora* striatin-interacting phosphatase and kinase (STRIPAK) complex. *PLoS One* 10:1–27.
170. Nordzieke S, Zobel T, Fränzel B, Wolters DA, Kück U, Teichert I. 2015. A fungal sarcolemmal membrane-associated protein (SLMAP) homolog plays a fundamental role in development and localizes to the nuclear envelope, endoplasmic reticulum, and mitochondria. *Eukaryot Cell* 14:345–358.
171. Bernhards Y, Pöggeler S. 2011. The phocein homologue SmMOB3 is essential for vegetative cell fusion and sexual development in the filamentous ascomycete *Sordaria macrospora*. *Curr Genet* 57:133–149.

172. Li L, Xue C, Bruno K, Nishimura M, Xu JR. 2004. Two PAK kinase genes, *CHM1* and *MST20*, have distinct functions in *Magnaporthe grisea*. *Mol Plant-Microbe Interact* 17:547–556.
173. Park G, Servin JA, Turner GE, Altamirano L, Colot H V., Collopy P, Litvinkova L, Li L, Jones CA, Diala FG, Dunlap JC, Borkovich KA. 2011. Global analysis of serine-threonine protein kinase genes in *Neurospora crassa*. *Eukaryot Cell* 10:1553–1564.
174. Wang C, Zhang S, Hou R, Zhao Z, Zheng Q, Xu Q, Zheng D, Wang G, Liu H, Gao X, Ma JW, Kistler HC, Kang Z, Xu JR. 2011. Functional analysis of the kinome of the wheat scab fungus *Fusarium graminearum*. *PLoS Pathog* 7:e1002460.
175. Chinnici JL, Fu C, Caccamise LM, Arnold JW, Free SJ. 2014. *Neurospora crassa* female development requires the PACC and other signal transduction pathways, transcription factors, chromatin remodeling, cell-to-cell fusion, and autophagy. *PLoS One* 9:e110603.
176. Virgilio S, Bertolini MC. 2018. Functional diversity in the pH signaling pathway: An overview of the pathway regulation in *Neurospora crassa*. *Curr Genet* 64:529–534.
177. Lamb TM, Xu W, Diamond A, Mitchell AP. 2001. Alkaline response genes of *Saccharomyces cerevisiae* and their relationship to the RIM101 pathway. *J Biol Chem* 276:1850–1856.
178. Arst HN, Peñalva MA. 2003. pH regulation in *Aspergillus* and parallels with higher eukaryotic regulatory systems. *Trends Genet* 19:224–231.
179. Peñalva MA, Tilburn J, Bignell E, Arst HN. 2008. Ambient pH gene regulation in

- fungi: Making connections. Trends Microbiol 16:291–300.
180. Virgilio S, Cupertino FB, Ambrosio DL, Bertolini MC. 2017. Regulation of the reserve carbohydrate metabolism by alkaline pH and calcium in *Neurospora crassa* reveals a possible cross-regulation of both signaling pathways. BMC Genomics 18:1–15.
 181. Su SSY, Mitchell AP. 1993. Identification of functionally related genes that stimulate early meiotic gene expression in yeast. Genetics 133:67–77.
 182. Li W, Mitchell AP. 1997. Proteolytic activation of Rim1p, a positive regulator of yeast sporulation and invasive growth. Genetics 145:63–73.
 183. Nahlik K, Dumkow M, Bayram Ö, Helmstaedt K, Busch S, Valerius O, Gerke J, Hoppert M, Schwier E, Opitz L, Westermann M, Grond S, Feussner K, Goebel C, Kaefer A, Meinicke P, Feussner I, Braus GH. 2010. The COP9 signalosome mediates transcriptional and metabolic response to hormones, oxidative stress protection and cell wall rearrangement during fungal development. Mol Microbiol 78:964–979.
 184. Busch S, Schwier EU, Nahlik K, Bayram O, Helmstaedt K, Draht OW, Krappmann S, Valerius O, Lipscomb WN, Braus GH. 2007. An eight-subunit COP9 signalosome with an intact JAMM motif is required for fungal fruit body formation. Proc Natl Acad Sci 104:8089–8094.
 185. Busch S, Eckert SE, Krappmann S, Braus GH. 2003. The COP9 signalosome is an essential regulator of development in the filamentous fungus *Aspergillus nidulans*. Mol Microbiol 49:717–730.
 186. Braus GH, Irrniger S, Bayram Ö. 2010. Fungal development and the COP9

- signalosome. *Curr Opin Microbiol* 13:672–676.
187. Christmann M, Schmalzer T, Gordon C, Huang X, Bayram Ö, Schinke J, Stumpf S, Dubiel W, Braus GH. 2013. Control of multicellular development by the physically interacting deneddylases DEN1/DenA and COP9 signalosome. *PLoS Genet* 9:e1003275.
 188. Ronald Morris N, Kirk KE. 2012. TubB α -tubulin is essential for sexual development in *Aspergillus nidulans*. *Mol Biol its Appl to Med Mycol* 97–104.
 189. Krappmann S, Jung N, Medic B, Busch S, Prade RA, Braus GH. 2006. The *Aspergillus nidulans* F-box protein GrrA links SCF activity to meiosis. *Mol Microbiol* 61:76–88.
 190. Shiu PKT, Raju NB. 2001. Meiotic silencing by unpaired DNA. *Cell* 107:905–916.
 191. Shiu PKT, Zickler D, Raju NB, Ruprich-Robert G, Metzenberg RL. 2006. SAD-2 is required for meiotic silencing by unpaired DNA and perinuclear localization of SAD-1 RNA-directed RNA polymerase. *Proc Natl Acad Sci* 103:2243–2248.
 192. Tessé S, Storlazzi A, Kleckner N, Gargano S, Zickler D. 2003. Localization and roles of Ski8p protein in *Sordaria* meiosis and delineation of three mechanistically distinct steps of meiotic homolog juxtaposition. *Proc Natl Acad Sci* 100:12865–12870.
 193. Nolting N, Pöggeler S. 2006. A STE12 homologue of the homothallic ascomycete *Sordaria macrospora* interacts with the MADS box protein MCM1 and is required for ascosporeogenesis. *Mol Microbiol* 62:853–868.
 194. Werner A, Otte K, Stahlhut G, Hanke LM, Pöggeler S. 2021. The glyoxysomal protease LON2 is involved in fruiting-body development, ascosporeogenesis and

- stress resistance in *Sordaria macrospora*. *J Fungi* 7:1–18.
195. Werner A, Herzog B, Voigt O, Valerius O, Braus GH, Pöggeler S. 2019. NBR1 is involved in selective pexophagy in filamentous ascomycetes and can be functionally replaced by a tagged version of its human homolog. *Autophagy* 15:78–97.
 196. Voigt O, Pöggeler S. 2013. Autophagy genes *Smatg8* and *Smatg4* are required for fruiting-body development, vegetative growth and ascospore germination in the filamentous ascomycete *Sordaria macrospora*. *Autophagy* 9:33–49.
 197. Nelson A, Merino ST, Mekenberg RL. 1997. A putative rhamnogalacturonase required for sexual development of *Neurospora crassa*. *Genetics* 146:531–540.
 198. Traeger S, Nowrousian M. 2015. Functional analysis of developmentally regulated genes *chs7* and *sec22* in the ascomycete *Sordaria macrospora*. *G3 Genes, Genomes, Genet* 5:1233–1245.
 199. Nair U, Jotwani A, Geng J, Gammoh N, Richerson D, Yen WL, Griffith J, Nag S, Wang K, Moss T, Baba M, McNew JA, Jiang X, Reggiori F, Melia TJ, Klionsky DJ. 2011. SNARE proteins are required for macroautophagy. *Cell* 146:290–302.
 200. Cao S, He Y, Hao C, Xu Y, Zhang H, Wang C, Liu H, Xu JR. 2017. RNA editing of the *AMD1* gene is important for ascus maturation and ascospore discharge in *Fusarium graminearum*. *Sci Rep* 7:1–11.
 201. Liu H, Li Y, Chen D, Qi Z, Wang Q, Wang J, Jiang C, Xu JR. 2017. A-to-I RNA editing is developmentally regulated and generally adaptive for sexual reproduction in *Neurospora crassa*. *Proc Natl Acad Sci* 114:E7756–E7765.
 202. Glass NL, Metznerberg RL, Raju NB. 1990. Homothallic Sordariaceae from nature:

- The absence of strains containing only the a mating type sequence. *Exp Mycol* 14:274–289.
203. Ferreira AVB, Saupe S, Glass NL. 1996. Transcriptional analysis of the *mt A* idiomorph of *Neurospora crassa* identifies two genes in addition to *mt A-1*. *Mol Gen Genet* 250:767–774.
204. Singh G, Ashby AM. 1998. Cloning of the mating-type loci from *Pyrenopeziza brassicae* reveals the presence of a novel mating type gene within a discomycete *MAT1-2* locus encoding a putative metallothionein-like protein. *Mol Microbiol* 30:799–806.
205. Amselem J, Cuomo CA, van Kan JAL, Viaud M, Benito EP, Couloux A, Coutinho PM, de Vries RP, Dyer PS, Fillinger S, Fournier E, Gout L, Hahn M, Kohn L, Lapalu N, Plummer KM, Pradier JM, Quévillon E, Sharon A, Simon A, Have A, Tudzynski B, Tudzynski P, Wincker P, Andrew M, Anthouard V, Beever RE, Beffa R, Benoit I, Bouzid O, Brault B, Chen Z, Choquer M, Collémare J, Cotton P, Danchin EG, Da Silva C, Gautier A, Giraud C, Giraud T, Gonzalez C, Grossetete S, Güldener U, Henrissat B, Howlett BJ, Kodira C, Kretschmer M, Lappartient A, Leroch M, Levis C, Mauceli E, Neuvéglise C, Oeser B, Pearson M, Poulain J, Poussereau N, Quesneville H, Rasclé C, Schumacher J, Ségurens B, Sexton A, Silva E, Sirven C, Soanes DM, Talbot NJ, Templeton M, Yandava C, Yarden O, Zeng Q, Rollins JA, Lebrun MH, Dickman M. 2011. Genomic analysis of the necrotrophic fungal pathogens *Sclerotinia sclerotiorum* and *Botrytis cinerea*. *PLoS Genet* 7:e1002230.
206. Palmer JM, Kubatova A, Novakova A, Minnis AM, Kolarik M, Lindner DL. 2014.

- Molecular characterization of a heterothallic mating system in *Pseudogymnoascus destructans*, the fungus causing white-nose syndrome of bats. *G3 Genes, Genomes, Genet* 4:1755–1763.
207. Bihon W, Wingfield MJ, Slippers B, Duong TA, Wingfield BD. 2014. *MAT* gene idiomorphs suggest a heterothallic sexual cycle in a predominantly asexual and important pine pathogen. *Fungal Genet Biol* 62:55–61.
208. Pöggeler S, Kück U. 2000. Comparative analysis of the mating-type loci from *Neurospora crassa* and *Sordaria macrospora*: Identification of novel transcribed ORFs. *Mol Gen Genet* 263:292–301.
209. Kanamori M, Kato H, Yasuda N, Koizumi S, Peever TL, Kamakura T, Teraoka T, Arie T. 2007. Novel mating type-dependent transcripts at the mating-type locus in *Magnaporthe oryzae*. *Gene* 403:6–17.
210. Yu J, Sun W, Yu M, Yin X, Meng X, Zhao J, Huang L, Huang L, Liu Y. 2015. Characterization of mating-type loci in rice false smut fungus *Villosiclava virens*. *FEMS Microbiol Lett* 362:1–9.
211. Martin SH, Wingfield BD, Wingfield MJ, Steenkamp ET. 2011. Structure and evolution of the *Fusarium* mating type locus: New insights from the *Gibberella fujikuroi* complex. *Fungal Genet Biol* 48:731–740.
212. Kothe GO, Free SJ. 1998. The isolation and characterization of *nrc-1* and *nrc-2*, two genes encoding protein kinases that control growth and development in *Neurospora crassa*. *Genetics* 149:117–130.
213. Ivey FD, Kays AM, Borkovich KA. 2002. Shared and independent roles for a Gai protein and adenylyl cyclase in regulating development and stress responses in

Neurospora crassa. Eukaryot Cell 1:634–642.

AUTHOR BIOGRAPHIES

Andi Wilson is a postdoctoral researcher at the Forestry & Agricultural Biotechnology Institute at the University of Pretoria, South Africa. She also completed her PhD at this institute under the primary supervision of Prof. BD Wingfield. Andi's primary research focus is on elucidating the underlying genetic mechanisms that control sexual development and mating in ascomycete fungi. Using a combination of comparative genomics and transcriptomics as well as classical genetics and CRISPR-Cas9 gene editing, she has unraveled some of the pathways that allow for unisexual development in filamentous ascomycete fungi.


Markus Wilken completed his Ph.D. (Genetics) at the University of Pretoria. He studied the uni-directional mating type switching system as the reproductive mode for many Ceratocystidaceae species. Thereafter, he completed two post-doctoral fellowships under the SARCHI Chair in Fungal Genomics run by Prof. BD Wingfield, where he used genomics to characterize the mating system of other Ceratocystidaceae species. Currently, Markus is a lecturer in the Department of Biochemistry, Genetics and Microbiology at the University of Pretoria where he runs a research program linked to studying sexual reproduction in plant pathogenic fungi.

Mike Wingfield, the founding director of the Forestry and Agricultural Biotechnology Institute, serves as an advisor to the University of Pretoria Executive. His research focuses on the health of forest trees, having published widely on this topic together with more than a hundred past PhD students. He served a five-year term as President of the

International Union of Forestry Research Organisations (IUFRO) and has been widely recognized for his research including receiving the Kwame Nkrumah Science Award from the African Union and being awarded honorary doctorates from the University of British Columbia (Canada) and North Carolina State University (USA).

Brenda Wingfield has made the study of the global movement and evolution of fungal pathogens, particularly those on trees, her research focus for the past 30 years. Prof. Wingfield holds the DSI-NRF SARCHI chair in Fungal Genomics. She pioneered fungal genomics at the University of Pretoria and was responsible for the first fungal genome to be sequenced in Africa. Her success is reflected in the internationally recognized work of many of her past PhD students. She is currently the Secretary General of the International Society of Plant Pathology and was the vice president of the Academy of Science of South Africa.

FIGURES AND FIGURE LEGENDS



ScEYence Studios
 ASM Journals
 MMBR00020-21
 Dr. Wilson
 Figure: 01

FIG 1 Generalized sexual cycle of heterothallic Pezizomycotina fungi. (1) The cycle begins with an interaction between compatible isolates, (2) with male structures from one of the partners secreting pheromones which are (3) recognized by the female structures from a suitable mating partner. This recognition results in (4) growth of the trichogyne towards the male cells, resulting in physical contact between the two partners, (5) fertilization and (6) the subsequent production of a protoascoma. Within the ascus, (7-12) multiple rounds of meiosis and mitosis produce the mature ascospores which are subsequently (13) released from the (14) mature ascoma. This cycle would be identical in homothallic species, except that an opposite mating partner would not be required to initiate the process. The different ascomatal types are depicted below the sexual cycle- illustrating the morphological diversity that can be observed in the Pezizomyctonia.





FIG 2 Mating-type idiomorphs, *MAT* genes and their distribution in fungi utilizing the various sexual strategies exhibited by Pezizomycotina fungi. Open circles represent nuclei in each cell, while closed circles represent the ascospores produced. Red indicates the presence of genes from the *MAT1-1* idiomorph and blue indicates the presence of genes from the *MAT1-2* idiomorph.




FIG 3 A cladogram of the Pezizomycotina highlighting the lack of studies on sexual reproduction in this group. Classes or Orders shown in red have no described *MAT* genes and/or no published functional studies relating to the sexual process. Orders shown in white have at least one species with described *MAT* genes and/or at least one functional study related to sexual reproduction. Details can be found in the Supplementary information (Table S1). The cladogram was constructed in phyloT using NCBI taxonomic identification numbers, and was edited in iTol v6 (<https://itol.embl.de/>).




FIG 4 The pheromone response pathway in *Neurospora crassa*. The MAT1-1-1 and MAT1-2-1 proteins control the expression of the α - and **a**-factor pheromones, respectively. Recognition of the pheromones by their respective cognate receptors results in the activation of the coupled G-protein and the dissociation of its subunits. These subunits subsequently activate one of the three MAPK cascades, MAK2, that ultimately results in protoascomatal development. Two other MAPK cascades, MAK1 and OS-2, are also important for protoascomatal development but respond to different environmental signals.





FIG 5 Structure and processing of the **a**- and **α**-factor mating pheromones. Both fungal mating pheromones are initially expressed as large pre-proteins which are processed into small, mature pheromone factors. The first step in the processing of the **a**-factor includes the addition of a farnesyl (indicated by the **+**) by RAM1/RAM2. This is followed by cleavage of the terminal AAX amino acids by RCE1/STE24 and the addition of a carboxymethyl group (indicated by the **●**) by STE14. Two final cleavage events, facilitated by STE24 and AXL1, release the mature pheromone. The first step in **α**-factor processing is the removal of the signal peptide (indicated in dark orange). KEX2-, KEX1- and STE13-mediated cleavage releases numerous copies of the mature pheromone.






FIG 6 Important protein complexes and pathways involved in protoascomatal maturation and ascomatal development in various filamentous fungi. A) The STRIPAK is important for sexual development in the homothallic *Sordaria macrospora*. B) The PACC pathway plays an essential role in protoascomatal development in the heterothallic *Neurospora crassa*. C) The COP9 signalosome is important for sexual development and other developmental processes in the homothallic *Aspergillus nidulans*. [Adapted from (161, 174, 185)]. Protein names and/or abbreviations: GPI1 (glycosylphosphatidylinositol-anchored protein), MOB3 (monopolar spindle one-binder protein), PP2Ac and PP2Aa (protein phosphatase subunits), KIN3 and KIN24 (kinases 3 and 24), CLA4 (from the p21-activated kinase family), PALA, PALB, PALC, PALF & PALH (various proteins from the PAL/RIM pathway), PACC (terminal transcription factor of the PACC pathway), CSN1-8 (COP9 signalosome proteins 1-8).

TABLES

TABLE 1 *MAT* genes described from the Pezizomycotina

Gene name	Representative taxa	Representative protein NCBI accession number	Conserved protein domain	References
Primary <i>MAT</i> genes				
<i>MAT1-1-1</i>	<i>Neurospora crassa</i>	ESA43845	PF04769 – MAT α1 HMG-box	(74, 202)
<i>MAT1-2-1</i>	<i>N. crassa</i>	AAA33598	PF00505 - HMG box	(55, 75)
Secondary <i>MAT</i> genes				
<i>MAT1-1-2</i>	<i>N. crassa</i>	EAA35087	PF17043 – MAT1-1-2 domain of unknown function	(203)
<i>MAT1-1-3</i>	<i>N. crassa</i>	EAA35088	PF00505 – HMG box	(204)
<i>MAT1-1-4</i>	<i>Pyrenopeziza brassicae</i>	CAA06845	None	(204)
<i>MAT1-1-5</i>	<i>Botrytis cinerea</i>	AHX22633	None	(205)
<i>MAT1-1-6</i>	<i>Pseudogymnoascus destructans</i>	AIG95729	None	(206)
<i>MAT1-1-7</i>	<i>Coccidioides immitis</i>	ABS19617	None	(67)
<i>MAT1-1-8</i>	<i>Diplodia sapinea</i>	AHA91691	None	(207)
<i>MAT1-1-9^a</i>	-	-	-	-
<i>MAT1-1-10</i>	<i>Morchella importuna</i>	MG681026	None	(68)
<i>MAT1-1-11</i>	<i>M. importuna</i>	MG681027	None	(68)
<i>MAT1-1-12</i>	<i>Teratosphaeria zuluensis</i>	QEJ80698	None	(69)
<i>MAT1-2-2</i>	<i>N. crassa</i>	^b	None	(85, 208)
<i>MAT1-2-3</i>	<i>Co. immitis</i>	KMP00264	PF11051 – Mannosyl trans3	(67)
<i>MAT1-2-4</i>	<i>Aspergillus fumigatus</i>	XP754990	None	(63)
<i>MAT1-2-5</i>	<i>D. sapinea</i>	AHA91682	None	(207)
<i>MAT1-2-6</i>	<i>Magnaporthe grisea</i>	BAE66607	PF00505 – HMG box	(209)
<i>MAT1-2-7</i>	<i>Huntiella omanensis</i>	AOY41711	None	(71)
<i>MAT1-2-8</i>	<i>Villosiclava virens</i>	AKE48512	None	(210)
<i>MAT1-2-9</i>	<i>Fusarium fujikuroi</i>	AEP03799	None	(211)
<i>MAT1-2-10</i>	<i>B. cinerea</i>	CDF43998	None	(205)

MAT1-2-11	<i>Ps. destructans</i>	AIG95713	None	(206)
MAT1-2-12	<i>T. zuluensis</i>	QEJ80704	None	(69)
MAT1-2-13	<i>Ca. pauciramosa</i>	QKY89075	None	(70)
MAT1-2-14	<i>Letharia columbiana</i>	QEL51134	None	(81)
MAT1-2-15	<i>Colletotrichum fructicola</i>	XM032032861	None	(80)

^a A recent paper synonymized *MAT1-1-7* and *MAT1-1-9*, leaving the *MAT1-1-9* gene name vacant (81). See text for detail.

^b The *N. crassa* *MAT1-2-2* gene was predicted, but never annotated in the Genbank database. See (56) for detail.

TABLE 2 Comparison between the functions of proteins involved in the pheromone response pathway and MAPK cascades in heterothallic and homothallic species

Protein	<i>Neurospora crassa</i> (Heterothallic)		<i>Sordaria macrospora</i> (Homothallic)		
	Gene Name	Mutant Phenotypes	Gene Name	Mutant Phenotypes	
α-factor pheromone	<i>ccg4</i>	Male sterility* in MAT1-1 partner (106)	<i>ppg1</i>	No sexual defects (128)	<i>Δppg1/Δpre1</i> : Greatly reduced ^b ascomatal production (128)
a-factor pheromone	<i>mfa-1</i>	Male sterility in MAT1-2 partner (106) Aberrant female development (108)	<i>ppg2</i>	No sexual defects (128)	<i>Δppg1/Δppg2</i> : Reduced ^a ascomatal production (128) <i>Δppg2/Δpre2</i> : Greatly reduced ascomatal production (128)
α-factor receptor	<i>pre2</i>	No protoascomatal maturation in MAT1-2 partner (24)	<i>pre2</i>	No sexual defects (128)	<i>Δpre1/Δpre2</i> : No protoascomatal maturation (128)
a-factor receptor	<i>pre1</i>	No protoascomatal maturation in MAT1-1 partner (107)	<i>pre1</i>	No sexual defects (128)	
Gα subunit	<i>gna1</i>	Unusually shaped protoascomata that do not grow towards spermatia (150)	<i>gsa1</i>	Delayed and reduced ascomatal production (152)	<i>Δgsa1/Δgsa2</i> & <i>Δgsa1/Δgsa3</i> : Greatly deduced protoascomatal production and no protoascomatal maturation (152)
	<i>gna2</i>	No sexual defects (151)	<i>gsa2</i> <i>gsa3</i>	No sexual defects (152)	Greatly reduced ascospore germination (152)
Gβ subunit	<i>gnb1</i>	No protoascomatal maturation (111,153)	-	-	
Gγ subunit	<i>gnγ1</i>	No protoascomatal maturation [†] (153)	-	-	
MAPKKK	<i>nrc1</i>	No protoascomatal production [‡] (212)	<i>mik2</i>	No protoascomatal maturation (160)	
MAPKK	-	-	<i>mek2</i>	No protoascomatal maturation (160)	

MAPK	<i>mak2</i>	No protoascomatal production (155)	<i>mak2</i>	No protoascomatal maturation (160)
Transcriptional Regulator	<i>pp-1</i>	No protoascomatal production (155)	<i>ste12</i>	Reduced asci production, fragile walls of asci and ascospores (193)
MAPKKK	<i>mik1</i>	No protoascomatal production (110)	<i>mik1</i>	No protoascomatal maturation (159)
MAPKK	<i>mek1</i>	No protoascomatal production (110)	<i>mek1</i>	No protoascomatal maturation (159)
MAPK	<i>mak1</i>	No protoascomatal production (110)	<i>mak1</i>	No protoascomatal maturation (159)
	<i>Rrg-1</i>	No protoascomatal production (156)	-	-
MAPKKK	<i>os-4</i>	No protoascomatal production (157)	-	-
MAPKK	<i>os-5</i>	No protoascomatal production (157)	-	-
MAPK	<i>os-2</i>	No protoascomatal production (156)	-	-
Adenylyl Cyclase	<i>cr-1</i>	Delayed ascomatal development (213)	<i>sac1</i>	Reduced ascomatal size and ascospore germination (152)

* Male sterility indicates that an isolate is unable to act as the male partners

† No protoascomatal maturation indicates that protoascomata are produced, but do not mature into ascomata

‡ No protoascomatal production indicates that protoascomata are not formed at all

Wild type (proto)ascomatal production > ^a Reduced (proto)ascomatal production > ^b Greatly reduced (proto)ascomatal production