

Nadeeshani Karannagoda<sup>\*1,2</sup>, Antanas Spokevicius<sup>1</sup>, Steven Hussey<sup>3</sup>, Hua Cassan-Wang<sup>4</sup>, Jacqueline Grima-Pettenati<sup>4</sup>, Gerd Bossinger<sup>1</sup>


<sup>1</sup>School of Ecosystem and Forest Sciences, The University of Melbourne, Creswick, Victoria 3363, Australia

<sup>2</sup>Agriculture Victoria, AgriBio, Centre for AgriBioscience, Bundoora, Victoria 3083, Australia

<sup>3</sup>Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa, 0002

<sup>4</sup>Laboratoire de Recherche en Sciences Végétales, Université de Toulouse III, CNRS, UPS, UMR 5546, 24 Chemin de Borde Rouge, 31320 Castanet-Tolosan, France

\*Corresponding author: nadeeshani.karannagoda@agriculture.vic.gov.au



Online Resource 3

Comparison of average changes in (a) cell wall thickness, (b) cell wall area, (c) lumen area and (d) cross-sectional area of *Populus alba* xylem fibres transformed with IAA13+. The black dot represents the mean value. Error bars represent 95% confidence intervals.