

# Groups with a given number of nonpower subgroups

C. S. Anabanti<sup>1</sup> and S. B. Hart<sup>2</sup>

<sup>1</sup>chimere.anabanti@up.ac.za; Dept. of Mathematics and Applied Mathematics, University of Pretoria

<sup>2</sup>s.hart@bbk.ac.uk; Dept. of Economics, Mathematics and Statistics, Birkbeck, University of London

## Abstract

It is well-known that no group has either exactly 1 or exactly 2 nonpower subgroups. In this paper, we obtain a classification of groups containing exactly 3 nonpower subgroups. Moreover, we show that there is a unique finite group with exactly 4 nonpower subgroups. Finally, we show that given any integer  $k$  greater than 4, there are infinitely many groups with exactly  $k$  nonpower subgroups.

## 1 Introduction

A subgroup  $H$  of a group  $G$  is called a *power subgroup* of  $G$  if there exists a non-negative integer  $m$  such that  $H = \langle g^m : g \in G \rangle$ . Any subgroup of  $G$  which is not a power subgroup is called a *nonpower subgroup* of  $G$ . Zhou et al.[3] proved that cyclic groups have no nonpower subgroups, and infinite noncyclic groups have an infinite number of nonpower subgroups. They showed further that no group has either exactly 1 or exactly 2 nonpower subgroups, and then asked: for each integer  $k$  greater than 2, does there exist at least one group possessing exactly  $k$  nonpower subgroups? This question was recently answered positively in [1], where it was also proved that for any integer  $k$  greater than 4 and composite, there are infinitely many groups with exactly  $k$  nonpower subgroups.

Let  $p$  be an odd prime. For each positive integer  $n$ , we define the group  $G_{n,p}$  as follows:

$$G_{n,p} := \langle x, y : x^{2^n} = 1 = y^p, yx = xy^{-1} \rangle.$$

We note that  $G_{1,p}$  is the dihedral group of order  $2p$ , and  $G_{2,p}$  is the generalized quaternion group of order  $4p$  (we obtain its usual presentation  $\langle a, b : a^{2p} = 1, b^2 = a^p, ba = a^{-1}b \rangle$  by setting  $a = x^2y$  and  $b = x$ ). More generally, for any positive integer  $n$ ,  $G_{n,p}$  is the semidirect product  $C_p \rtimes C_{2^n}$ , and has order  $2^n p$ . We may now state our first result.

**Theorem 1.** *There are infinitely many groups with an odd prime number of nonpower subgroups. In particular, for any odd prime  $p$  and each positive integer  $n$ , the group  $G_{n,p}$  has exactly  $p$  nonpower subgroups.*

---

*Key words and phrases:* Counting subgroups, nonpower subgroups, finite groups  
*2010 Mathematics Subject Classification:* 20D25, 20D60, 20E34.

Theorem 1, combined with the fact that for composite  $k$  greater than 4 there are infinitely many groups with  $k$  nonpower subgroups [1, Theorem 5], gives the following immediate corollary.

**Corollary 2.** *Let  $k$  be an integer greater than 4. Then there are infinitely many groups with exactly  $k$  nonpower subgroups.*

The only unresolved cases are therefore  $k = 3$  and  $k = 4$ . Our second main result deals with these cases.

**Theorem 3.** (a) *A group  $G$  contains exactly three nonpower subgroups if and only if  $G$  is isomorphic to one of  $C_2 \times C_2$ ,  $Q_8$  or  $G_{n,3}$  for  $n \in \mathbb{Z}^+$ .*

(b) *Up to isomorphism,  $C_3 \times C_3$  is the only group containing exactly four nonpower subgroups.*

For the rest of this section, we recall some preliminaries. We note that each power subgroup is characteristic and hence normal in  $G$ . Following [1], we write  $s(G)$  for the number of subgroups in a group  $G$ ,  $ps(G)$  for the number of power subgroups of  $G$  and  $nps(G)$  for the number of nonpower subgroups of  $G$ .

**Lemma 4.** [1, Lemma 3] *If  $A$  and  $B$  are finite groups such that  $|A|$  and  $|B|$  are coprime, then*

$$nps(A \times B) = nps(A)s(B) + ps(A)nps(B).$$

We denote by  $\Phi(G)$  the Frattini subgroup of  $G$ , that is, the intersection of the maximal subgroups of  $G$ . It is a characteristic subgroup of  $G$ .

**Theorem 5** (Burnside's Basis Theorem). *Let  $G$  be a  $p$ -group and suppose  $[G : \Phi(G)] = p^d$ .*

(a)  *$G/\Phi(G)$  is elementary abelian of order  $p^d$ . Moreover, if  $N \trianglelefteq G$  and  $G/N$  is elementary abelian, then  $\Phi(G) \leq N$ .*

(b) *Every minimal system of generators of  $G$  contains exactly  $d$  elements.*

(c)  *$\Phi(G) = G^p G'$ . In particular, if  $p = 2$ , then  $\Phi(G) = G^2$ .*

**Lemma 6** ([2] Theorem 1.10(a)). *Let  $G$  be a non-cyclic  $p$ -group, where  $p > 2$ . Then the number of subgroups of order  $p$  in  $G$  is congruent to  $1 + p$  modulo  $p^2$ .*

**Remark.** *It is well-known that the only 2-groups with a unique involution are cyclic or generalised quaternion.*

## 2 Proof of main results

We begin with a proof of Theorem 1.

*Proof of Theorem 1.* Let  $p$  be an odd prime. Our goal is to show that for any positive integer  $n$ , and any odd prime  $p$ , the group  $G_{n,p} = \langle x, y : x^{2^n} = 1 = y^p, yx = xy^{-1} \rangle$  contains exactly  $p$  nonpower subgroups. We have that  $|G_{n,p}| = 2^n p$ . We first obtain a count on the number of subgroups in  $G_{n,p}$ . Since the Sylow 2-subgroup  $\langle x \rangle$  is not a normal subgroup, the number of

Sylow 2-subgroups of  $G_{n,p}$  must be  $p$ . On the other hand, since  $y^x = y^{-1}$ , there is a unique normal Sylow  $p$ -subgroup, namely the cyclic subgroup  $\langle y \rangle$  of order  $p$ . Since  $x^2$  is central in  $G_{n,p}$  and each Sylow 2-subgroup of  $G_{n,p}$  is cyclic, there is a unique subgroup of order  $2^k$  (for each  $k \in \{0, \dots, n-1\}$ ) and a unique subgroup of order  $2^k p$  (for each  $k \in \{1, \dots, n\}$ ). Along with the  $p$  subgroups of order  $2^n$ , we see that  $s(G_{n,p}) = 2n + p + 1$ . As the subgroups of order  $2^n$  are not normal, we know immediately that they are nonpower subgroups. Hence  $nps(G_{n,p}) \geq p$ . We now show that any subgroup of  $G_{n,p}$  that is not a Sylow 2-subgroup of  $G_{n,p}$  is a power subgroup of  $G_{n,p}$ . First, the unique subgroup of order  $p$  is  $G_{n,p}^{2^n}$ . Secondly, for each  $k \in \{0, \dots, n-1\}$ , the subgroup of order  $2^k$  is  $G_{n,p}^{2^{n-k}p}$ . Finally, for each  $k \in \{1, \dots, n\}$ , the subgroup of order  $2^k p$  is  $G_{n,p}^{2^{n-k}}$ . Therefore,  $ps(G_{n,p}) = 2n + 1$ ; whence  $nps(G_{n,p}) = p$ .  $\square$

We now move onto the proof of Theorem 3. Let  $G$  be a finite noncyclic group. Then  $G$  falls into one of the following three categories: (i) a noncyclic  $p$ -group; (ii) a noncyclic nilpotent group that is not a  $p$ -group; (iii) a non-nilpotent group. For each of these cases above, we classify all the finite groups with exactly 3 or 4 nonpower subgroups.

**Proposition 7.** *Let  $G$  be a finite noncyclic  $p$ -group. Then  $nps(G) = 3$  if and only if  $G$  is  $C_2 \times C_2$  or  $Q_8$ , and  $nps(G) = 4$  if and only if  $G$  is  $C_3 \times C_3$ .*

*Proof.* Let  $G$  be noncyclic of order  $p^n$ . It was shown in [3] that if  $N \triangleleft G$  and  $A/N$  is a nonpower subgroup of  $G/N$ , then  $A$  is a nonpower subgroup of  $G$ . Suppose  $G$  has exactly  $k$  nonpower subgroups, where  $k \in \{3, 4\}$ . Now,  $G/\Phi(G) \cong C_p \times \dots \times C_p$  ( $d$ -times), and  $d \geq 2$  as  $G$  is not cyclic. The  $\frac{p^d-1}{p-1}$  cyclic subgroups of order  $p$  in  $C_p^d$  are nonpower subgroups. Thus  $G/\Phi(G)$ , and hence  $G$ , has at least  $1 + p + \dots + p^{d-1}$  nonpower subgroups. Hence,  $d = 2$ , either  $p = 2$  or  $p = 3$ , and  $G$  has  $p + 1$  maximal subgroups that are nonpower subgroups.

The power subgroups of  $G$  are  $G^1 = G, G^p, G^{p^2}, \dots, G^{p^m}$ , where  $p^m$  is the exponent of  $G$ . There are thus at most  $m + 1$  distinct power subgroups. Since  $G$  is not cyclic, this means  $m < n$ ; so  $ps(G) \leq n$ .

What about  $s(G)$ ? There is at least one subgroup of order  $p^i$  for  $0 \leq i \leq n$  (just take any composition series). This gives at least  $n + 1$  subgroups. But there are  $p + 1$  maximal subgroups (of order  $p^{n-1}$ ) arising from the  $p + 1$  nontrivial proper subgroups of  $G/\Phi(G)$ . Thus  $s(G) \geq n + p + 1$ .

Suppose  $p = 2$ . If  $G$  is not generalised quaternion (and by assumption  $G$  is not cyclic), then  $G$  has at least 3 involutions, and hence at least 3 subgroups of order 2. So, if  $n > 2$ , then  $s(G) \geq n + 5$ , meaning that  $nps(G) \geq 5$ , a contradiction. Thus, either  $G$  is generalised quaternion or  $n = 2$ , which means  $G \cong C_2 \times C_2$ , and in this case  $nps(G) = 3$ . If  $G$  is generalised quaternion, then  $G$  has  $2^{n-1} + 2$  elements of order 4, resulting in  $2^{n-2} + 1$  subgroups of order 4. If  $n > 3$ , we get that  $s(G) \geq n + 1 + 2^{n-2} \geq n + 5$ . Again, this means that  $nps(G) \geq 5$ . Thus,  $n = 3$ , and then  $G \cong Q_8$ . Again,  $nps(Q_8) = 3$ .

The remaining case is  $p = 3$ . By Lemma 6, there are at least four subgroups of order 3 in  $G$ . If  $n > 2$ , then these are distinct from the four maximal subgroups, and so we get  $s(G) \geq n + 7$ . This forces  $nps(G) \geq 7$ , a contradiction. The only possibility is that  $n = 2$ . A quick check shows that  $nps(C_3 \times C_3) = 4$ .

Thus,  $nps(G) = 3$  if and only if  $G$  is  $C_2 \times C_2$  or  $Q_8$ , and  $nps(G) = 4$  if and only if  $G$  is  $C_3 \times C_3$ .  $\square$

**Lemma 8.** *Let  $G$  be a finite noncyclic nilpotent group. If  $G$  is not a  $p$ -group, then  $nps(G) \geq 6$ .*

*Proof.* Recall that a finite group is nilpotent if and only if it is the direct product of its Sylow subgroups, each of which is normal. Since  $G$  is noncyclic, at least one of these Sylow subgroups is noncyclic. Let  $p_1, \dots, p_r$  be the primes dividing  $|G|$ , and let  $P_1, \dots, P_r$  be the respective Sylow subgroups. Assume, without loss of generality, that  $P_1$  is noncyclic. Write  $Q = P_2 \times \dots \times P_r$ ; so  $G \cong P_1 \times Q$ . Since  $G$  is not a  $p$ -group, we have that  $Q \neq \{1\}$ . By Lemma 4 therefore,

$$nps(G) = nps(P_1)s(Q) + ps(P_1)nps(Q) \geq nps(P_1)s(Q).$$

As  $Q \neq \{1\}$ , we have that  $s(Q) \geq 2$ . As  $P_1$  is not cyclic,  $nps(P_1) \geq 3$ . Hence  $nps(G) \geq 6$ .  $\square$

**Lemma 9.** *If  $G$  is a finite non-nilpotent group such that  $nps(G) \in \{3, 4\}$ , then  $nps(G) = 3$  and  $G \cong G_{n,3} = \langle x, y : x^{2^n} = 1 = y^3, yx = xy^{-1} \rangle$ , for some positive integer  $n$ .*

*Proof.* Suppose  $G$  is finite, non-nilpotent and  $nps(G) = k \in \{3, 4\}$ . If  $G$  had a unique Sylow  $p$ -subgroup for each  $p$  dividing  $|G|$ , then  $G$  would be nilpotent. So there is at least one such  $p$  for which  $G$  has more than one Sylow  $p$ -subgroup. For any such  $p$ , the number,  $n_p$ , of Sylow  $p$ -subgroups is congruent to 1 mod  $p$ . So  $n_p \geq p + 1$ . These groups are not normal, so are not power subgroups. Therefore, as  $nps(G) \in \{3, 4\}$ , we have that either  $p = 2$  and  $n_2 = 3$ , or  $p = 3$  and  $n_3 = 4$ . For all other primes  $q$  dividing  $|G|$ , there must be a unique Sylow  $q$ -subgroup. If any subgroup of  $G$ , other than the Sylow  $p$ -subgroups, were non-normal, then it and its conjugates could not be power subgroups. Thus there would be at least two further nonpower subgroups, forcing  $nps(G) \geq 5$ , a contradiction. Therefore, every subgroup of  $G$ , other than the Sylow  $p$ -subgroups, is normal.

Let  $P$  be one of the Sylow  $p$ -subgroups. Let  $q_1, \dots, q_r$  be the primes other than  $p$  dividing  $|G|$ . Let  $Q_1, \dots, Q_r$  be the corresponding normal Sylow subgroups. Each  $Q_i$  is normal and the  $Q_i$  intersect trivially. Therefore, defining  $H = Q_1 Q_2 \dots Q_r$ , we have that  $H \cong Q_1 \times Q_2 \times \dots \times Q_r$  is a normal subgroup of  $G$ , with  $G = PH$ . Now,  $P \trianglelefteq N_G(P)$ , and setting  $K = H \cap N_G(P)$ , we have that  $K \trianglelefteq G$  (because certainly  $K$  is not a Sylow  $p$ -subgroup). But  $P$  is normal in  $N_G(P) = PK$ ; so  $N_G(P) \cong P \times K$ . Let  $h \in H - N_G(P)$ . Then  $(PK)^h = P^h K \neq PK$ . This means that  $PK$  is not normal in  $G$ ; a contradiction unless  $K = \{1\}$ . Therefore,  $K = \{1\}$ , and  $P = N_G(P)$ . In particular,  $n_p = |G : P| = |H|$ .

Suppose first that  $p = 3$ . Then  $|H| = 4$ . If  $H \cong C_2 \times C_2$ , then each of its cyclic subgroups would be normal, and hence the involutions they contain would be central. But that would imply that  $P$  is normal in  $G$ , a contradiction. Therefore  $H \cong C_4$ . Let  $z$  be a generator of  $H$ . We have  $H \leq C_G(z) \leq G$ . Thus,  $|z^G| = 3^i$  for some  $i$  with  $0 \leq i \leq n$ . But  $z^G \subseteq \{z, z^{-1}\}$ . The only possibility is that  $z^G = \{z\}$ , and  $z$  is central in  $G$ . Again, this implies that  $P$  is normal in  $G$ , a contradiction. Therefore,  $p \neq 3$ .

The remaining case is when  $p = 2$ . In this case,  $H \cong C_3$ . Let  $A_1, A_2$ , and  $A_3$  be the three Sylow 2-subgroups. Every proper subgroup of  $P$  is not one of  $A_1, A_2$  and  $A_3$ , so is normal in  $G$  and hence contained in all of  $A_1, A_2$  and  $A_3$ . If  $P$  were not cyclic, then each of its generators would generate a proper cyclic subgroup, and would hence be contained in  $A_1, A_2$  and  $A_3$ . This implies  $P \leq A_1 \cap A_2 \cap A_3$ ; a contradiction. Therefore,  $P$  is cyclic of order  $2^n$ . Write  $P = \langle x \rangle$  and  $H = \langle y \rangle$ . Certainly,  $y^x \neq y$ ; so the only possibility is that  $y^x = y^{-1}$ . Therefore,

$$G = \langle x, y : x^{2^n} = 1, y^3 = 1, yx = xy^{-1} \rangle$$

for some integer  $n \geq 1$ . That is,  $G \cong G_{n,3}$ . By Theorem 1, we have  $nps(G) = 3$ .  $\square$

Theorem 3 follows immediately from Proposition 7, Lemma 8 and Lemma 9.

## References

- [1] C. S. Anabanti, A. B. Aroh, S. B. Hart and A. R. Oodo, *A question of Zhou, Shi and Duan on nonpower subgroups of finite groups*, *Quaestiones Mathematicae* (2021), DOI: 10.2989/16073606.2021.1924891.
- [2] Y. Berkovich. *Groups of Prime Power Order*, Volume 1. De Gruyter Expositions in Mathematics **46** (2008).
- [3] W. Zhou, W. Shi and Z. Duan, *A new criterion for finite noncyclic groups*, *Communications in Algebra*, **34** (2006), 4453–4457.