

Supplementary Table 3. Hematology, biochemistry and hemostatic variables in healthy unaffected dogs and the three dogs in which a macrothrombi was identified.

Variable	Healthy unaffected dogs*	Case 1	Case 2	Case 3
Complete blood count				
Hemoglobin (g/dL)	17.7 (17.0, 18.5)	16.6	13.2	7.0
Red blood cell count ($\times 10^6/\text{mm}^3$)	7.82 (7.16, 8.5)	6.95	5.49	2.55
Hematocrit (%)	0.54 (0.51, 0.56)	49	36	20
Mean corpuscular volume (fL)	68.6 (66.2, 72.1)	70.5	65.0	79.5
Mean corpuscular hemoglobin concentration (g/dL)	33.4 (32.8, 33.5)	33.8	36.9	34.3
Red cell distribution width (%)	13.6 (13.1, 13.9)	14.2	16.4	21.5
White cell count ($\times 10^3/\text{mm}^3$)	8.27 (7.24, 9.80)	23.95	38.56	22.69
Neutrophils ($\times 10^3/\text{mm}^3$)	6.45 (1.84)	19.16	32.0	31.11
Band neutrophils ($\times 10^3/\text{mm}^3$)	0 (0, 0)	1.2	1.93	0.7
Lymphocytes ($\times 10^3/\text{mm}^3$)	1.34 (0.53)	2.4	1.54	1.75
Monocytes ($\times 10^3/\text{mm}^3$)	0.41 (0.30, 0.49)	1.2	3.08	1.4
Eosinophil ($\times 10^3/\text{mm}^3$)	0.38 (0.30, 0.55)	0	0	0
Basophil ($\times 10^3/\text{mm}^3$)	0	0	0	0
Platelets ($\times 10^3/\text{mm}^3$)	261 (222, 300)	343	149	61
Biochemistry				
C-reactive protein ($\mu\text{g/L}$)	<10 (<10, <10)	287	315	74
Serum amyloid A ($\mu\text{g/L}$)	<2 (<2, <2)	1697	1597	133
Hemostasis				
TEG R time (min)	3.53 (1.15)	4.9	2.3	5.6
TEG K (min)	1.8 (1.5, 2.0)	1.5	1.2	16.1
TEG angle (degrees)	65.2 (62.1, 69.2)	69.2	74.4	29.7

TEG MA (mm)	59.7 (58.0, 62.5)	73.6	79.7	23.4
TEG G (dyn/cm)	7.4 (6.9, 8.4)	14.0	19.7	1.5
TEG lysis 30 (%)	0 (0, 0)	0.2	0	0
TEG lysis 60 (%)	1.10 (0.30, 1.95)	2.1	0.7	0
Fibrinogen (mg/dL)	324 (261, 368)	113	136	102
Factor X (%)	100 (16.7)	52.6	67.4	31.0
Factor VII (%)	100 (32.2)	45.4	69.1	21.8
D-dimer (ng/mL)	287 (200, 424)	2485	651	6000
Antithrombin (%)	107 (10.08)	80.4	68.5	59.5

*Presented as the mean (standard deviation) for normally distributed data and the median (interquartile range) for non-normal data.