

**Supplemental table 3:** The frequency of image defined risk factors related to risk stratification

<b>All risk stratifications</b>				
		<b>N (%)</b>		
		<b>IDRFs</b>	<b>No IDRFs</b>	<b>Total</b>
IDRFs according to risk stratification	Low risk	15 (41.7)	21 (58.3)	36
	Intermediate risk	13 (46.4)	15 (53.6)	28
	High Risk	78 (38.2)	126 (61.8)	204
	Unknown	0 (0.0)	3 (100)	3
	<b>Total</b>	106 (39.1)	165 (60.9)	271
<b>High-risk neuroblastoma</b>				
		<b>N (%)</b>		
General number of IDRFs	No IDRFs	126/204 (61.8)		
	IDRFs	69/204 (33.8)		
	VE/MBC/OI/AO/SE	44/69 (63.8)		
	Other IDRFs	25/69 (36.2)		
	>1	4/69 (5.8)		
	VE/MBC/OI/AO/SE	4/4 (100.0)		
	Pleural effusions or ascites	10/204 (4.9)		
	<b>Total</b>	204*		
IDRFs per category	Vascular encasement	4/78 (5.1)		
	Multiple body compartments	6/78 (7.7)		
	Organ infiltration	6/78 (7.7)		
	Airway obstruction	9/78 (11.5)		
	Intraspinal tumour extension	23/78 (29.5)		
	Other IDRFs	25/78 (32.1)		
	Pleural effusions and ascites	10/78 (12.8)		
	IDRF + pleural	1/10 (10.0)		
	<b>Total</b>	78*		

\* One IDRF had a pleural effusion or ascites – which accounts for the percentages adding up to more than 100%

**Abbreviations:** AO - airway obstruction; IDRF – image defined risk factor; MBC - multiple body compartments; OI - Organ infiltration; SE - intraspinal tumour extension; VE - vascular encasement