

MLG42					4.2					
MLG43		33.3								
MLG44		2.2								
MLG45		4.4		4.2						
MLG46		4.4		4.2						
MLG47		2.2								
MLG48								6.9		
MLG49								3.4		
MLG50								3.4		
MLG51						38.5		37.9		
MLG52					4.2					
MLG53						7.7				
MLG54								3.4	11.1	
MLG55					4.2					
MLG56									2.8	
MLG57		2.2								
MLG58		2.2		16.7						
MLG59				4.2						
MLG60				12.5						
MLG61									2.8	
MLG62									2.8	
MLG63								6.9		
MLG64									19.4	
MLG65									11.1	
MLG66					8.3	10.0			2.8	
MLG67									2.8	
MLG68		24.4		37.5				3.4	2.8	
No. of isolates/ collection	72	45	18	24	24	10	13	29	36	25
No. of genotypes/ collection	5	14	4	8	13	7	4	15	15	7

Isolates were obtained from pitch canker diseased trees of *Pinus greggii* at GRE10 (Demagtenberg, KwaZulu-Natal), GRE12 (KwaZulu-Natal, Karkloof), GRE5 (KwaZulu-Natal, Karkloof), and *Pinus patula* at PAT12+18 (Limpopo, Soutpansberg), PAT4+6 (Limpopo, Soutpansberg), PAT3 (Limpopo, Soutpansberg). *Pinus patula* establishment disease trees at PATrc3 (Carolina, Mpumalanga) and seedlings at PATnii (Ngondwana, Mpumalanga), PATniii (Sabie, Mpumalanga) and PATni (Tzaneen, Limpopo)