

Diarrhoeal diseases in Soweto, South Africa, 2020: a cross-sectional community survey

Authors and affiliations

Siobhan L. Johnstone^{1,2}, Nicola A. Page^{1,3}, Juno Thomas¹, Shabir A. Madhi⁴, Portia Mutevedzi⁴, Nellie Myburgh⁴, Carlos Herrera⁴, Michelle J. Groome⁴

¹Center for Enteric Diseases, National Institute for Communicable Diseases, Johannesburg, South Africa; ²School of Public Health, Faculty of Health Science, University of the Witwatersrand, Johannesburg, South Africa; ³Department of Medical Virology, Faculty of Health Sciences, University of Pretoria, Private Bag X323, Arcadia, 0007; ⁴South African Medical Research Council: Vaccines and Infectious Diseases Analytics Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Supplementary Tables and Figures




Figure S1: Health seeking for reported diarrhoeal episodes

Table S1: Factors associated with ORS knowledge

	Some knowledge on ORS ^a N = 192	No knowledge on ORS N = 182	p-value
Gender			
Male (%)	29 (15.8)	89 (51.1)	
Female (%)	154 (84.2)	85 (48.9)	<0.001
Age in years (mean; IQR)			
	46.4 (44.1 – 48.7)	44.5 (41.8 - 47.3)	0.300
Children less than 5 years old in the house			
Yes	77 (40.1)	50 (27.5)	
No	115 (59.9)	132 (72.5)	0.010
Children between 5 and 15 years old in the house			
Yes	104 (54.2)	70 (38.5)	
No	88 (45.8)	112 (61.5)	0.002
Wealth Index			
	84.7 (83.4 – 85.9)	86.1 (84.7 – 87.5)	0.132

^a Some knowledge on ORS defined as knowing the recipe or being able to name ingredients.