



REFERENCES

- ACOCKS, J.P.H. 1988.** Veld types of South Africa. *Memoirs of the Botanical Society of South Africa*. No. 57.
- AL-HOUTY, W. & AL-MUSALAM, F. 1997.** Dung preference of the dung beetle *Scarabaeus cristatus* F (Coleoptera – Scarabaeidae) from Kuwait. *Journal of Arid Environments*. 35: 511-516.
- AOAC. 1984.** *Official Methods of Analysis*. (14th ed.) Association of Official Analytical Chemists Inc. Arlington, Virginia, USA.
- ATKINSON, W.D. & SHORROCKS, B. 1981.** Competition on a divided and ephemeral resource: a simulation model. *Journal of Animal Ecology*. 50: 461-471.
- BARTH, D., KARRER, M., HEINZE-MUTZ, E.M & ELSTER, N. 1994.** Colonisation and degradation of cattle dung: aspects of sampling, fecal composition and artificially formed pats. *Environmental Entomology*. 23(3): 571-578.
- BARTH, D., KARRER, M. & HEINZE-MUTZ, E.M. 1995.** Significance of moisture content of dung pats for colonisation and degradation of cattle dung. *Applied Parasitology*. 36: 11-21.
- BARTHOLOMEW, G.A. & HEINRICH, B. 1978.** Endothermy in African dung beetles during flight, ball making and ball rolling. *Journal of Experimental Biology*. 73: 65-83.
- BRAY, J.R. & CURTIS, J.T. 1957.** An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs*. 27: 325-349.

BEGON, M., HARPER, J.L. & TOWNSEND, C.R. 1995. *Ecology, individuals, populations and communities* (2nd ed.). Blackwell Science, Cambridge.

BERGSTROM, R.C., MAKI, L.R. & WERNER, B.A. 1976. Small dung beetles as biological control agents: laboratory studies of beetle action on Trichostrongylid eggs in sheep and cattle feces. *Proceedings of the Helminthological Society of Washington*. 43(2): 171-174.

BORGHESIO, L., LUZATTO, M., PALESTRINI, C. 1999. Interactions between dung, plants and the dung fauna in a heathland in northern Italy. *Pedobiologia*. 43: 97-109.

BORNEMISSZA, G.F. 1960. Could dung-eating insects improve our pastures? *Journal of Australian Institute of Agricultural Science*. 26: 54-56.

BREYTENBACH, W. & BREYTENBACH, G.J. 1986. Seasonal patterns in dung feeding Scarabaeidae in the southern Cape. *Journal of the Entomological Society of Southern Africa*. 49: 359-366.

CAMBEFORT, Y. 1991. Dung beetles in tropical savannas, pp. 283-304. *In*: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.

CASTLE, M.E. & MACDAID, E. 1972. The decomposition of cattle dung and its effect on pasture. *Journal of the British Grassland Society*. 27: 133-137.

CAVENEY, S, SCHOLTZ, C.H. & McINTYRE, P. 1995. Patterns of daily flight activity in onitine dung beetles (Scarabaeinae: Onitini). *Oecologia*. 103: 444-452.

CHOWN, S.L., & SCHOLTZ, C.H. 1993. Temperature regulation in the nocturnal melolonthine *Sparrmannia flava*. *Journal of Thermal Biology*. 18: 25-33.

- CHOWN, S.L., SCHOLTZ, C.H., KLOK, C.J., JOUBERT, F.J. & COLES, K.S. 1995.** Ecophysiology, range contraction and survival of a geographically restricted African dung beetle (Coleoptera: Scarabaeidae). *Functional Ecology*. 9: 30-39.
- CHOWN, S.L., PISTORIUS, P., SCHOLTZ, C.H. 1998.** Morphological correlates of flightlessness in southern African Scarabaeinae (Coleoptera: Scarabaeidae): testing a condition of the water-conservation hypothesis. *Canadian Journal of Zoology*. 76: 1123-1133.
- COLLINS, N.M. & THOMAS, J.A. 1990.** *The conservation of insects and their habitats*. Academic Press. London.
- CHRISTIANSEN, F.B. & FENCHEL, T.M. 1977.** *Theories of populations and communities*. Berlin.
- DANCKWERTS, J.E. 1982.** The grazing capacity of sweetveld: 2. A model to estimate grazing capacity in the false thornveld of the Eastern Cape. *Proclamations of the Grasslands Society of Southern Africa*. 17: 94-98.
- DANCKWERTS, J.E. & STUART-HILL, G.C. 1988.** The effect of severe drought and management after drought on the mortality and recovery of semi-arid grassveld. *Journal of the Grassland Society of Southern Africa*. 5(4): 218-222.
- DAVIS, A.L.V. 1977.** *The endocoprid dung beetles of southern Africa (Coleoptera: Scarabaeidae)*. M.Sc. thesis, Rhodes University, Grahamstown, South Africa.
- DAVIS, A.L.V. 1987.** Geographical distribution of dung beetles (Coleoptera: Scarabaeidae) and their seasonal activity in south-western Cape Province. *Journal of the Entomological Society of Southern Africa*. 50: 275-285.

- DAVIS, A.L.V. 1993.** Alpha-diversity patterns of dung beetle assemblages (Coleoptera: Scarabaeidae, Aphodiidae, Staphylinidae, Histeridae, Hydrophilidae) in the winter rainfall region of South Africa. *African Entomology*. 1(1): 67-80.
- DAVIS, A.L.V. 1994.** Habitat fragmentation in southern Africa and distributional response patterns in five specialist or generalist dung beetle families (Coleoptera). *African Journal of Ecology*. 32: 192-207.
- DAVIS, A.L.V. 1996.** Habitat associations in a South African, summer rainfall, dung beetle community (Coleoptera: Scarabaeidae, Aphodiidae, Staphylinidae, Histeridae, Hydrophilidae). *Pedobiologia*. 40: 260-280.
- DAVIS, A.L.V., DOUBE, B.M. & McLENNAN, P. 1988.** Habitat associations and seasonal abundance of coprophilous Coleoptera (Staphylinidae, Hydrophilidae, Histeridae) in the Hluhluwe Region of South Africa. *Bullitin of Entomological Research*. 78: 425-434.
- DEMPSTER, J.P. 1991.** Fragmentation, isolation and mobility of insect populations, pp. 143-153. In: Collins, N.M. & Thomas, J.A. (eds.). *The conservation of insects and their habitats*. Academic Press, London.
- DENNO, R.F., McCLURE, M.S. & OTT, J.R. 1995.** Interspecific interactions in phytophagous insects: competition reexamined and resurrected. *Annual Review of Entomology* 40: 297-331.
- DICKINSON, C.H., UNDERHAY, V.S.H. & ROSS, V. 1981.** Effect of season, soil fauna and water content on the decomposition of cattle dung pats. *The New Phytologist*. 88: 129-141.

- DOUBE, B.M. 1983.** The habitat preference of some bovine dung beetles (Coleoptera: Scarabaeidae) in Hluhluwe Game Reserve, South Africa. *Bulltin of Entomological Research*. 73: 357-71.
- DOUBE, B.M. 1987.** Spatial and temporal organization in communities associated with dung pads and carcasses, pp 224-248. In: Gee, J.H. & Giller D.S. (eds.). *Organization of communities past and present*. Blackwell Scientific Publications, Oxford.
- DOUBE, B.M. 1990.** A functional classification for analysis of the structure of dung beetle assemblages. *Ecological Entomology*. 15: 371-383.
- DOUBE, B.M. 1991.** Dung beetles of southern Africa, pp. 133-155. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.
- DOUBE, B.M., GILLER, P.S. & MOOLA, F. 1988.** Dung burial strategies in some South African coprine and onitine dung beetles (Scarabaeidae, Scarabaeinae). *Ecological Entomology*. 13: 251-261.
- DOUBE, B.M., MACQUEEN, A. & FAY H.A.C. 1988.** Effects of dung fauna on survival and size of Buffalo flies (*Haematobia spp.*) breeding in the field in South Africa and Australia. *Journal of Applied Ecology*. 25: 523-536.
- DOUBE, B.M., MACQUEEN, A., RIDSDILL-SMITH, T.J. & WEIR, T.A. 1991.** Native and introduced dung beetles in Australia, pp. 133-155. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.
- DOWNING, B.H. 1978.** Environmental consequences of agricultural expansion in South Africa since 1850. *South African Journal of Science*. 74: 420-422.

DUFRÊNE, M. & LEGENDRE, P. 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs*. 67(3): 345-366.

DUNN, C.H. 1994. Gaps in GAP. *Plant Science Bulletin*. 40: 120-121.

DU PLESSIS, S.F. 1969. *The past and present geographical distribution of the Perissodactyla and Artiodactyla in southern Africa*. M.Sc. thesis, Pretoria University, Pretoria, South Africa.

ECKHOLM, E. 1985. Land, pp. 40-49. In: Myers, N. eds. *The Gaia atlas of planet management*. Pan books, London & Sydney.

EDWARDS, P.B. 1986. Phenology and field biology of the dung beetle *Onitis caffer* (Boheman) (Coleoptera: Scarabaeidae) in Southern Africa. *Bulletin of Entomological Research*. 76: 433-446.

EDWARDS, P.B. 1991. Seasonal variation in the dung of African grazing mammals, and its consequences for coprophagous insects. *Functional Ecology*. 5: 617-628.

ENDRÖDY-YOUNGA, S. 1982. An annotated checklist of dung-associated beetles of the Savanna ecosystem project study area, Nylsvley. *South African National Scientific Programmes Report*. 59: 1-30.

ESTRADA, A, COATES-ESTRADA, R, DADDA, A.A. & CAMMARANO, P. 1998. Dung and carrion beetles in tropical rain forest fragments and agricultural habitats at Los Tuxtlas, Mexico. *Journal of Tropical Ecology*. 14, 577-593.

FAY, H.A.C. & DOUBE, B.M. 1983. The effect of some coprophagous and predatory beetles on the survival of immature stages of the African buffalo fly, *Haematobia thirouxi potans*, in bovine dung. *Zeitschrift für angewandte Entomologie*. 95: 460-466.

- FAY, H.A.C., MACQUEEN, A. & DOUBE, B.M. 1990.** Impact of fauna on mortality and size of *Haematobia spp.* (Diptera: Muscidae) in natural dung pads in Australia and South Africa. *Bulletin of Entomological Research*. 80: 385-392.
- FINCHER, G.T. 1973.** Dung beetles as biological control agents for gastrointestinal parasites of livestock. *Journal of Parasitology*. 59: 396-399.
- FINCHER, G.T. 1981.** The potential value of dung beetles in pasture ecosystems. *Journal of the Georgia Entomological Society*. 16(2): 301-316.
- FINCHER, G.T., STEWART, T.B. & DAVIS, R. 1970.** Attraction of coprophagous beetles to faeces of various animals. *Journal of Parasitology*. 56(2): 378-383.
- FINCHER, G.T., DAVIS, R. & BONNER STEWART, T. 1971.** Flight activity of coprophagous beetles on a swine pasture. *Annals of the Entomological Society of America*. 64 (4): 855-860.
- FLOATE, K.D. & GILL, B.D. 1998.** Seasonal activity of dung beetles (Coleoptera: Scarabaeidae) associated with cattle dung in southern Alberta and their geographic distribution in Canada. *The Canadian Entomologist*. 130: 131-151.
- FOURIE, J.H. & DU TOIT, P.F. 1983.** Weidingstudies in die Vrystaatstreek: Die beplanning van veldbestand. *Glen Agric*. 12(1): 5-9.
- FRY, R. & LONSDALE, D. 1991.** Habitat conservation for insects – a neglected green issue. *The Amateur Entomologist Society*. Feltham, Middlesex, England.

GALANTE, E. 1994. Biogeography of Aphodiinae from the phytogeographic Orocantabrica Province, Cantabrian range, Spain (Col. Scarabaeoidea). *Acta Zoologica Cracov.* 37 (2): 39-58.

GALANTE, E., GARCIA-ROMAN, M. BARRERA, I. & GALINDO, P. 1991. Comparison of spatial distribution patterns of dung-feeding scarabs (Coleoptera: Scarabaeidae, Geotrupidae) in wooded and open pastureland in the Mediterranean "Dehesa" area of the Iberian Peninsula. *Environmental Entomology.* 20(1):90-97.

GALANTE, E, MENA, J. & LUMBRERAS, C. 1993. Study of the spatio-temporal distribution in a coprophagous community in a mediterranean holm-oak ecosystem (Coleoptera: Scarabaeoidea: Scarabaeidae: Geotrupidae). *Elytron.* 7: 87-97.

GALANTE, E., MENA, J. & LUMBRERAS, C. 1995. Dung beetles (Coleoptera: Scarabaeidae, Geotrupidae) attracted to fresh cattle dung in wooded and open pasture. *Environmental Entomology.* 24(5): 1063-1068.

GILL, B.D. 1991. Dung beetles in Tropical American Forests, pp. 133-155. *In:* Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology.* Princeton University Press, Princeton.

GILLARD, P. 1967. Coprophagous beetles in pasture ecosystems. *Journal of the Australian Institute of Agricultural Science.* 33: 30-34.

GILLER, P.S. & DOUBE, B.M. 1989. Experimental analysis of inter-and intraspecific competition in dung beetle communities. *Journal of Animal Ecology.* 58; 129-142.

GILLER, P.S. & DOUBE, B.M. 1994. Spatial and temporal co-occurrence of competitors in Southern African dung beetle communities. *Journal of Animal Ecology.* 63: 629-643.

GITTINGS, T. & GILLER, P.S. 1998. Resource quality and the colonisation and succession of coprophagous dung beetles. *Ecography*. 21: 581-592.

GORDON, R.D. 1983. Studies on the genus *Aphodius* of the United States and Canada (Coleoptera: Scarabaeidae). VII. Food and habitat; distribution; key to eastern species. *Proceedings of the Entomological Society of Washington*. 85(4): 633-652.

GORDON, R.D. & CARTWRIGHT, O.L. 1974. Survey of food preferences of some American Canthonini (Coleoptera: Scarabaeidae). *Entomological News*. 85: 181-185.

GORMALLY, M.J. 1993. The effect of dung beetle activity on the discharge of *Pilobolus sporangia* in cattle faeces. *Medical and Veterinary Entomology*. 7: 197-198.

GOUDIE, A. 1990. *The human impact on the natural environment (3rd ed.)*. Basil Blackwell Ltd., Oxford.

HALFFTER, G. & MATTHEWS E.G. 1966. The natural history of dung beetles of the subfamily Scarabaeinae (Coleoptera: Scarabaeidae). *Folia Entomologica Mexico*. 12-14:1-312.

HANSKI I. 1980(a). Spatial patterns and movements in coprophagous beetles. *Oikos*. 34: 293-210.

HANSKI I. 1980(b). Patterns of beetle succession in droppings. *Annales Zoologica Fennici*. 17: 17-25.

HANSKI, I. 1986. Individual behaviour, population dynamics and community structure of *Aphodius* (Scarabaeidae) in Europe. *Acta Ecologica*. 7 (2): 171-187.

HANSKI, I. 1987(a). Nutritional ecology of dung- and carrion-feeding insects. In: Slansky, F. & Rodriguez, J.G. (Eds.). *Nutritional ecology of insects, mites, and spiders*. John Wiley & Sons, New York.

HANSKI, I. 1987(b). Colonization of ephemeral habitats, pp. 155-186. In: Crawley, M. Edwards. P. & Gray A.J. (eds.). *Colonization, succession and stability*. Blackwell Scientific Publications. Oxford.

HANSKI, I. 1990. Dung and carrion insects, pp. 127-145. In: Shorrocks, B. & Swingland, T. (eds.). *Living in a patchy environment*. Oxford University Press, Oxford.

HANSKI, I. 1991. The dung insect community, pp. 5-21. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.

HANSKI, I & CAMBEFORT, Y. 1991(a). Spatial processes, pp. 283-304. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.

HANSKI, I & CAMBEFORT, Y. 1991(b). Competition in dung beetles, pp. 305-329. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.

HANSKI, I. & KRIKKEN, J. 1991. Dung beetles in tropical forests in south-east Asia, pp.179-197. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.

HANSKI, I. & KOSKELA, H. 1977. Niche relations among dung-inhabiting beetles. *Oecologia*. 28: 203-231.

HANSKI, I. & KOSKELA, H. 1978. Stability, abundance, and niche width in the beetle community inhabiting cow dung. *Oikos*. 31: 290-298.

HANSKI, I. & KOSKELA, H. 1979. Resource partitioning in six guilds of dung-inhabiting beetles (Coleoptera) *Annales Entomologica Fennici*. 45: 1-12.

HERRICK, J.E. & LAL, R. 1996. Soil physical property changes during dung decomposition in a tropical pasture. *Soil Science Society of America Journal*. 59(3): 908-912.

HIRSCHBERGER, P. 1998. Spatial distribution, resource utilisation and intraspecific competition in the dung beetle *Aphodius ater*. *Oecologia*. 116:136-142.

HOLTER, P. 1979(a). Abundance and reproductive strategy of the dung beetle *Aphodius rufipes* (L) (Scarabaeidae). *Ecological Entomology*. 4: 317-326.

HOLTER, P. 1979(b). Effect of dung-beetles (*Aphodius spp.*) and earthworms on the disappearance of cattle dung. *Oikos*. 32: 393-402.

HOLTER, P. 1982. Resource utilization and local coexistence in a guild of scarabaeid dung beetles (*Aphodius spp.*). *Oikos*. 39: 213-227.

HOWDEN, H.F. & NEALIS, V.G. 1975. Effects of clearing in a tropical rain forest on the composition of the coprophagous scarab beetle fauna (Coleoptera). *Biotropica*. 7(2): 77-83.

HOUSTON, W.W.K & McINTYRE, P. 1985. The daily onset of flight in the crepuscular dung beetle *Onitis alexis* (Coleoptera: Scarabaeidae). *Entomol. Exp. Appl.* 39: 223-32.

HUDSON, E.E. & STILING, P. 1997. Exploitative competition strongly affects the herbivorous insect community on *Baccharis halimifolia*. *Oecologia*. 109: 521-528.

HUGHES, R.D., TYNDALE-BISCOE, M. & WALKER, J. 1978. Effects of introduced dung beetles (Coleoptera: Scarabaeidae) on the breeding and abundance of the Australian bushfly, *Musca vetustissima* Walker (Diptera: Muscidae). *Bulletin of Entomological Research*. 68: 361-372.

IVES, A.R. 1991. Aggregation and coexistence in a carrion fly community. *Ecological Monographs*. 61: 75-94.

JANZEN, D.H. 1983. Seasonal change in abundance of large nocturnal dung beetles (Scarabaeidae) in a Costa Rican deciduous forest and adjacent horse pasture. *Oikos*. 41: 274-283.

JAMESON, M.L. 1989. Diversity of coprophagous Scarabaeidae (Coleoptera) in grazed versus ungrazed sandhills prairie in Western Nebraska. *Transactions of the Nebraska Academy of Sciences*. 17: 29-35.

JENKINSON, D.S. 1988. Determination of microbial biomass C and N in soil, pp. 368-386. In: Wilson, J.R. (ed.). *Advances in Nitrogen Cycling in Agricultural Ecosystems*. CAB International, Wallingford.

JONES, R.M. & RATCLIFF, D. 1983. Patchy grazing and its relation to deposition of cattle dung pats in pastures in coastal subtropical Queensland. *Journal of the Australian Institute of Agricultural Science*. 49: 109-111.

KEOUGH, M.J. 1984. Effects of patch size on the abundance of sessile marine invertebrates. *Ecology*. 65: 423-437.

- KIRK, A.A. 1992.** The effect of the dung pad fauna on the emergence of *Musca tempestiva* (Dipt.: Muscidae) from dung pads in southern France. *Entomophaga*. 37(4): 507-514.
- KIRK, A.A. & WALLACE, M.M.H. 1990.** Seasonal variations in numbers, biomass and breeding patterns of dung beetles (Coleoptera: Scarabaeidae) in southern France. *Entomophaga*. 35(4): 569-581.
- KNEIDEL, K.A. 1985.** Patchiness, aggregation and the coexistence of competitors for ephemeral resources. *Ecological Entomology*. 10: 441-448.
- KOSKELA, H. 1979.** Patterns of diel flight activity in dung-inhabiting beetles: an ecological analysis. *Oikos*. 33: 419-439.
- KOSKELA, H. & HANSKI, I. 1977.** Structure and succession in a beetle community inhabiting cow dung. *Annales Zoologica Fennici* 14: 204-223.
- LANDIN, B.O. 1961.** Ecological studies on dung beetles (Col. Scarabaeidae). *Opuscula Entomologica Supplementum*. 19: 1-228.
- LEE, J.M. & PENG, Y.S. 1982.** Influence of manure availability and nesting density on the progeny of *Onthophagus gazella*. *Environmental Entomology*. 11:38-41.
- LOBO, J.M. 1993.** The relationship between distribution and abundance in a dung-beetle community (Col., Scarabaeoidea). *Acta Ecologica*. 14(1): 43-55.
- LOVELL, R.D. & JARVIS, S.C. 1996.** Effect of cattle dung on soil microbial biomass C and N in a permanent pasture soil. *Soil Biology and Biochemistry*. 28(3): 291-299.



- LUMARET, J.P. & KIRK, A.A. 1987. Ecology of dung beetles in the French Mediterranean region (Coleoptera, Scarabaeidae). *Acta Zoologica Mexicana*. 24: 1-60.
- LUMARET, J.P. & KIRK, A.A. 1991. South temperate dung beetles, pp. 97-113. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.
- LUMARET, J.P. & STIERNET, N. 1991. Montane dung beetles, pp. 242-254. In: Hanski, I. & Cambefort, Y. (eds.). *Dung beetle ecology*. Princeton University Press, Princeton.
- LUMARET, J.P., KADIRI, N. & BERTRAND. 1992. Changes in resources: consequences for the dynamics of dung beetle communities. *Journal of Applied Ecology*. 29: 349-356.
- LUMARET, J.P. & IBORRA, O. 1996. Separation of trophic niches by dung beetles (Coleoptera, Scarabaeoidea) in overlapping habitats. *Pedobiologia*. 40: 392-404.
- MAGURRAN, A.E. 1988. *Ecological diversity and its measurement*. Croom Helm, London.
- MACQUEEN, A., WALLACE, M.M.H. & DOUBE, B.M. 1986. Seasonal changes in favourability of cattle dung in central Queensland for three species of dung-breeding insects. *Journal of the Australian Entomological Society*. 25: 23-29.
- MARSHALL LEE, J. & PENG, Y. 1981. Influence of adult size of *Onthophagus gazella* on manure pat degradation, nest construction, and progeny size. *Environmental Entomology*. 10: 626-630.

- MATTHIESSEN, J.N. & HAYLES, L. 1983.** Seasonal changes in characteristics of cattle dung as a resource for an insect in southwestern Australia. *Australian Journal of Ecology*. 8, 9-16.
- MENENDEZ, R. & GUTIERREZ, D. 1996.** Altitudinal effects on habitat selection of dung beetles (Scarabaeoidea: Aphodiidae) in the northern Iberian peninsula. *Ecography*. 19: 313-317.
- MERRITT, R.W. & ANDERSON, J.R. 1977.** The effects of different pasture and rangeland ecosystems on the annual dynamics of insects in cattle droppings. *Hilgardia*. 45(2): 31-71.
- MICO, E., VERDU, J. & GALANTE, E. 1998.** Diversity of dung beetles in Mediterranean wetlands and bordering brushwood. *Annals of the Entomological Society of America*. 91(3):298-302.
- MILLNER, A. 1961.** The mouth parts and digestive tract of adult dung beetles (Coleoptera: Scarabaeidae), with reference to the ingestion of helminth eggs. *Journal of Parasitology*. 47: 735-744.
- MOON, R.D., LOOMIS, E.C. & ANDERSON, J.R. 1980.** Influence of two species of dung beetles on larvae of Face fly. *Environmental Entomology*. 9: 607-612.
- MOHR, C.O. 1943.** Cattle droppings as ecological units. *Ecological Monographs*. 13. 276-259.
- MOORE, N.W. 1991.** Observe extinction or conserve diversity, pp. 1-8. In: Collins, N.M. & Thomas, J.A. (eds.). *The conservation of insects and their habitats*. Academic Press, London.

- NEALIS, V.G. 1977. Habitat associations and community analysis of south Texas dung beetles (Coleoptera: Scarabaeinae). *Canadian Journal of Zoology*. 55: 138-147.
- NEL, C. 1991. Veld condition in South Africa. *Farmer's Weekly*. (February 15, pp. 38, 39 & 42).
- ODUM, E.P. 1983. *Basic Ecology*. Saunders, New York.
- ODUM, E.P. 1993. *Ecology and our endangered life-support systems*. Sinauer Associates, Inc. Publishers. Sunderland, Massachusetts.
- OMALIKO, C.P.E. 1984. Dung decomposition and its effects on the soil component of a tropical grassland ecosystem. *Tropical Ecology*. 25: 214-220.
- OPPERMAN, D.P.J. 1980. Die benutting en die bewaring van die weidingsekosisteem in die R.S.A. *Journal of the Grassland Society of Southern Africa*. 15: 13-17.
- OSBERG, D.C., DOUBE, B.M. & HANRAHAN, S.A. 1993. Habitat specificity in African dung beetles: the effect of soil type on dung burial by two species of ball-rolling dung beetles (Coleoptera: Scarabaeidae). *Tropical Zoology*. 6: 243-251.
- OTRONEN, M. & HANSKI, I. 1983. Movement patterns in *Sphaeridium*: differences between species, sexes, and feeding and breeding individuals. *Journal of Animal Ecology*. 52: 663-680.
- OWEN-SMITH, N. 1989. *Megaherbivores: the influence of very large body size on ecology*. Cambridge University Press, Cambridge.

PECK, S.B. & FORSYTH, A. 1982. Composition, structure, and competitive behaviour in a guild of Ecuadorian rain forest dung beetles (Coleoptera; Scarabaeidae). *Canadian Journal of Zoology*. 60: 1624-1634.

PEITMEIER, B.A., CAMPBELL, J.B. & THOMAS, G.D. 1992. Insect fauna of bovine dung in northeastern Nebraska and their possible effect on the face fly *Musca autumnalis* (Diptera: Muscidae). *Journal of the Kansas Entomological Society*. 65(3): 267-274.

PEREZ-MENDOZA, J., HAGSTRUM, D.W., DOVER, B.A., HOPKINS, T.L., BAKER, J.E. 1999. Flight response, body weight, and lipid content of *Rhyzopertha dominica* (F.) (Coleoptera: Bostrichidae) as influenced by strain, season and phenotype. *Journal of Stored Products Research* 35: 183-196.

PETERSEN, R.G., WOODHOUSE, W.W. & LUCAS, H.L. 1956. The distribution of excreta by freely grazing cattle and its effect on pasture fertility: II. Effect of returned excreta on the residual concentration of some fertilizer elements. *Agron.* 48: 444-449.

POTGIETER, D.J. 1975. *Standard encyclopaedia of Southern Africa*. Vol. 11. pp. 116, 448. Nasou Ltd., Cape Town.

PRICE, P.W. 1984. *Insect ecology* (2nd ed.). John Wiley & Sons, New York.

RAINIO, M. 1966. Abundance and phenology of some coprophagous beetles in different kinds of dung. *Annales Zoologica Fennici*. 3: 88-98.

REEVE, J.D.; RHODES, D.J. & TURCHIN, P. 1998. Scramble competition in the southern pine beetle, *Dendroctonus frontalis*. *Ecological Entomology*. 23: 433-443.

RIDSDILL-SMITH, T.J. 1981. Some effects of three species of dung beetles (Coleoptera: Scarabaeidae) in south-western Australia on the survival of the bush fly, *Musca vetustissima* Walker (Diptera: Muscidae), in dung pads. *Bulletin of Entomological Research*. 71: 425-433.

RIDSDILL-SMITH, T.J. 1986. The effect of seasonal changes in cattle dung on egg production by two species of dung beetles (Coleoptera: Scarabaeidae) in south-western Australia. *Bulletin of Entomological Research* 76: 63-68.

RIDSDILL-SMITH, T.J. 1990. Competition in dung breeding insects, pp. 264-292. In: Bailey, W.J. & Risdill-Smith, T.J. (eds.). *Reproductive behaviour in insects – individuals and populations*. Chapman & Hall. London.

RIDSDILL-SMITH, T.J. & HALL, G.P. 1984. Beetles and mites attracted to fresh cattle dung in southwestern Australian pastures. *CSIRO Division of Entomology Report*. No. 34.

RIDSDILL-SMITH, T.J., HALL, G.P. & CRAIG, G.F. 1982. Effect of population density on reproduction and dung dispersal by the dung beetle *Onthophagus binodis* in the laboratory. *Entomology Experiment and Application*. 32: 80-85.

RIDSDILL-SMITH, T.J. & HAYLES, L. 1987. Mortality of eggs and larvae of the bush fly, *Musca vetustissima* Walker (Diptera: Muscidae), caused by scarabaeine dung beetles (Coleoptera: Scarabaeidae) in favourable cattle dung. *Bulletin of Entomological Research*. 77: 731-736.

ROMOSER, W.S. 1981. *The science of entomology (2nd ed)*. Macmillan Publishing Co., Inc., New York.

SAMWAYS, M.J. 1994. *Insect conservation biology*. Chapman & Hall. London.

SCHOENER, T.W. 1983. Field experiments on interspecific competition. *American Naturalist*. 122: 240-285.

SCHOENER, T.W. 1986. Patterns in terrestrial vertebrate versus arthropod communities: do systematic differences in regularity exist? pp. 556-586. *In*: Diamond, J. & Case, T.J. (eds.). *Ecology*. Harper & Row, New York, USA.

SCHMIDT-NIELSEN, K. 1988. *Animal physiology: Adaptation and environment*. Cambridge University Press, Cambridge.

SKINNER, T.E. 1981. Droughts in perspective. *Proclamations of the Grassland Society of Southern Africa*. 16: 13-16

SNOWBALL, G.J. 1944. A consideration of the insect population associated with cow dung at Crawley, W.A. *J. Proc. R. Soc. West. Aust.* 28:219-245.

SNYMAN, H.A. & FOUCHÉ. 1993. Estimating seasonal herbage production of a semi-arid grassland based on veld condition, rainfall and evapotranspiration. *African Journal of Rangeland Science*. 10(1): 21-24.

SOUTHWOOD, T.R.E. 1978. *Ecological methods*. Chapman and Hall, London.

SOWIG P. 1997. Niche separation in coprophagous beetles: a comparison of two multivariate approaches. *Bulletin of Entomological Research*. 87:625-631.

STEWART, T.B. 1967. Food preferences of coprophagous beetles with special reference to *Phanaeus spp.* *Journal of the Georgia Entomological Society*. 2(3): 69-77.

STOTT, P. 1998. Biogeography and ecology in crisis: the urgent need for a new metalanguage. *Journal of Biogeography*. 25: 1-2.

TEAGUE, W.R. & SMIT, G.N. 1992. Relations between woody and herbaceous components and the effects of bush clearing in Southern African savannas. *Journal of the Grassland Society of Southern Africa*. 9(2): 60-69.

TOKESHI, M. 1993. Species abundance patterns and community structure. *Advances in Ecological Research*. 24: 11-186.

TRIBE, G.D. 1976. *The ecology and ethology of ball-rolling dung beetles (Coleoptera: Scarabaeidae)*. M.Sc. thesis, University of Natal, Pietermaritzburg, South Africa.

TROLLOPE, W.S.W. 1975. Vegetation and veld management. In: *The agricultural potential of the Ciskei*. Institute for Planning Research, Stellenbosch.

TYNDALE-BISCOE, M. 1988. The phenology of *Onitis alexis* (Coleoptera: Scarabaeidae) in the Araluen Valley: Survival in a marginal environment. *Australian Journal of Ecology*. 13: 431-443.

TYNDALE-BISCOE, M. 1994. Dung burial by native and introduced dung beetles (Scarabaeidae). *Australian Journal of Agriculture Research*. 45:1799-1808.

USHER, M.B. & JEFFERSON, R.G. 1991. Creating new and successional habitats for Arthropods, pp. 263-283. In: Collins, N.M. & Thomas, J.A. (eds.). *The conservation of insects and their habitats*. Academic Press, London.

VALIELA, I. 1974. Composition, food webs and population limitation in dung arthropod communities during invasion and succession. *The American Midland Naturalist*. 92(2): 370-385.

VAN OUDTSHOORN, F. 1991. *Gids tot grasse van Suid-Afrika*. Briza Publikasies, Arcadia.

VILJOEN, A.J. 1979. *Die plant-ekologie van die Sandveld Natuurresewaat, Hoopstad*. M.Sc. thesis, University of the Orange Free State, Bloemfontein, South Africa.

WALKER, J.M. & DOUBE, B.M. 1984. Susceptibility to drowning of eggs of the dung breeding flies *Haematobia irritans exigua* and *Musca vetustissima* (Dipt., Muscidae) in relation to dung beetle activity. *Zeitschrift für angewandte Entomologie*. 98: 518-527.

WASSMER, T. 1994. Seasonality of coprophagous beetles in the Kaiserstuhl area near Freiburg (SW-Germany) including the winter months. *Acta Ecologica*. 15 (5): 607-631.

WATERHOUSE, D.F. 1974. The biological control of dung. *Scientific American*. 230(4):101-109.

WHITE, P.S. & PICKETT, S.I.A. 1985. Natural disturbance and patch dynamics: an introduction. In: Pickett, S.T.A. & White, P.S. (eds.). *The ecology of natural disturbance and patch dynamics*. Academic Press, New York.

WINGO, C.W., THOMAS, G.D., CLARK, G.N. & MORGAN, C.E. 1974. Succession and abundance of insects in pasture manure: relationship to Face fly survival. *Annals of the Entomological Society of America*. 67: 386-390.

WOLDA, H. 1978. Fluctuations in abundance of tropical insects. *American Naturalist*. 112:1017-1045.