

REFERENCES

- ¹ King H, Aubert RE, Herman WH. Global burden of diabetes, 1995-2025: prevalence, numerical estimates, and projections. **Diabetes Care** 1998;21:1414-1431.
- ² Songer TJ, Zimmer PZ. Epidemiology of Type II Diabetes. **Pharmacoeconomics** 1995; 8 Suppl 1:1-9.
- ³ Omar MAK, Deedat MA, Motala AA. The prevalence of Diabetes Mellitus and impaired glucose tolerance in a group of urban South African Blacks. **S Afr Med J** 1993;83:641-43.
- ⁴ Gu K, Cowie CC, Harris MI. Mortality in adults with and without diabetes in a national cohort of the US population, 1971-1993. **Diabetes Care** 1998;21: 1138-45.
- ⁵ South African Medical Research Council: Burden of disease unit at <http://www.mrc.ac.za/bod/bod.htm> accessed on 12 September 2003.
- ⁶ Rosenbloom AL, Joe JR, Young RS, Winter WE. Emerging epidemic of Type 2 DM in youth. **Diabetes Care** 1999;22:345-54.
- ⁷ Wareham NJ. The effectiveness of interventions aimed at weight loss, and other effects of diet and physical activity in achieving control of diabetes and preventing its complications. In: Williams R, Herman W, Kinmonth AL, Wareham NJ, editors. *The Evidence Base for Diabetes Care*. West Sussex: John Wiley & Sons; 2002. p. 355-78.

- ⁸ Huddle KRL. Diabetes Care in South Africa. In: Gill G, Mbanya JC, Alberti G, editors. *Diabetes in Africa*. Cambridge: Biddles Ltd.; 1997. p. 189-96.
- ⁹ Department of Statistics, Tswane Metro, Pretoria; 2003.
- ¹⁰ Department of Health, Tswane Metro, Skinner Street, Pretoria; 2003.
- ¹¹ Rheeder P, Van Wyk JT, Hokken JWE, Hueting HM. Monofilament assessment of neuropathy in a community diabetes clinic. **S Afr Med J** 2002;92:715-19.
- ¹² Boule NG, Haddad E, Kenny GP, Wells GA, Sigal RJ. Effect of exercise on glycaemic control and body mass in Type 2 DM mellitus: a meta-analysis of controlled clinical trials. **JAMA** 2001;286:1218-27.
- ¹³ Tuomilehto J. Missed and newly recovered potential for the prevention of Type 2 DM: a commentary. In: Williams R, Herman W, Kinmonth AL, Wareham NJ, editors. *The Evidence Base for Diabetes Care*. West Sussex: John Wiley & Sons; 2002. p. 180-85.
- ¹⁴ Albright A, Franz M, Hornsby G, Kriska A, Marrero D, Ullrich I. American College of Sports Medicine position stand: exercise and Type 2 DM. *Med Sci Sports Exerc* 2000;32:1345-60.
- ¹⁵ Willey K, Singh MAF. Battling Insulin resistance in elderly obese people with Type 2 DM. **Diabetes Care** 2003;26:1580-88.
- ¹⁶ Foster C, Cadwell K, Crenshaw B, Dehart-Beverly M, Hatcher S, Karlsdottir AE, et al Physical activity and exercise training prescriptions for patients. **Cardiol Clin** 2001;19:447-55.
- ¹⁷ Myers J, Atwood JE, Froelicher V, editors. *Active lifestyle and Diabetes*. **Circulation** 2003;107:2392-4.

- ¹⁸ Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C et al. Physical activity and public health: a recommendation from the Centres for Disease Control and Prevention and the American College of Sports Medicine. **JAMA** 1995;273:402-7.
- ¹⁹ Chale S. Economics of diabetes care in Africa. In: Gill G, Mbanya JC, Alberti G, editors. Diabetes in Africa. Cambridge: Biddles Ltd.; 1997. p. 199-213.
- ²⁰ American College of Sports Medicine. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness in health adults. **Med Sci Sports Exerc** 1998;30:975-91.
- ²¹ Oldridge NB. Cardiac rehabilitation, self-responsibility and quality of life. **J Cardiopulm Rehabil** 1986;6:153-6.
- ²² Tshabalala G, Gill GV. Cultural aspects of diabetes in Africa. In: Gill G, Mbanya JC, Alberti G, editors. Diabetes in Africa. Cambridge: Biddles Ltd.; 1997. p.275-87.
- ²³ Schapera I, Goodwin JH. Work and Health. In: Schapera I editor. Bantu-speaking tribes of South Africa. London: Routledge & Kegan Paul Ltd.; 1962.p. 150-2.
- ²⁴ Knowles JH, editor. Responsibility for health. **Science** 1977;198:4322.
- ²⁵ American Diabetes Association. Standards of medical care for patients with Diabetes Mellitus. **Diabetes Care** 2003;26 Suppl 1:33-47.
- ²⁶ Goldburgh S. Treatment priorities for patients with Type 2 DM. **S Afr Fam Pract** 2003;4:30-6.

- ²⁷ American College of Sports Medicine. Guidelines for exercise testing and prescription. 6th ed. Philadelphia: Lippincott Williams and Wilkins; 2000.
- ²⁸ South African Medical Research Council: Burden of disease unit at <http://www.mrc.ac.za/bod/bod.htm> accessed on 12 September 2003.
- ²⁹ Dunn SM, Bryson JM, Hoskins PL, Alford JB, Handelsman DJ, Turtle JR. Development of the Diabetes Knowledge (DKN) Scales: Forms DKNA, DKNB and DKNC. **Diabetes Care** 1984;7:36-41.
- ³⁰ Voorrips LE, Ravelli PCA, Dongelmans P, Deurenberg P, Van Staveren WA. A physical activity questionnaire for the elderly. **Med Sci Sports Exerc.** 1991;23:974-9.
- ³¹ Anderson RM, Donnelly MB, Dedrick RF. Measuring the attitudes of patients towards diabetes and its treatment. **Patient Educ Couns** 1990; 16:231-45.
- ³² Maillet NA, Melkus GD, Spollett G. Using focus groups to characterize the health beliefs and practices of black women with Non-insulin-dependent Diabetes. **Diabetes Educator** 1996;22:39-46.
- ³³ Altman DC. Practical for medical research. 2nd ed. London: Chapman & Hall; 1992: p.189, 325.
- ³⁴ Stewart KJ. Exercise training and the cardiovascular consequences of Type 2 DM and Hypertension. **JAMA** 2002; 288:1622-31.
- ³⁵ Expert Committee on the diagnosis and classification of Diabetes Mellitus **Diabetes Care** 2003;26 Suppl 1:5-9.

- ³⁶ World Health Organization. Study group on Prevention of Diabetes Mellitus. Geneva, Switzerland; 1994. Technical report series No: 844.
- ³⁷ Bennett PH. Epidemiology of Type 2 Diabetes Mellitus. In: Le Roith D, Taylor SI, Olefsky JM, editors. Diabetes Mellitus. 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2000. p. 544-56.
- ³⁸ Meyer BJ, Van Papendorp DH, Meij HS, Viljoen, Claassen N, Haag M. Human Physiology: chemical, physical and physiological principles. Lansdowne: Juta; 2002.p. 18.42-18.50.
- ³⁹ Duncan GE, Perri MG, Theriaque DW, Hutson AD, Eckel RH, Stacpoole PW. Exercise training, without weight loss, increases Insulin sensitivity and postheparin plasma lipase activity in previously sedentary adults. **Diabetes Care** 2003;26:557-62.
- ⁴⁰ McLarty D, Pollitt C, Swai A, Alberti KGMM. Epidemiology of diabetes in Africa. In: Gill G, Mbanya JC, Alberti G, editors. Diabetes in Africa. Cambridge: Biddles Ltd.; 1997.p. 1-17.
- ⁴¹ De Courten M, Bennett PH, Tuomilehto J, Zimmet P. Epidemiology of NIDDM in Non-Europids. In: Alberti KGMM, Zimmet P, DeFronzo RA, Keen H, editors. International Textbook of Diabetes Mellitus 2nd ed. West Sussex: John Wiley & Sons Ltd.; 2001.p. 150.
- ⁴² Motala AA, Pirie FJ, Gouws E, Amod A, Omar MAK. High incidence of Type 2 DM mellitus in South African Indians: a 10-year follow-up study. **Diabet Med** 2003;20:23-30.
- ⁴³ Motala AA. NIDDM in Africans: is it increasing? **IDF Bulletin** 1995;40:23-6.
- ⁴⁴ Skene DJ, Dali P, Bam WJ, et al Diabetes in a developing country. Epidemiology and management problems. **S Afr Med J** 1982; 61:463-64.

- ⁴⁵ Levitt NS, Steyn K, Lambert EV, Reagon G, Lombard CJ, Fourie JM, et al Modifiable risk factors for Type 2 DM mellitus in a peri-urban community in South Africa. **Diabet Med** 1999;16:946-50.
- ⁴⁶ Kriska AM, LaPorte RE, Pettitt DJ, Charles M, Nelson R, Kuller L, et al The association of physical activity with obesity, fat distribution and glucose intolerance in Pima Indians. **Diabetologia** 1993;36:863.
- ⁴⁷ Omar MAK, Asmal AC. Family histories of diabetes mellitus in young African and Indian diabetics. **Br Med J** 1983;286:1786.
- ⁴⁸ American Diabetes Association. Type 2 DM in children and adolescents: Consensus statement. **Diabetes Care** 2000;23:381-89.
- ⁴⁹ National Institutes of Health. Physical Activity and cardiovascular health: Consensus Statement. Leon AS editor. Champaign: Human Kinetics; 1997. p. 3-13.
- ⁵⁰ Zimmet PZ. Challenge in diabetes epidemiology -from west to the rest: Kelly West Lecture 1991. **Diabetes Care** 1992;15:232-52.
- ⁵¹ Huddle KRL. Diabetes Care in South Africa. In: Gill G, Mbanya JC, Alberti G, editors. Diabetes in Africa. Cambridge: Biddles Ltd.; 1997. p. 189-96.
- ⁵² Caspersen CJ, Matthew MZ. The prevalence of physical inactivity in the United States. In: National Institutes of Health: Consensus Statement: Physical Activity and cardiovascular health. Leon AS, editor. Champaign: Human Kinetics; 1997. p. 35-9.
- ⁵³ Gill GV, Huddle KR. Patterns and problems of diabetes in Soweto. **Cent Afr J Med** 1984;30:189-95.

- ⁵⁴ Gill GV, Rolfe M. Manifestations of diabetes. In: Gill G, Mbanya JC, Alberti G, editors. *Diabetes in Africa*. Cambridge: Biddles Ltd.; 1997.p. 29-33.
- ⁵⁵ Diabetes Control and Complications Trial Research Group (UKPDS) The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. **N Engl J Med** 1993;329:977-86.
- ⁵⁶ UK Prospective Diabetes Study Group: Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with Type 2 DM (UKPDS 33). **Lancet** 1998;352:837-53.
- ⁵⁷ UK Prospective Diabetes Study Group: Effect of intensive blood-glucose control with metformin on complications in overweight patients with Type 2 DM (UKPDS 34). **Lancet** 1998;352:854-65.
- ⁵⁸ Guyton AC, Hall JE. *Textbook of Medical Physiology*. 10th ed. Philadelphia: WB Saunders Company.; 2000. p.874-97.
- ⁵⁹ Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, et al Association of glycaemia with macrovascular and microvascular complications of Type 2 DM (UKPDS 35): prospective observational study. **BMJ** 2000;321:405-12.
- ⁶⁰ Shichiri M, Kishikawa H, Ohkubo Y, Wake N. Long-term results of the Kumamoto Study on optimal diabetes control in Type 2 diabetic patients. **Diabetes Care** 2000;23 Suppl 2:21-9.
- ⁶¹ Wing RR, Koeske R, Epstein LH, Nowalk MP, Gooding W, Becker D. Long-term effects of modest weight loss in Type 2 diabetic patients. **Arch Intern Med** 1987;147:1749-53.

- ⁶² Guare JC, Wing RR, Grant A. Comparison of obese NIDDM and nondiabetic women: short and long-term weight loss. **Obes Res** 1995;3:329-35.
- ⁶³ Brown SA, Upchurch S, Winter M, Ramirez G, Anding R. Promoting weight loss in Type 2 diabetics. **Diabetes Care** 1996;19:613-24.
- ⁶⁴ Expert Panel on the Identification, Evaluation and Treatment of Overweight and Obesity in Adults: Clinical Guidelines. In: Delahanty LM. Evidence-based trends for achieving weight loss and increased physical activity: applications for diabetes prevention and treatment. **Diabetes Spectrum** 2002;15:183-90.
- ⁶⁵ Joslin EP, Root HF, White P, Marble A. The treatment of diabetes mellitus. 5th ed. Philadelphia: Lea & Febiger; 1935: p.299.
- ⁶⁶ Goodyear LJ, Smith RJ. Exercise and Diabetes. In: Kahn CR, Weir GC, editors. Joslin's Diabetes Mellitus. 13th ed. Philadelphia: Lea & Febiger; 1994. p. 451-7.
- ⁶⁷ Kemmer FW, Gudat U, Berger M. Exercise. In: Alberti KGMM, Zimmet P, DeFronzo RA, editors. 2nd ed. Chichester: John Wiley & Sons; 1994. p. 799-815.
- ⁶⁸ Horton ES. Exercise in patients with Type 2 DM Mellitus. In: LeRoith D, Taylor SI, Olefsky JM, editors. Diabetes Mellitus. 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2000. p. 765-9.
- ⁶⁹ Ivy JL. Role of exercise training in the prevention and treatment of insulin resistance and Non-insulin-dependent Diabetes Mellitus. **Sports Med** 1997;5:321-36.

- ⁷⁰ Cartee GD, Young DA, Sleeper MD et al Prolonged increase in insulin-stimulated glucose transport in muscle after exercise. **Am J Physiol** 1989;256(4Pt 1): E494-9.
- ⁷¹ Mayer-Davies E, D'Agostino R, Karter A, Haffner S, Rewers M, Saad M, et al Intensity and amount of physical activity in relation to insulin sensitivity: the Insulin Resistance Atherosclerosis Study. **JAMA** 1998;279:669-74.
- ⁷² Reaven GM. Banting Lecture 1988. Role of insulin resistance in human disease. **Diabetes** 1988;37:1595-1607.
- ⁷³ Hickner R, Fisher JS, Hansen PA, et al Muscle glycogen accumulation after endurance exercise in trained and untrained individuals. **J Appl Physiol** 1997;83:897-903.
- ⁷⁴ Laakso M, Edelman SV, Brechtel G, Baron AD. Impaired insulin-mediated skeletal muscle blood flow in patients with NIDDM. **Diabetes** 1992;41:1076-83.
- ⁷⁵ Edelman SV, Laakso M, Wallace P, Olefsky JM, Baron AD. Kinetics of insulin-mediated and non-insulin-mediated glucose uptake in humans. **Diabetes** 1990;39:955-64.
- ⁷⁶ Dela F, Larsen JJ, Mikines KJ, Ploug T, Petersen LN, Galbo H. Insulin-stimulated muscle glucose clearance in patients with NIDDM. **Diabetes** 1995;44:1010-20.
- ⁷⁷ Olefsky JM. The insulin receptor: Its role in insulin resistance in obesity and diabetes. **Diabetes** 1976;25:1154-64.
- ⁷⁸ Dohm GL, Sinha MK, Caro JF. Insulin receptor binding and protein kinase activity in muscles of trained rats. **Am J Physiol** 1987;252(2 Pt 1): E 170-175.

- ⁷⁹ Dela F, Ploug T, Handberg A, Petersen LN, Larsen JJ, Mikines KJ. Physical training increases muscle GLUT4 protein and mRNA in patients with NIDDM. **Diabetes** 1994;43:862-5.
- ⁸⁰ Ivy JL, Holloszy JO. Persistent increase in glucose uptake by rat skeletal muscle following exercise. **Am J Physiol** 1981;241:C200-3.
- ⁸¹ Kelley DE, Mokan M, Mandarino LJ. Intracellular defects in glucose metabolism in obese patients with NIDDM. **Diabetes** 1992;41:698-706.
- ⁸² Holm G, Krotkiewski M. Potential importance of the muscles for the development of insulin resistance in obesity. **Acta Med Scand** 1988;723(Suppl):95-101.
- ⁸³ Hickey MS, Carey JO, Azevedo JL, et al Skeletal muscle fibre composition is related to adiposity and in vitro glucose transport rate in humans. **Am J Physiol** 1995; 268(3Pt 1): E453-57.
- ⁸⁴ Krotkiewski M, Bjorntorp P. Muscle tissue in obesity with different distribution of adipose tissue. Effects of physical training. **Int J Obesity** 1986;10:331-41.
- ⁸⁵ Saltin B, Henriksson J, Nygaard E, et al Fibre Types and metabolic potentials of skeletal muscles in sedentary man and endurance runners. In: Ivy JL. Role of exercise training in the prevention and treatment of insulin resistance and Non-insulin-dependent Diabetes Mellitus. **Sports Med** 1997;24:321-36.
- ⁸⁶ Nestel PJ, Ishikawa T, Goldrick RB. Diminished plasma free fatty acid clearance in obese subjects. **Metabolism** 1978;27:589-97.

- ⁸⁷ Craig BW, Hammons GT, Garthwaite SM, et al Adaptations of fat cells to exercise: response of glucose uptake and oxidation to insulin. **J Appl Physiol** 1981;51:1500-6.
- ⁸⁸ Piatti PM, Monti ID, Pacchioni M, Pontiroli AE, Pozza G. Forearm insulin- and non-insulin-mediated glucose uptake and muscle metabolites in man: role of free fatty acids and blood glucose levels. **Metabolism** 1991;40:926-33.
- ⁸⁹ Schneider SH, Amorosa LF, Kachadorian AK, Ruderman NB. Studies of the mechanisms of improved glucose control during regular exercise in Type 2 (non-insulin-dependent) diabetes. **Diabetologia** 1984;26:355-60.
- ⁹⁰ King H, Zimmet PZ, Gareeboo H et al Risk factors for diabetes in three Pacific populations. **Am J Epidemiol** 1984;119:396-409.
- ⁹¹ Dowse GK, Zimmet PZ, Gareeboo H, Alberti KGMM, Tuomilehto J, Finch CF, et al Abdominal obesity and physical inactivity as risk factors for NIDDM and impaired glucose tolerance in Indians, Creole and Chinese Mauritians. **Diabetes Care** 1991;14:271-82.
- ⁹² Zimmet P, Faaiuso S, Ainuu S, Whitehouse S, Milne B, DeBoer W. The prevalence of diabetes in the rural and urban Polynesian population of Western Samoa. **Diabetes** 1981;30:45-51.
- ⁹³ Manson JE, Nathan DM, Krolewski AS, Stampfer MJ, Willett WC, Hennekens CH. A prospective study of exercise and incidence of diabetes among US male physicians. **JAMA** 1992;268:63-7.
- ⁹⁴ Perry IJ, Wannamethee SG, Walker MK, Thomson AG, Whincup PH, Shaper AG. Prospective study of risk factors for development of non-insulin dependent diabetes in middle aged British men. **BMJ** 1995;310:560-4.

- ⁹⁵ Kriska AM, Harris SB, Hanley AJG, Zinman B. Physical activity, physical fitness and Insulin and Glucose concentrations in an isolated Native Canadian Population experiencing rapid lifestyle change. **Diabetes Care** 2001;24:1787-92.
- ⁹⁶ Kriska AM, Blair SN, Pereira MA. The potential role of physical activity in the prevention of non-insulin-dependent diabetes mellitus: the epidemiological evidence. **Exerc Sports Sci Rev** 1991;22:121-143.
- ⁹⁷ Helmrich SP, Ragland DR, Leung RW, Paffenberg RS. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. **N Engl J Med** 1991;325:147-52.
- ⁹⁸ Schranz A, Tuomilehto J, Marti B, Jarrett RJ, Grabauskas V, Vassallo A. Low physical activity and worsening of glucose tolerance results from a 2-year follow-up of a population sample in Malta. **Diabetes Res Clin Pract** 1991;11:127-36.
- ⁹⁹ Hu FB, Sigal RJ, Rich-Edwards JW, et al Walking compared with vigorous physical activity and risk of Type 2 DM in women: A prospective study. **JAMA** 1999;282:1433.
- ¹⁰⁰ Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Walter WC, et al **JAMA** 1999;282:1433.
- ¹⁰¹ Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, et al Diet, lifestyle and the risk of Type 2 DM Mellitus in women. **N Engl J Med** 2001;234:790-97.
- ¹⁰² Hu FB, Stampfer MJ, Solomon C et al Physical activity and risk for cardiovascular events in diabetic women. **Ann Intern Med** 2001;134:96-105.

- ¹⁰³ Kelley DE, Goodpaster BH. Effects of exercise on glucose homeostasis in Type 2 DM mellitus. **Med Sci Sports Exerc** 2001;33 Suppl 6:495-501.
- ¹⁰⁴ Balkau B, King H, Zimmet P, Raper LR. Factors associated with the development of diabetes in the Micronesian population of Nauru. **Am J Epidemiol** 1985;122:594-605.
- ¹⁰⁵ James SAL, Jamjoum TE, Raghunathan DS, Strogatz DS, Firth ED, Khazanie PG. Physical activity and NIDDM in African-Americans. The Pitt County Study. **Diabetes Care** 1998;21:555-62.
- ¹⁰⁶ Eriksson KF, Lindgarde F. Prevention of Type 2 (non-insulin-dependent) diabetes mellitus by diet and physical exercise. **Diabetologia** 1991;34:891-8.
- ¹⁰⁷ Pan X-R, Cao H-B, Li G-W, Liu P-A, Hu Y-H, Jiang X-G, et al The effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. **Diabetes Care** 1997;20:537-44.
- ¹⁰⁸ McNabb W, Cook S, Quinn M, Karrison T, Kerver J. The PATHWAYS Church-based weight loss program for urban African-American women at risk for diabetes. **Diabetes Care** 1997;10:1518-23.
- ¹⁰⁹ Diabetes Prevention Programme Research Group. The Diabetes Prevention Programme (DPP). **Diabetes Care** 2002;25:2165-71.
- ¹¹⁰ Narayan KM Hoskin, M, Kozak D, Hanson RL, Pettitt DJ, Bennett PH, et al Randomized clinical trial of lifestyle interventions in Pima Indians: a pilot study. **Diabet Med** 1998;15:66-72.

- ¹¹¹ Eriksson J, Lindstrom J, Valle T, Aunola S, Hamalainen H, Ilanne-Parikka P, et al Prevention of Type II diabetes in subjects with impaired glucose tolerance: the Diabetes Prevention Study (DPS) in Finland- Study design and one-year interim report on the feasibility of the lifestyle intervention programme. **Diabetologia** 1999;42:793-801.
- ¹¹² Wing RR, Polley BA, Venditti E, Lang W, Jakicic JM. Lifestyle intervention in overweight individuals with a family history of diabetes. **Diabetes Care** 1998;21:350-59.
- ¹¹³ Irwin ML, Durstine JL, Mayer-Davis EJ, Stolarczyk LM, Addy CL, Ainsworth BE, et al Moderate-intensity physical activity and fasting insulin levels in women: The Cross-cultural Activity Participation Study. **Diabetes Care** 2000; 23:449-54.
- ¹¹⁴ Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, et al Prevention of Type 2 DM by changes in lifestyle among subjects with impaired glucose tolerance. **N Engl J Med** 2001;344:1343-50.
- ¹¹⁵ Gaede p, Vedel P, Larsen N, Pedersen. Multifactorial intervention and cardiovascular disease in patients with Type 2 DM. **N Engl J Med** 2003;348:383-93.
- ¹¹⁶ Tanasescu M, Leitzmann MF, Rimm EB et al Physical activity in relation to cardiovascular disease and total mortality among men with Type 2 DM. **Circulation** 2003;107:2435-9.
- ¹¹⁷ Delahanty LM. Evidence-based trends for achieving weight loss and increased physical activity: applications for diabetes prevention and treatment. **Diabetes Spectrum** 2002;15:183-90.

- ¹¹⁸ US Dept of Health and Human Services. *Healthy People 2000: National Health Promotion and Disease Prevention Objectives*. Washington, DC: US Dept of Health and Human Services; 1991. DHHS publication PHS 91-50212.
- ¹¹⁹ Wei M, Gibbons LW, Kampert JB, Nichaman MZ, Blair SN. Low cardiorespiratory fitness and physical inactivity as predictors of mortality in men with Type 2 DM. **Ann Intern Med** 2000;132:605-11.
- ¹²⁰ Dunn AL, Marcus BH, Kampert JB, Garcia ME, Kohl HW, Blair SN. Reduction in cardiovascular disease risk factors: 6-month results from Project Active. **Preventive Med** 1997;26:883-92.
- ¹²¹ American Diabetes Association. Diabetes Mellitus and exercise. **Diabetes Care** 2002;25 Suppl 1:64-68.
- ¹²² Brown SA. Interventions to promote diabetes self-management: state of the science. **Diabetes Educator** 1999; 25 Suppl 6:52-61.
- ¹²³ Reaven GM, Laws A. Insulin resistance, compensatory hyperinsulinaemia, and coronary heart disease. **Diabetologia** 1994;37:948-52.
- ¹²⁴ American Diabetes Association. Physical activity / Exercise and Diabetes Mellitus. **Diabetes Care** 2003;26 Suppl 1:73-77.
- ¹²⁵ Dattilo A, Kris-Etherton P. Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. **Am J Clin Nutr** 1992;56:320-28.
- ¹²⁶ Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, et al. Effect of exercise on total and intra-abdominal body fat in postmenopausal women. **JAMA** 2003;289:323-30.

- ¹²⁷ Eriksson J, Taimela S, Koivisto VA. Exercise and the metabolic syndrome. **Diabetologia** 1997;40:125-35.
- ¹²⁸ Keim N, Barbieri T, Van Loan M, et al Energy expenditure and physical performance in overweight women: response to training with and without caloric restriction. **Metabolism** 1990;39:651-57.
- ¹²⁹ Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults: Second report of the National Cholesterol Education Program. **JAMA** 1993;269:3015-23.
- ¹³⁰ Nielsen FS, Rossing P, Gall MA, Skoot F, Smidt UM, Parving HH. Impact of lisinopril and atenolol on kidney function in hypertensive NIDDM subjects with diabetic nephropathy. **Diabetes** 1994;43:1108-13.
- ¹³¹ Huang ES, Meigs JB, Singer DE. The effect of interventions to prevent cardiovascular disease in patients with Type 2 DM mellitus. **Am J Med** 2001;111:633-42.
- ¹³² UKPDS Group. Tight blood pressure control and risk of macrovascular and microvascular complications in Type 2 DM: UKPDS 38. **BMJ** 1998;317:703-13.
- ¹³³ Joint National Committee on Prevention, Detection, Evaluation and Treatment of High Blood Pressure: The sixth report. **Arch Intern Med** 1997;157:2413-46.
- ¹³⁴ Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomised, controlled trials. **Ann Intern Med** 2002;136:493-503.
- ¹³⁵ Kelley GA, Kelley KS, Tran ZV. Walking and resting blood pressure in adults: a meta-analysis. **Prev Med** 2001;33(Suppl 2, pt1):120-7.

- ¹³⁶ Marks JB, Raskin P. Cardiovascular risk in diabetes: a brief review. **J Diabetes Complications** 2000;14:108-15.
- ¹³⁷ Brown JB, Pedula KL, Bakst AW. The progressive cost of complications in Type 2 DM mellitus. **Arch Intern Med** 1999;159:1873-80.
- ¹³⁸ Blair S, Kampert J, Kohl H, Barlow CE, Macera CA, Paffenbarger RS, et al Influence of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. **JAMA** 1996;276:205-10.
- ¹³⁹ Knatterud G, Klimt CR, Goldner MG, et al University Group Diabetes Program. Effects of hypoglycaemic agents on vascular complications in patients with adult-onset diabetes. **JAMA** 1978;240:37-42.
- ¹⁴⁰ Abaira C, Colwell JA, Nuttal FQ, Sawin CT, Nagel NJ, Comstock JP, et al Veterans Affairs Cooperative Study on glycaemic control and complications in Type II diabetes (VA CSDM). **Diabetes Care** 1995;18:1113-23.
- ¹⁴¹ Gaede P, Vedel P, Parving H, Pedersen O. Intensified multifactorial intervention in patients with Type 2 DM mellitus and microalbuminuria: the Steno Type 2 randomised study. **Lancet** 1999;353:617-22.
- ¹⁴² Clark CM, Fradkin JE, Hiss RG, Lorenz RA, Vinicor F, Warren-Boulton E. The National Diabetes Education Program, changing the way diabetes is treated. **Diabetes Care** 2001;24:617.
- ¹⁴³ Horton ES. Role and management of exercise in diabetes Mellitus. **Diabetes Care** 1988;11:201-11.
- ¹⁴⁴ Pronk NP, Wing RR. Physical activity and long-term maintenance of weight loss. **Obes Res** 1994;2:587-99.

- ¹⁴⁵ Perri MG, McAdoo WG, McAllister DA, et al Enhancing the efficacy of behaviour therapy for obesity. **J Consult Clin Psychol** 1986;54:670-75.
- ¹⁴⁶ Di Loreto C, Parlanti N, Fanelli C, Santeusanio F, Lucidi P, Brunetti P, et al Validation of a counseling strategy to promote the adoption and the maintenance of physical activity by Type 2 diabetic subjects. **Diabetes Care** 2003;26:404-8.
- ¹⁴⁷ Dunstan DW, Burke V, Mori TA, Morton AR, Puddey IB, Stanton KG, et al The independent and combined effects of aerobic exercise and dietary fish intake on serum lipids and glycaemic control in NIDDM. **Diabetes Care** 1997;20:913-21.
- ¹⁴⁸ Wing RR, Epstein LH, Paternostro-Bayles M, et al Exercise in a behavioral weight control programme for obese patients with Type 2 (non-insulin-dependent) diabetes. **Diabetologia** 1988;31:902-9.
- ¹⁴⁹ Walker KZ, Jones JA, Piers LS, O'Dea K, Putt RS. Effects of regular walking on cardiovascular risk factors and body composition in normoglycaemic women and women with Type 2 DM. **Diabetes Care** 1999;22:555-61.
- ¹⁵⁰ Jakicic JM, Winters C, Lang W, Wing RR. Effects of intermittent exercise and use of home exercise equipment on adherence, weight loss and fitness in overweight women. **JAMA** 1999;282:1554.
- ¹⁵¹ Brown SA, Kouzekanani K, Garcia AA, Hanis CL. Culturally competent diabetes self-management education for Mexican Americans: The Starr County Border Health Initiative. **Diabetes Care** 2002;25:259-68.

- ¹⁵² Keyserling TC, Sameul-Hodge CD, Ammerman AS, Ainsworth BE, Bangdiwala SI, Henriquez-Roldan CF, et al A randomized trial of an intervention to improve self-care behaviors of African-American women with Type 2 DM. **Diabetes Care** 2002;25:1576-83.
- ¹⁵³ Goldhaber-Fiebert JD, Goldhaber-Fiebert SN, Tristan ML, Nathan DM. Randomized controlled community-based nutrition and exercise intervention improves glycaemia and cardiovascular risk factors in Type diabetic patients in rural Costa Rica. **Diabetes Care** 2003;26:24-9.
- ¹⁵⁴ US Department of Health and Human Services: Physical activity and Health: Report of the Surgeon General. Executive Summary 1-14. Atlanta, Centres for Disease Control and Prevention, 1996.
- ¹⁵⁵ Lauzon N, Tudor-Locke C, Myers A, Rodger W, Rhonda B, Chan C, et al Increased physical activity and improved health measures with a pedometer-based physical activity intervention for Type 2 DM. **Diabetes** 2003; 52:236.
- ¹⁵⁶ Brown A, Wolff K, Misra S, Elasy T. A randomized controlled trial to increase daily activity in out-patients with Type 2 DM. **Diabetes** 2003;52: 234.
- ¹⁵⁷ Dunstan DW, De Courten M, Shaw J, Zimmet P, Daly RM, Jolley D, et al High-intensity resistance training improves glycaemic control in older patients with Type 2 DM. **Diabetes Care** 2002;25:1729-36.
- ¹⁵⁸ Eeley EA, Stratton IM, Hadden DR, Turner RC, Holman RR. Estimated dietary intake in type 2 diabetic patients randomly allocated to diet, sulphonylurea or insulin therapy: UKPDS 18. **Diabet Med** 1996;13:656-62.

- ¹⁵⁹ American College of Sports Medicine: Appropriate intervention strategies for weight loss and prevention of weight regain for adults. **Med Sci Sports Exerc** 2001;33:2145-56.
- ¹⁶⁰ Rice B, Janssen I, Hudson R, Ross R. Effects of aerobic or resistance exercise and/or diet on glucose tolerance and plasma insulin levels in obese men. **Diabetes Care** 1999;22:684-91.
- ¹⁶¹ Lowther M, Mutrie N, Scott EM. Promoting physical activity in a socially and economically deprived community: a 12 month randomized control trial of fitness assessment and exercise consultation. **J Sports Sci** 2002;20:577-88.
- ¹⁶² Guyatt GH, Sullivan MJ, Thompson PJ, Fallen EL, Pugsley SO, Taylor DW, et al The 6-minute walk: a new measure of exercise capacity in patients with chronic heart failure. **Can Med Assoc J** 1985;132:919-22.
- ¹⁶³ Eales CJ, Stewart AV. A proposed field test for evaluating fitness in elderly hypertensive patients. **SA J Physiother** 1997; 53:16-9.
- ¹⁶⁴ Borg GAV. Psychophysical bases of perceived exertion. **Med Sci Sports Exerc** 1982;14:377-81.
- ¹⁶⁵ Colberg SR, Swain DP, Vinik AI. Use of Heart Rate Reserve and Rating of Perceived Exertion to prescribe exercise intensity in Diabetic Autonomic Neuropathy. **Diabetes Care** 2003;26:986-90.
- ¹⁶⁶ Samaras K, Ashwell S, Campbell LV, Chisholm DJ, Mackintosh A-M, Exercise in NIDDM: Are we missing the point. Guest Editorial. **Diabet Med** 1996;13:780-81.

- ¹⁶⁷ Pollock ML, Franklin BA, Balady GJ et al Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety and prescription: an advisory from the Committee on Exercise, Rehabilitation and Prevention, Council on Clinical Cardiology, American Heart Association. **Circulation** 2000;101:828-33.
- ¹⁶⁸ Franklin BA. Program factors that influence exercise adherence: practical adherence skills for clinical staff. In: Dishman R, editor. *Exercise Adherence: Its impact on Public Health*. Champaign, IL: Human Kinetics;1988. p.237-58.
- ¹⁶⁹ Bartlett E. Historical glimpses of patient education in the United States. **Patient Educ Counsel** 1986;8:135-49.
- ¹⁷⁰ De Weerdts I, Visser A, Van der Veen E. Attitude behavior theories and diabetes education programmes. **Patient Educ Counsel** 1989;14:3-19.
- ¹⁷¹ Roter D, Kinmonth A-L. What is the evidence that increasing participation of individuals in self-management improves the processes and outcomes of care? In: Williams R, Herman W, Kinmonth AL, Wareham NJ, editors. *The Evidence Base for Diabetes Care*. West Sussex: John Wiley & Sons; 2002. p.682-700.
- ¹⁷² Eales CJ, Stewart AV. Health and responsibility, self-efficacy, self-care and self-responsibility. **SA J Physiother** 2001;57:20-5.
- ¹⁷³ World Health Organization. Health education in self-care: possibilities and limitations. Report of a scientific consultation. Geneva, Switzerland; November 1983: 21-5.
- ¹⁷⁴ Oldridge NB, Streiner DL. The health belief model: predicting compliance and dropout in cardiac rehabilitation. **Med Sci Sports Exerc** 1990;22:678-83.

- ¹⁷⁵ Rudd J, Glanz K. How individuals use information for health action: consumer information processing. In: Eales CJ, Stewart AV. Health and responsibility, self-efficacy, self-care and self-responsibility. **SA J Physiother** 2001;57:20-5.
- ¹⁷⁶ Rickheim PL, Flader JL, Weaver TW, Kendall DM. Assessment of group versus individual diabetes education. **Diabetes Care** 2002;25:269-74.
- ¹⁷⁷ Janz NK, Becker MH. The health belief model: a decade later. **Health Educ Q** 1987;11:1-47.
- ¹⁷⁸ Perry CL, Baranowski T, Parcel GS. How individuals, environments, and health behavior interact: Social Learning Theory. In: In: Eales CJ, Stewart AV. Health and responsibility, self-efficacy, self-care and self-responsibility. **SA J Physiother** 2001;57:20-5.
- ¹⁷⁹ Bandura A. Self-efficacy: The exercise of control. New York: WH Freeman and Co; 1997. p. 198-421.
- ¹⁸⁰ Prochaska J, DiClemente C. Stages and processes of self-change of smoking: toward an integrative model of change. **J Counseling and Clinical Psychology** 1983;51:390-5.
- ¹⁸¹ Bury T, Mead J. Evidence-based Healthcare: a practical guide for therapists. Oxford: Butterworth-Heinemann; 2000. p. 66-83.
- ¹⁸² Glanz K, Lewis FM, Rimer BK, editors. Health behavior and health education: theory, research and practice. In: In: Eales CJ, Stewart AV. Health and responsibility, self-efficacy, self-care and self-responsibility. **SA J Physiother** 2001;57:20-5.

- ¹⁸³ Norris SL, Engelgau MM, Narayan KMV. Effectiveness of self-management training in Type 2 DM. **Diabetes Care** 2001;24:561-87.
- ¹⁸⁴ Norris SL, Engelgau MM, Lau J, Smith SJ, Schmid CH. Self-management education for adults with Type 2 DM: a meta-analysis of the effect on glycaemic control. **Diabetes Care** 2002;25:1159-71.
- ¹⁸⁵ Jones H, Edwards L, Vallis TM, Ruggiero L, Rossi SR, Rossi JS, et al Changes in diabetes self-care behaviors make a difference in glycemic control. **Diabetes Care** 2003;26:732-37.
- ¹⁸⁶ Egede LE. Lifestyle modification to improve blood pressure control in individuals with diabetes. **Diabetes Care** 2003;26:602-7.
- ¹⁸⁷ Kirk A, MacIntyre P, Mutrie N, Fisher M. Increasing physical activity in people with Type 2 DM. **Diabetes Care** 2003;26:1186-92.
- ¹⁸⁸ Glasgow R, Toobert D, Strycker L. Problem-solving as a key to successful long-term diabetes self-management. **Diabetes** 2003;52:413.
- ¹⁸⁹ Hill-Briggs F, Gary T, Batts-Turner M, et al Social problem solving and diabetes control in urban African Americans with Type 2 DM. **Diabetes** 2003;52:38.
- ¹⁹⁰ Kinmonth AL. Compliance and negotiation with patients with Type 2 DM. Abstract Book. Pre-EASD Symposium. 1999.
- ¹⁹¹ Glasgow RE, Hampson SE, Strycker LA, Ruggiero LA. Personal-model beliefs and social environmental barriers related to diabetes self-management. **Diabetes Care** 1997;20:556-61.

- ¹⁹² Kerner MS, Grossman AH. Scale construction for measuring attitude, beliefs, perception of control and intention to exercise. **J Sports Med Phys Fitness** 2001;41:124-31.
- ¹⁹³ Stewart AV, Eales CJ, De Charmoy S. The Knowledge Profile of Patients with Hypertension. **SA J Physiother** 2000;56:17-21.
- ¹⁹⁴ Oldridge NB, Donner AP, Buck CW ,et al Predictors of dropout from cardiac rehabilitation. Ontario Exercise Heart Collaboraqtive Study. **Am J Cardiol** 1983;51:70-4.
- ¹⁹⁵ Bopape MW. The beliefs and attitudes of patients with diabetes (non insulin dependent) in the Northern Province of South Africa. [dissertation]. University of the North: South Africa; 2000. p.1-113.
- ¹⁹⁶ Jacobson AM, Hauser ST, Anderson BJ, Polonsky W. Psychosocial aspects of Diabetes. In: Kahn CR, Weir GC, editors. *Joslin's Diabetes Mellitus*. 13th ed. Philadelphia:Lea & Febiger; 1994. p. 431-47.
- ¹⁹⁷ Aikens JE, Kiolbasa TA, Sobel R. Psychological of glycemic change with relaxation training in non-insulin-dependent Diabetes Mellitus. **Psychother Psychosom** 1997;66:302-6.
- ¹⁹⁸ Grant I, Kyle GC, Teichman A, et al Recent life events and diabetes in adults. In: Surwit RS, Schneider MS. *Role of stress in etiology and treatment of Diabetes Mellitus*. **Pschosom Med**1993;55:380-93.
- ¹⁹⁹ Surwit RS, Schneider MS. Role of stress in the etiology and treatment of Diabetes Mellitus. **Psychosom Med** 1993;55:380-93.
- ²⁰⁰ Peyrot M, McMurray JF. Psychosocial factors in diabetes control: adjustment of insulin-treated adults. **Psychosom Med** 1985;47:542-57.

- ²⁰¹ Hammond-Tooke D. *Rituals and Medicines: Indigenous Healing in South Africa*. Cape Town: AD Donker Publishers; 1989. p. 14-163.
- ²⁰² De Villiers S. Beliefs and behaviour in transcultural health care. **S Afr Fam Pract** 1991;12:44-9.
- ²⁰³ Sameul-Hodge CD, Keyserling TC, Headen SW, Jackson EJ, Skelley AH, Ammerman AS, et al Influences on day-to-day self-management of Type 2 DM among African-American women. **Diabetes Care** 2000;23:928-33.
- ²⁰⁴ De la Harpe R, De la Harpe P *Medicinal plants of South Africa*. Cape Town: Struik Publishers LTD; 1997. p. 10.
- ²⁰⁵ Snoek FJ. Quality of life: a closer look at measuring patients' well-being. **Diabetes Spectrum** 2000;13:24-31.
- ²⁰⁶ Redekop WK, Koopmanschap MA, Niessen LW, Stolk RP, Rutten GEHM, Wolffenbuttel BHR. Health-related quality of life and treatment satisfaction in Dutch patients with Type 2 DM. **Diabetes Care** 2002;25:458-63.
- ²⁰⁷ Eiser C, Tooke JE. Quality of life in Type II diabetes. **Pharmacoeconomics** 1995; 8 Suppl1:17-22.
- ²⁰⁸ Bradley C. What is the impact of treatment on quality of life in diabetes? Abstract Book . Pre-EASD symposium. 1999.
- ²⁰⁹ Petterson T, Young B, Lee P, Newton P, Hollis S, Dornan T. Well-being and treatment satisfaction in older people with diabetes. **Diabetes Care** 1998;21:930-5.
- ²¹⁰ Snoek FJ, Skinner TC. Psychological counselling in problematic diabetes: does it help? **Diabet Med** 2002;19:265-73.

- ²¹¹ Surwit RS, Feinglos MN, Van Tilburg MAL, Edwards CL, Zucker N, Williams P, et al Stress management improves long-term glycemic control in Type 2 DM. **Diabetes Care** 2002;25:30-4.
- ²¹² Lerman C, Glanz K Stress, coping and health behavior. In: Glanz K, Lewis FM, Rimer BK, eds: Health behavior and health education. 2nd ed. San Francisco: Jossey-Bass Publishers; 1997. p. 113-38.
- ²¹³ Lane JD, Feinglos MN, McCaskill CC, et al Relaxation training for NIDDM. **Diabetes Care** 1993;8:1087-93.
- ²¹⁴ Sotile WM Psychosocial interventions for cardiopulmonary patients: a guide for health professionals. Champaign: Human Kinetics; 1969. p. 97 – 101.
- ²¹⁵ Stone RA, DeLeo J. Psychotherapeutic control of hypertension. **N Engl J Med** 1976;294:80-4.
- ²¹⁶ McGrady A, Bailey BK, Good MP. Controlled study of biofeedback-assisted relaxation in Type I diabetes. **Diabetes Care** 1991;14:360-5.
- ²¹⁷ Glaser NS. Non-insulin-dependent diabetes mellitus in childhood and adolescence. **Pediatr Clin North Am** 1997;44:307-37.
- ²¹⁸ Aikens JE, Kiolbasa TA, Sobel R. Psychological of glycemic change with relaxation training in non-insulin-dependent Diabetes Mellitus. **Psychother Psychosom** 1997;66:302-6.
- ²¹⁹ Tu KS, McDaniel G, Gay JT. Diabetes self-care knowledge, behaviours and metabolic control of older adults – the effect of a post-educational follow-up program. **Diabetes Educator** 1993;19:25-30.

- ²²⁰ Ajzen I, Fishbein M. Understanding attitudes and predicting social behavior. Englewood Cliffs NJ: Prentice Hall; 1980.
- ²²¹ Anderson RM, Fitzgerald JT. The relationship between diabetes-related attitudes and patients' self-reported adherence. **Patient Educ Couns** 1993;21:41-50.
- ²²² Beeney LJ, Dunn SM, Welch G. Measurement of diabetes knowledge-the development of the DKN-scales. 2nd ed. Amsterdam: Harwood Academic Publishers; 1996. p.159-89.
- ²²³ Manson JE, Colditz G, Stampfer MJ, Willett WC, Rosner B, Monson RR et al A prospective study of obesity and risk of coronary disease in women. **N Engl J Med** 1990;322:882-9.
- ²²⁴ Westaway MS, Viljoen E, Rheeder P, Masemola T. Does blood glucose control affect the health-related quality of life (HRQOL) of urban black South African Type diabetes mellitus patients? **Diabetes Res** 2000;34:209-17.
- ²²⁵ Ligtenberg PC, Hoekstra JBL, Zonderland ML, Erkenlens DW. Physical activity and Diabetes Mellitus. **Eur J Int Med** 1995;6:95-108.
- ²²⁶ Greeff OBW. The management of Diabetes Mellitus. **Med J** 2000;42:19-25.
- ²²⁷ Nunnely JC. Psychometric theory. 2nd ed. New York: McGraw-Hill; 1978. p.230-45.
- ²²⁸ Campbell E. Behaviour change in the prevention and treatment of Non-insulin-dependent Diabetes Mellitus. In: Bradley C, editor. Handbook of Psychology and Diabetes. Amsterdam: Harwood Academic Publishers; 1996.p. 170.

- ²²⁹ Nilsson PM, Johansson SE, Sundquist J. Low educational status is a risk factor for mortality among diabetic people. **Diabet Med** 1998;15:213-19.
- ²³⁰ Beckles GLA, Engelgau MM, Narayan KMV, Herman WH, Aubert RE, Williamson DF. Population-based assessment of the level of care among adults with diabetes in the U.S.. **Diabetes Care** 1998;21:1432-37.
- ²³¹ Westaway MS, Viljoen E Rheeders P. Investigating the reliability and validity of a quality of life measure for urban black South African diabetic patients: a pilot study. **Diabetes Res** 1999;34:139-47.
- ²³² Lockington TJ, Farrant S, Meadows KA, Dowlatsahi D, Wise PH. Knowledge profile and control in diabetic patients. **Diabet Med** 1988; 5:381-6.
- ²³³ Beeney LJ, Dunn SM. Knowledge improvement and metabolic control in diabetes education : approaching the limits? **Patient Educ Counsel** 1990;16:231-45.
- ²³⁴ Cauley JA, La Porte PHR, Sandler RB, Schramm MM, Kriska AM. Comparison of methods to measure physical activity in postmenopausal women. **Am J Clin Nutr** 1987;45:14-22.
- ²³⁵ Ligtenberg PC, Hoekstra JBL, Bol E, Zonderland ML, Erkelens DW. Effects of physical training on metabolic control in elderly Type 2 DM Mellitus patients. **Clin Sci** 1997;93:127-35.
- ²³⁶ Voorrips LE, Meijers JHH, Sol P, Seidell JC Van Staveren WA. History of body weight and physical activity of elderly women differing in current physical activity. **Int J Obesity** 1992;16:199-205.

- ²³⁷ Blair SN, Applegate WB, Dunn AL, Ettinger WH, Haskell WL, King AC, et al The Activity Counselling Trial Research Group. **Med Sci Sports Exerc** 1998;1097-1106.
- ²³⁸ Hays LM, Clark DO. Correlates of physical activity in a sample of older adults with Type 2 DM. **Diabetes Care** 1999;22:706-12.
- ²³⁹ Skelley AH, Marshall JE, Haughey BP, Davis PJ, Dunford RG. Self-efficacy and confidence in outcomes as determinants of self-care practices in Inner-city African-American women with Non-insulin-dependent Diabetes. **Diabetes Educator** 1995;21:38-46.
- ²⁴⁰ Zgibor JC, Songer TJ. External barriers to Diabetes care: addressing personal and health systems issues. **Diabetes Spectrum** 2001;14:23-31.
- ²⁴¹ Orchard TJ, Dorman JS, Maser RE, Becker DJ, Ellis D, LaPorte RE, et al Factors associated with avoidance of severe complications after 25 years of IDDM: Pittsburgh Epidemiology of Diabetes Complications Study I. **Diabetes Care** 1990;13: 741-7.
- ²⁴² Tripp-Reimer T, Choi E, Kelley LS, Enslein JC. Cultural barriers of care: Inverting the problem. **Diabetes Spectrum** 2001;14:13-27.
- ²⁴³ Rheeder P, Stolk RP, Veenhouwer JF, Grobbee DE. The metabolic syndrome in black hypertensive women – waist circumference more strongly related than body mass index. **S Afr Med J** 2002;92:637-41.
- ²⁴⁴ Zgibor IC, Songer TJ, Kelsey SF, et al The association of diabetes specialist care with health care practices and glycemic control in patients with type 1 diabetes: a cross sectional analysis from the Pittsburgh Epidemiology of Diabetes Complications Study. **Diabetes Care** 2000;23:472-6.

- ²⁴⁵ Mbanya JCN, Chungong SSS, Mfopou KJ. Metabolic effects of African diets. In: Gill G, Mbanya JC, Alberti G, editors. Diabetes in Africa. Cambridge: Biddles Ltd.; 1997.p. 262.
- ²⁴⁶ Van Rooijen AJ, Rheeder P, Eales CJ, Becker PJ. Black female patients with Type 2 DM Mellitus: Knowledge, Attitudes and Physical Activity. **SA J Physiother** 2001;57:20-7.
- ²⁴⁷ Quatromoni PA, Milbauer M, Posner BM, et al Use of focus groups to explore nutrition practices and health beliefs of urban Caribbean Latinos with diabetes. **Diabetes Care** 1994;17:869-73.
- ²⁴⁸ Betts NM, Baranowski T, Hoerr SL. Recommendations for planning and reporting focus group research. **J Nutr Educ** 1996;28:279-81.
- ²⁴⁹ Côte-Arsenault D, Morrison-Beedy D. Practical advice for planning and conduction focus groups. **Nurs Res** 1999;48:280-3.
- ²⁵⁰ McDougall P. Focus groups: an overview of their use as a research method. **Community Pract** 1999;72:48-9.
- ²⁵¹ Strauss A, Corbin J. Basics of qualitative research. Newbury Park Cal; Sage Pub Inc: 1990.
- ²⁵² World Health Organization. Ageing and Physical activity: Geneva, Switzerland;1994.
- ²⁵³ Kriska A. Ethnic and cultural issues in assessing physical activity. **Res Q Exerc Sport** 2000;71:47-59.
- ²⁵⁴ Wannamethee SG, Shaper AG. Physical activity in the prevention of cardiovascular disease: an epidemiological perspective. [Review] **Sports Med** 2001;31:101-14.

- ²⁵⁵ Blair SN, Jackson AS. Physical fitness and activity as separate heart disease risk factors: a meta-analysis. **Med Sci Sports Med** 2001;33:754-61.
- ²⁵⁶ Crosson P, Zhu W, Timm G. African American Women's Exercise Barriers: a focus group study. **Res Q Exerc Sport** 2000;71:100.
- ²⁵⁷ Oldridge NB, Spencer J. Exercise habits and perceptions before and after graduating or dropout from supervised cardiac exercise rehabilitation. **J Cardiac Rehabil** 1985;5 313-19.
- ²⁵⁸ Van Rooijen AJ, Rheeder P, Eales CJ, Molatoli HM. Barriers to and expectations of performing physical activity in female subjects with Type 2 DM. **SA J Physiother** 2002;58: 3-11.
- ²⁵⁹ Rose GA, Balckburn H, Gillium RF, Prineas RJ. Cardiovascular survey methods. WHO Geneva 1982.
- ²⁶⁰ Westaway MS, Rheeder P, Van Zyl DG, Seager JR. Development and testing of a 25-item patient satisfaction scale for black South African diabetic out-patients. **Curationis** 2002;25:68-75.
- ²⁶¹ Stewart AL. The Short-Form-20 Health Survey. In: McDowell I, Newell C. Measuring health: a guide to rating scales and questionnaires. 2nd ed. New York: Oxford University Press; 1996.
- ²⁶² Bradley C, Lewis KS. Measures of psychological well-being and treatment satisfaction developed from responses of people with tablet treated diabetes. **Diabetes Medicine** 1990;7:445-51.

- ²⁶³ Bailey GJ, Weiner DA, McCabe CH, Ryan TJ. A review of common errors in the indirect measurement of blood pressure. *Archives of Internal Medicine* 1993;153:2741-48.
- ²⁶⁴ Borg GAV. Psychophysical bases of perceived exertion. *Medicine and Science in Sports and Exercise* 1982;14:377-81.
- ²⁶⁵ SYNCHRON LX. Systems Chemistry information Manual P/N 962288, 1998.
- ²⁶⁶ Ainsworth BA, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, et al Compendium of Physical activities: an update of activity codes and MET intensities *Medicine and Science in Sports and Exercise* 2000;32: (Suppl 1): 498-516.
- ²⁶⁷ Andersen RE, Wadden TA, Bartlett SJ, et al Effects of lifestyle activity vs structured aerobic exercise in obese women. *JAMA* 1999;281:335-40.
- ²⁶⁸ Pollock ML, Gaesser GA, Buthcer JD, Després, Dishman RK, Franklin BA, et al The recommended quantity and quality of exercise for developing and maintaining cardio respiratory and muscular fitness and flexibility in adults. American College of Sports Medicine Position Stand **Med Sci Sports Exerc** 1998;30:975-91.
- ²⁶⁹ Lamonte MJ, Ainsworth BE. Quantifying energy expenditure and physical activity in the context of dose response. **Med Sci Sports Exerc** 2001; 33 Suppl 1:370-8.
- ²⁷⁰ Noakes T. IMMDA Advisory Statement on Guidelines for fluid replacement during marathon running. *IAAF Technical Q* 2002;17:15-24.

- ²⁷¹ Kesaniemi AY, Danforth E JR, Jensen MD, Kopelman PG, Lefébvre P, Reeder BA. Dose-response issues concerning physical activity and health: an evidence- based symposium. *Medicine and Science in Sports and Exercise* 2001;33:Suppl,S351-S358.
- ²⁷² Cassimjee N. Expert opinion. Neuro-psychology, Dept of Psychology, University of Pretoria: 2003.
- ²⁷³ Mouton J. How to succeed in your Master's & Doctoral studies. Van Schaik Publishers, Pretoria South Africa 2003: p. 106.
- ²⁷⁴ Sonnaville JJJ, Snoek FJ, Colly LP, Deville W, Wijkel D, Heine RJ. Well-being and symptoms in relation to insulin therapy in Type 2 DM. **Diabetes Care** 1998;6:919-24.
- ²⁷⁵ Beling J. The validity and reliability of a portable heart rate monitor and carotid pulse palpation among health adults. Platform presentation. 14th International WCPT Congress 2003. Barcelona.
- ²⁷⁶ Mbanya JCN, Chungong SSS, Mfopou KJ. Metabolic effects of African diets. In: Gill G, Mbanya JC, Alberti G, editors. *Diabetes in Africa*. Cambridge: Biddles Ltd.; 1997. p.262.
- ²⁷⁷ Dubbert PM. Physical activity in women. In: National Institutes of Health: Consensus Statement: Physical Activity and cardiovascular health. Leon AS (ed) *Human Kinetics*, Champaign 1997: 204-205.
- ²⁷⁸ Wilson IB, Cleary PD. Linking clinical variables with Health-related Quality of Life. **JAMA** 1995;273:59-65.