

REFERENCES

- Amemura-Maekawa J., Mitsuhashi T. and Watanabe H. (1996). Identification of a novel periplasmic catalase-peroxidase KatA of *Legionella pneumophila*. *FEMS Microbiol. Lett.* 176 (2): 339-344.
- Abu Kwaik Y., Gao L.Y., Harb O.S. and Stone B.J. (1997). Transcriptional regulation of the macrophage-induced gene (gspA) of *Legionella pneumophila* and phenotypic characterisation of a null mutant. *Mol. Microbiol.* 24 (3): 629-642.
- Adams S.L. (1999). Legionella: A new millenium bug. *Clin. Lab. Sci.* 12 (5): 309-315.
- Adeleke A.A. (1996). Legionella-like amoebal pathogens – phylogenetic status and possible role in respiratory disease. *Emer. Infect. Dis.* 2 (3): <http://www.cdc.gov/ncidod/eid/vol2no3/adeleke.htm>
- Adeleke A.A., Fields B.S., Benson R.F., Daneshvar M.I., Pruckler J.M., Ratcliff R.M., Harrison T.G., Weyant R.S., Birtles R.J., Raoult D. and Halablab M.A. (2001). *Legionella drozanskii* sp. nov., *Legionella rowbothamii* sp. nov. and *Legionella fallonii* sp. nov., three unusual new Legionella species. *Int. J. Sys. Evol. Microbiol.* 51 (3): 1151-1160.
- Alary M. and Joly J.R. (1992). Factors contributing to the contamination of hospital water distribution systems by legionellae. *J. Infect. Dis.* 165: 565-569.
- Allen K.W., Prempeh H. and Osman M.S. (1999). Legionella pneumonia from a novel industrial aerosol. *Commun. Dis. Public Health* 2 (4): 294-296.
- Amano K. and Williams J.C. (1983). Peptidoglycan of *Legionella pneumophila*: apparent resistance to lysosome hydrolysis correlates with a high degree of peptide cross-linking. *J. Bacteriol.* 153 (1): 520-526.
- Amemura-Maekawa J., Kura F. and Watanabe H. (1996). Cloning and nucleotide sequences of iron and copper-zinc superoxide dismutase genes of *Legionella pneumophila* and their distribution among Legionella species. *Jpn. J. Med. Sci. Biol.* 49 (4): 167-186 (abstract only).

Amemura-Maekawa J., Mishima-Abe S., Kura F., Takanashi T. and Watanabe H. (1999). Identification of a novel periplasmic catalase-peroxidase KatA of *Legionella pneumophila*. *FEMS Microbiol. Lett.* 176 (2): 339-344.

Andrews H.L., Vogel J.P. and Isberg R.R. (1998). Identification of linked *Legionella pneumophila* genes essential for intracellular growth and evasion of the endocytic pathway. *Infect. Immun.* 66 (3): 950-958.

Aragon V., Kurtz S., Flieger A., Neumeister B. and Cianciotto N.P. (2000). Secreted enzymatic activities of wild-type and pilD-deficient *Legionella pneumophila*. *Inf. Imm.* 68 (4): 1855-1863.

Armon R. and Payment P. (1990). A transparent medium for isolation of *Legionella pneumophila* from environmental water sources. *J. Microbiol. Methods* 11: 65-71.

Armon R., Potasman I. and Green M. (1990). Biochemical fingerprints of *Legionella* species by the BIOLOG system: presumptive identification of clinical and environmental isolates. *Letts. Appl. Microbiol.* 11: 290-292.

ASTM Guidelines D5952 (1996). Standard guide for inspecting water systems for legionellae and investigating possible outbreaks of legionellosis (Legionnaires' disease and Pontiac fever).

Australian Standard: Waters – Examination for legionellae. Standards Australia (AS 3896-1991).

Aye T., Wachmuth K., Feeley J.C., Gibson R.J. and Johnson S.R. (1981). Plasmid profiles of *Legionella* species. *Curr. Microbiol.* 6: 389-394.

Bandyopadhyay P. and Steinman H.M. (2000). Catalase-peroxidases of *Legionella pneumophila*: cloning of the katA gene and studies of KatA function. *J. Bacteriol.* 182 (23): 6679-6686.

Bangsborg J.M., Collins M.T. and Høiby N. (1986). Antigenic homogeneity among *Legionella pneumophila* serogroups 1 to 6 evaluated by crossed immunoelectrophoresis. *Int. J. Syst. Bacteriol.* 36 (3): 399-404.

Barker J., Lambers P.A. and Brown M.R. (1992). Influence of intra-amoebic and other

Barbaree J.M. (1993). Selecting a subtyping technique for use in investigations of legionellosis epidemics. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). *Legionella. Current status and emerging perspectives*. American Society for Microbiology, Washington, DC.

Suppl. 1: 24-31.

Barbaree J.M., Fields B.S., Feeley J.C., Gorman G.W. and Martin W.T. (1986). Isolation of protozoa from water associated with a legionellosis outbreak and demonstration of intracellular multiplication of *Legionella pneumophila*. *Appl. Environ. Microbiol.* 51 (2): 422-424.

Bej A.K., Mahubani M.H., Miller R., DeCesare J.L., Herf J. and Atlas R.M. (1991)

Barbaree J.M., Gorman G.W., Martin W.T., Fields B.S. and Morrill W.E. (1987). Protocol for sampling environmental sites for legionellae. *Appl. Environ. Microbiol.* 53 (7): 1454-1458.

Bej A.K., Mahubani M.H. and Atlas R.M. (1991). Detection of legionellae

Barker J., Brown M.R.W., Collier P.J., Farrell I. and Gilbert P. (1992). Relationship between *Legionella pneumophila* and *Acanthamoeba polyphaga*: physiological status and susceptibility to chemical inactivation. *Appl. Environ. Microbiol.* 58 (8): 2420-2425.

Bej A.K., Mahubani M.H., DeCesare J.L. and Atlas R.M. (1991). Polymerase chain

Barker J., Brown M.R.W., Gilbert P., Collier P.J. and Farrell I.D. (1993). The physiological status of *Legionella pneumophila* and its susceptibility to chemical inactivation. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). *Legionella. Current status and emerging perspectives*. American Society for Microbiology, Washington, DC.

of *Legionella pneumophila* by polymerase chain reaction amplification and restriction

Barker J. and Brown M.R. (1995). Speculations on the influence of infecting phenotype on virulence and antibiotic susceptibility of *Legionella pneumophila*. *J. Antimicrob. Chemother.* 36 (1): 7-21.

Barker J. and Clement P. (1994). Reducing the risk of legionellosis: the role of amoebae. *Occup. Health. Rev.* Jan/Feb. 20-23.

Appl. Environ. Microbiol. 57: 2626-2630.

Barker J., Farrell I.D. and Hutchinson J.G. (1986). Factors affecting growth of *Legionella pneumophila* in liquid media. *J. Med. Microbiol.* 22 (2): 97-100.

Barker J., Lambert P.A. and Brown M.R. (1993). Influence of intra-amoebic and other growth conditions on the surface properties of *Legionella pneumophila*. *Infect. Immun.* 61 (8): 3503-3510.

Bartlett J. (2000). Treatment of community-acquired pneumonia. *Chemotherapy* 46: Suppl. 1: 24-31.

Bartie C. and Klugman K.P. (1997). Exposures to *Legionella pneumophila* and *Chlamydia pneumoniae* in South African mine workers. *Int. J. Occup. Env. Health* 3 (2): 120-127.

Bej A.K., Mahbubani M.H., Miller R., DiCesare J.L., Haff L. and Atlas R.M. (1990). Multiplex PCR amplification and immobilised capture probes for detection of bacterial pathogens and indicators in water. *Mol. Cell. Probes* 4: 353-365.

Bej A.K., Mahbubani M.H. and Atlas R.M. (1991). Detection of viable *Legionella pneumophila* in water by polymerase chain reaction and gene probe methods. *Appl. Environ. Microbiol.* 57 (2): 597-600.

Bej A.K., Mahbubani M.H., DiCesare J.L. and Atlas R.M. (1991). Polymerase chain reaction-gene probe detection of microorganisms by using filter-concentrated samples. *Appl. Environ. Microbiol.* 57 (12): 3529-3534.

Bej A.K., Mahbubani M.H. and Atlas R.M. (1993). Detection and molecular serogrouping of *Legionella pneumophila* by polymerase chain reaction amplification and restriction enzyme analysis. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). *Legionella. Current status and emerging perspectives*. American Society for Microbiology, Washington, DC.

Bej A.K., Mahbubani M.H., DiCesare J.L. and Atlas R.M. (1991). Polymerase chain reaction-gene probe detection of microorganisms by using filter-concentrated samples. *Appl. Environ. Microbiol.* 57: 3529-3534.

- Benson R.F. and Fields B.C. (1998). Classification of the genus *Legionella*. *Semin. Respir. Infect.* 13 (2): 90-99.
- * Benson R.F., Fields B.S., Besser R. and Benin A. (2000). Application of amplified fragment length polymorphism to subtyping of *Legionella pneumophila* SG1. Presented at the 5th International Conference on Legionella, September 26-29, Neu-Ulm, Germany.
- Bentham R.H. (2000). Routine sampling and the control of *Legionella* species in cooling water systems. *Curr. Microbiol.* 41 (4): 271-275.
- Bohach G.A. and Snyder I.S. (1983). Cyanobacterial stimulation of growth and oxygen uptake by *Legionella pneumophila*. *Appl. Environ. Microbiol.* 46 (2): 528-531.
- Boissinot M., Ramsay D., Barthe C. and Joly J.R. (1987). Antigenic variability of the outer membrane antigens of *Legionella pneumophila* SGs 1-8. *Can. J. Microbiol.* 33 (7): 607-613.
- * Bollin G.E., Plouffe J.F., Para M.F. and Hackman B. (1985). Aerosols containing *Legionella pneumophila* generated by showerhead and hot-water faucets. *Appl. Env. Microbiol.* 50 (5): 1128-1131.
- Bopp C.A., Sumner J.W., Morris G.K. and Wells J.G. (1981). Isolation of *Legionella* species from environmental water samples by low-pH treatment and use of a selective medium. *J. Clin. Microbiol.* 13 (4): 714-719.
- Boulanger C.A. and Edelstein P.H. (1995). Precision and accuracy of recovery of *Legionella pneumophila* from seeded tap water by filtration and centrifugation. *Appl. Environ. Microbiol.* 61: 1805-1809.
- Brand B.C., Sadosky A.B. and Shuman H.A. (1994). The *Legionella pneumophila* icm locus: a set of genes required for intracellular multiplication in human macrophages. *Mol. Microbiol.* 14 (4): 797-808.
- * Broadbent O.H. (1987). Practical measures to control Legionnaires' disease hazards. *Australian Refrigeration, Air Conditioning and Heating*, July 22-30.

Brenner D.J., Feeley J.C. and Weaver R.E. (1984). Family VII. Legionellaceae, p279. In NR Krieg and JG Wolf (eds). *Bergey's Manual of systematic bacteriology*. Vol 1. Williams and Wilkins, Baltimore.

Brenner D.J., Steigerwalt A.G. and McDade J.E. (1979). Classification of the Legionnaires' disease bacterium: *Legionella pneumophila*, genus novum, species nova, of the family Legionellaceae, family nova. *Ann. Intern. Med.* 90 (4): 656-658.

Brenner D.J. (1986). Classification of Legionellaceae. *Isr. J. Med. Sci.* 22 (9): 620-633.

* Brenner D.J. (1987). Classification of the legionellae. *Semin. Respir. Infect.* 2 (4): 190-205.

Brenner D.J., Steigerwalt A.G., Epple P., Bibb W.F., McKinney R.M., Starnes R.W., Colville J.M., Selander R.K., Edelstein P.H. and Moss C.W. (1988). *Legionella pneumophila* SG Lansing 3 isolated from a patient with fatal pneumonia, and descriptions of *L. pneumophila* subsp. *pneumophila* subsp. nov., *L. pneumophila* subsp. *fraseri* subsp. nov. and *L. pneumophila* subsp. *pascullei* subsp. nov. *J. Clin. Microbiol.* 26 (9): 1695-1703.

Breiman R.F. (1993). Modes of transmission in epidemic and non-epidemic *Legionella* infections: directions for further study. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology, Washington, DC.

* Breiman R.F., Fields B.S., Sanden G.N., Volmer L., Meier A. and Spika J.S. (1990). Association of shower use with Legionnaires' disease: possible role of amoebae. *J. Am. Med. Assoc.* 2924 (3):

Brindle R.J., Stannett P.J. and Cunliffe R.N. (1987). *Legionella pneumophila*: comparison of isolation from water specimens by centrifugation and filtration. *Epidem. Inf.* 99: 241-247.

Broadbent C.R. (1987). Practical measures to control Legionnaires' disease hazards. *Australian Refrigeration, Air Conditioning and Heating*, July: 22-30.

Clondeston N.R., Bengtson J.M., Eisenstein B.I. and Inglesby N.G. (1999).

* Brown C.M., Nuorti P.J., Breiman R.F., Hathcock A.L., Fields B.S., Lipman H.B., Llewellyn G.C., Hofman J. and Detron M. (1999). A community outbreak of Legionnaires' disease linked to hospital cooling towers: an epidemiological method to calculate dose of exposure. *Int. J. Epidemiol.* 28: 353-359.

Brown A., Vickers R.M., Elder E.M., Lema M. and Garrity G.M. (1982). Plasmid and surface antigen markers of endemic and epidemic *Legionella pneumophila* strains. *J. Clin. Microbiol.* 16 (2): 230-235.

Brundett G. (1989). Water Services and Maintenance Systems. *Occup Health* June: 159-161.

Butler C.A. and Hoffman P.S. (1990). Characterisation of a major 31-kilodalton peptidoglycan-bound protein of *Legionella pneumophila*. *J. Bacteriol.* 172 (5): 2401-2407.

* Butler C.A., Street E.D., Hatch T.P. and Hoffman P.S. (1985). Disulfide-bonded outer membrane proteins in the genus *Legionella*. *Infect. Immun.* 48 (1): 14-18.

Byrd T.F. and Horwitz M.A. (2000). Aberrantly low transferrin receptor expression on human monocytes is associated with non-permissiveness for *Legionella pneumophila* growth. *J. Infect. Dis.* 181 (4): 1394-1400.

Byrd J.J., Xu H.S. and Colwell R.R. (1991). Viable but non-culturable bacteria in drinking water. *Appl. Env. Microbiol.* 57 (3): 875-878.

Chandler F.W., Cole R.M., Hicklin M.D., Blackmon J.A. and Callaway C.S. (1979). Ultrastructure of the Legionnaires' disease bacterium. *Ann. Intern. Med.* 90 (4): 642-647.

* Chandler F.W., Roth I.L., Callaway C.S., Bump J.L., Thomason B.M. and Weaver R.E. (1980). Flagella on Legionnaires' disease bacteria. *Ann. Intern. Med.* 93: 711-714.

Cianciotto N.P., Bangsberg J.M., Eisenstein B.I. and Engleberg N.C. (1990). Identification of mip-like genes in the genus *Legionella*. *Infect. Immun.* 58: 2912-2919.

spacer length polymorphism. *J. Clin. Microbiol.* 39 (1): 162-169.

Cianciotto N.P. and Fields B.S. (1992). *Legionella pneumophila* mip gene potentiates intracellular infection of protozoa and human macrophages.

Distribution and regulation of the *Legionella* mip gene. In: Barbara J.M., Braiman R.F. and DuFour A.P. (eds) *Legionella: Current status and emerging*

Cianciotto (2001). Pathogenicity of *Legionella pneumophila*. *Int. J. Med. Microbiol.* 291 (5): 331-343.

Comparison of isolation methods

Ciecielski C.A., Blaser M.J. and Wang W.L. (1986). Serogroup specificity of *Legionella pneumophila* is related to lipopolysaccharide characteristics. *Infect. Immun.* 51 (2): 397-404.

The serological

Cirillo S.L., Bermudez L.E., El-Etr S.H., Duhamel G.E. and Cirillo J.D. (2001). *Legionella pneumophila* entry gene rtxA is involved in virulence. *Infect. Immun.* 69 (1): 508-517.

Coers J., Kagan J.C., Matthews M., Nagai H., Zuckman D.M. and Roy C.R. (2000). Identification of icm protein complexes that play distinct roles in the biogenesis of an organelle permissive for *Legionella pneumophila* intracellular growth. *Mol. Microbiol.* 38 (4): 719-736.

Characterization of body and

Colbourne J.S. and Dennis P.J. (1985). Distribution and persistence of *Legionella* in water systems. *Microbiol. Sc.* 2 (2): 40-43.

Dennis P.J., Dall N., Miles L., Pedrosa P., Hernandez A. and Poole C. (1987)

Colbourne J.S. and Dennis P.J. (1989). The ecology and survival of *Legionella pneumophila*. *J. IWEM.* 3:345-350.

J. Infect. Dis. 16 (12): 336-339.

Collins M.T., Cho S.N., Høiby N., Espersen F., Baek L. and Reif J.S. (1983). Crossed immunoelectrophoretic analysis of *Legionella pneumophila* SG1 antigens. *Inf. Immun.* 39 (3): 1428-1440.

Water quality - detection and

Craeger J.G., Black J.G. and Davison V.E. (eds). (1990). *Microbiology: Principles and Applications*. Prentice Hall, Englewood Cliffs, New Jersey 07632.

Dennis P.J., O'Connell W. and Cianciotto N.P. (1993). Distribution and regulation of

the *Legionella* mip gene. In: Barbara J.M., Braiman R.F. and DuFour A.P. (eds)

De Gheldre Y., Maes N., Presti F.L., Etienne J. and Struelens M. (2001). Rapid identification of clinically relevant legionella spp. by analysis of transfer DNA intergenic spacer length polymorphism. *J. Clin. Microbiol.* 39 (1): 162-169.

Dennis P.J. (1993). Potable water systems: insights into control. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). Legionella: current status and emerging perspectives. American Society for Microbiology. Washington, DC.

Dennis P.J., Bartlett C.L.R. and Wright A.E. (1993). Comparison of isolation methods for Legionella species. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). Legionella. Current status and emerging perspectives. American Society for Microbiology, Washington, DC.

Desai R., Welsh C., Summy M., Farone M. and Newsome A.L. (1999). The potential of in situ hybridisation and an immunogold assay to identify Legionella associations with other microorganisms. *J. Microbiol. Methods* 37 (2): 155-164.

Dietrich C., Heuner K., Brand B.C., Hacker J. and Steinert M. (2001). Flagellum of *Legionella pneumophila* positively affects the early phase of infection in eukaryotic host cells. *Infect. Immun.* 69 (4): 2116-2122.

Diogo A., Veríssimo A., Nobre M.F. and Da Costa M.S. (1999). Usefulness of fatty acid composition for differentiation of Legionella species. *J. Clin. Microbiol.* 37 (7): 2248-2254.

Dominguez J., Gali N., Matas L., Pedroso P., Hernandez A. and Padilla E. (1999). Evaluation of a rapid immunochromatographic assay for the detection of Legionella antigen in urine. *Eur. J. Clin. Microbiol. Infect. Dis.* 18 (12): 896-898.

Draft British Standard for Identification of Legionella from environmental samples (1992). DD 211:1992.

Draft International Standard ISO/DIS 11731 (1996). Water quality – detection and enumeration of Legionella.

Dumais-Pope C., O'Connell W. and Cianciotto N.P. (1993). Distribution and regulation of the Legionella mip gene. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds).

Legionella: Current Status and Emerging Perspectives. American Society for Microbiology. Washington, DC.

Edelstein P.H. (1981). Improved semiselective medium for isolation of *L pneumophila* from contaminated clinical and environmental specimens. *J. Clin. Microbiol.* 14 (3): 298-303.

Edelstein P.H. (1982). Comparative study of selective media for isolation of *Legionella pneumophila* from potable water. *J. Clin. Microbiol.* 16 (4): 697-699.

Edelstein P.H., Edelstein M.A., Higa F. and Falkow S. (1999). *Proc. Natl. Acad. Sci. USA* 96 (14): 190-195.

Ehret W., Von Specht B.U. and Ruckdeschel G. (1986). Discrimination between clinical and environmental strains of *Legionella pneumophila* by a monoclonal antibody. *Isr. J. Med. Sci.* 22 (10): 715-723.

Ehret W. and Ruckdeschel G. (1985). Membrane proteins of Legionellaceae. I. Membrane proteins of different strains and serogroups of *Legionella pneumophila*. *Zentralbl. Bakteriol. Mikrobiol. Hyg.* 259 (4): 433-445.

Ehret W., Anding G., Tartakovski I. and Ruckdeschel G. (1993). Molecular epidemiology of outbreak-associated Serogroup 1 isolates. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). *Legionella: current status and emerging perspectives*. American Society for Microbiology. Washington, DC.

Ehret W., Von Specht B.U. and Ruckdeschel G. (1986). Discrimination between clinical and environmental strains of *Legionella pneumophila* by a monoclonal antibody. *Isr. J. Med. Sci.* 22 (10): 715-723

Engelhart S., Sennekamp J., Gilges S., Pleischl S. and Exner M. (2000). Workplace-related complaints due to exposure to contaminated humidifier water and the VDI guideline 6022. *Eur. J. Med. Res.* 5 (3): 126.

England A.C., Fraser D.W., Plikaytis B.D., Tsai T.F., Storch G. and Broome C.V. (1981). Sporadic legionellosis in the United States: the first thousand cases. *Am. J. Med.* 70: 707-711.

Engleberg N.C., Howe D.C., Rogers J.E., Arroyo J. and Eisenstein B.I. (1991). Characterisation of a *Legionella pneumophila* gene encoding a lipoprotein antigen. *Mol. Microbiol.* 5 (8): 2021-2029.

Engleberg N.C. (1993). Genetic studies of Legionella pathogenesis. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). Legionella: Current Status and Emerging Perspectives. American Society for Microbiology. Washington, DC. Pp 63-68.

Feeley J.C., Gibson R.J., Gorman G.W., Langford N.C., Rasheed J.K., Mackel D.C. and Baine W.B. (1979). Charcoal yeast extract agar: primary isolation medium for *Legionella pneumophila*. *J. Clin. Microbiol.* 10 (4): 437-441.

Feeley J.C., Gorman G.W., Weaver R.E., Mackel D.C. and Smith H.W. (1978). Primary isolation media for Legionnaires' disease bacterium. *J. Clin. Microbiol.* 8 (3): 320-325.

Fields B.S. (1991). The role of amoebae in legionellosis. *Clin. Microbiol. News.* 13 (12): 92-93 (editorial).

Fields B.S. (1996). The molecular ecology of legionellae. *Trends Microbiol.* 4 (7): 286-290.

Fields B.S. (1997). Legionella and Legionnaires' disease, p 666-675. In: Walter M.V., Knudsen G.R., McInerney M.J., Stetzenbach L.D. and Hurst C.J. (eds). Manual of environmental microbiology. American Society for Microbiology, Washington, DC.

Fields B.S., Sanden G.N., Barbaree J.M., Morrill W.E., Wadowski R.M., White E.H. and Feeley J.C. (1989). Intracellular multiplication of *Legionella pneumophila* in amoebae isolated from hospital hot water tanks. *Curr. Microbiol.* 18: 131-137.

Finch R. (1988). Minimising the risk of Legionnaires' disease. *British Med. J.* 296: 1343-1344.

Fiore A.E., Butler J.C., Emori T.G. and Gaynes R.P. (1999). A survey of methods used to detect nosocomial legionellosis among participants in the National Nosocomial Infections Surveillance System. *Infect. Control Hosp. Epidemiol.* 20 (6): 412-416.

Flieger A., Gong S., Faigle M., Deeg M., Bartmann P. and Neumeister B. (2000). Novel phospholipase A activity secreted by *Legionella* species. *J. Bacteriol.* 182 (5): 1321-1327.

Fouchet P., Jayat C., Héchard Y., Ratinaud M.H. and Frelat G. (1993). Recent advances of flow cytometry in fundamental and applied microbiology. *Biol. Cell* 78: 95-109.

Freije M.R. (1996). Legionellae control in health care facilities. A guide for minimising risk. HC Information Resources, Inc., Indianapolis, USA.

Gabay J.E., Blake M., Niles W.D. and Horwitz M.A. (1985). Purification of *Legionella pneumophila* major outer membrane protein and demonstration that it is a porin. *J. Bacteriol.* 162 (1): 85-91.

Gabay J.E. and Horwitz M.A. (1985). Isolation and characterization of the cytoplasmic and outer membranes of the Legionnaires' disease bacterium (*Legionella pneumophila*). *J. Exp. Med.* 161: 409-422.

Girod J.C., Reichman R.C., Winn W.C., Laucke D.N., Vogt R.L. and Dolin R. (1982). Pneumonic and non-pneumonic forms of legionellosis: the result of a common-source exposure to *Legionella pneumophila*. *Arch. Intern. Med.* 142: 545-547.

Glick T.H., Gregg M.B., Berman B., Mallison G., Rhodes W.W. and Kassanoff I. (1978). Pontiac fever: an epidemic of unknown etiology in a health department: I. Clinical and epidemiological aspects. *Am. J. Epidemiol.* 107 (2): 149-160.

Goldstein B., Cole L.R., Koornhof H.J. and Dubery B. (1980). A fatal case of Legionnaires' disease. *SA Mediese Tydskrif* 58: 24-27.

Gorman G.W., Feeley J.C., Steigerwalt A., Edelstein P.H., Moss C.W. and Brenner D.J. (1985). *Legionella anisa*: a new species of Legionella isolated from potable waters and a cooling tower. *Appl. Environ. Microbiol.* 49 (2): 305-309.

Gosting L.H., Cabrian K., Sturge J.C. and Goldstein L.C. (1984). Identification of a species-specific antigen in *Legionella pneumophila* by a monoclonal antibody. *J. Clin. Microbiol.* 20 (6): 1031-1035.

Grabow N.A., Kfir R. and Grabow W.O.K. (1991). A most probable number method for the enumeration of Legionella bacteria in water. *Wat. Sci. Tech.* 24 (2): 143-147.

Grabow N.A., Pienaar E.J. and Kfir R. (1991). The occurrence of Legionella bacteria in cooling towers in South Africa. *Wat. Sci. Tech.* 24 (2): 149-152.

Graman P.S., Quinlan G.A., Rank J.A. (1997). Nosocomial legionellosis traced to a contaminated ice machine. *Infect. Control Hosp. Epidemiol.* 18 (9): 637-640.

Gregersen P., Grunnet K., Uldum S.A., Andersen B.H. and Madsen H. (1999). Pontiac fever at a sewage treatment plant in the food industry. *Scand. J. Work Environ. Health* 25 (3): 291-295.

Grimm D., Ludwig W., Brand B.C., Schleifer K.H., Michel R., Hacker J. and Steinert M. (2000). Specific detection of Legionella and its amoebic hosts by fluorescence in situ hybridisation with 16S and 18S rRNA oligonucleotide probes. Presented at the 5th International Conference on Legionella. September 26-29, Neu-Ulm, Germany.

Grimont P.A.D., Grimont F., Desplaces N. and Tchen P. (1985). DNA probe specific for *Legionella pneumophila*. *J. Clin. Microbiol.* 2: 431-437.

Grimont F., Lefevre M., Ageron E., and Grimont P.A.D. (1989). rRNA gene restriction patterns of Legionella species: a molecular identification system. *Res. Microbiol.* 140: 615-626.

Hébert G.A., Dalloway C.B. and Ewing E.P. Jr (1984). Comparison of Legionella pneumophila, L. micdadei, L. bozemanii and L. dumoffii by transmission electron microscopy. *J. Clin. Microbiol.* 19 (2): 116-121.

- Groothuis D.G. and Veenendal H.R. (1983). Heat treatment as an aid for the isolation of *Legionella pneumophila* from clinical and environmental samples. *Zentralbl. Bakteriol. Mikrobiol. Hyg.* 255 (1): 39-43.
- Hacker J., Ott M., Wintermeyer E., Ludwig B. and Fischer G. (1993). Analysis of virulence factors of *Legionella pneumophila*. *Int. J. Med. Microbiol. Parasitol. Infect. Dis.* 278 (2-3):348-358.
- Hales L.M. and Shuman H.A. (1999). The *Legionella pneumophila* rpoS gene is required for growth within *Acanthamoeba castellanii*. *J. Bacteriol.* 181 (16): 4879-4889.
- Hales L.M. and Shuman H.A. (1999). *Legionella pneumophila* contains a type II general secretion pathway required for growth in amoebae as well as for secretion of the Msp protease. *Infect. Immun.* 67 (7): 3662-3666.
- Haley C.E., Cohen M.L., Halter J. and Meyer R.D. (1979). Nosocomial Legionnaires' disease: a continuing common-source epidemic at Wadsworth Medical Centre. *Ann. Intern. Med.* 90 (4): 583-586.
- Harb O.S. and Abu Kwaik Y. (2000). Interaction of *Legionella pneumophila* with protozoa provides lessons. *ASM News* 66 (10): 609-616.
- Hariparsad D., Ramsaroop R., Pather M. and Wesley A. Legionnaires' disease in Durban. A case report. *SA Mediese Tydskrif* 59: 977-979.
- Hart I., Serman S., Lee P.A. and Lee J.V. (2000). Evaluation of the Oxoid Dryspot Legionella test range. Presented at the 5th International Conference on Legionella, September 26-29, Neu-Ulm, Germany.
- Hébert G.A. (1981). Hippurate hydrolysis by *Legionella pneumophila*. *J. Clin. Microbiol.* 13 (1): 240-242.
- Hébert G.A., Callaway C.S. and Ewing E.P. Jr (1984). Comparison of *Legionella pneumophila*, *L. micdadei*, *L. bozemanii* and *L. dumoffii* by transmission electron microscopy. *J. Clin. Microbiol.* 19 (2): 116-121.

Helbig J.H., Luck P.C., Steinert M., Jacobs E. and Witt M. (2001). Immunolocalisation of the Mip protein of intracellularly and extracellularly grown *Legionella pneumophila*. *Letts. Appl. Microbiol.* 32 (2): 83-88.

Hellard M.E., Sinclair M.I., Streeton C.L. and Fairly C.K. (1997). Commentary: Drinking water and microbial pathogens – issues and challenges for the year 2000. *J. Pub. Health Med.* 19 (2): 129-131.

Heuner K., Brand B.C. and Hacker J. (1999). The expression of the flagellum of *Legionella pneumophila* is modulated by different environmental factors. *FEMS Microbiol. Lett.* 175 (1): 69-77.

Heuner K., Dietrich C., Steinert M., Gobel U.B. and Hacker J. (2000). Cloning and characterisation of the *Legionella pneumophila*-specific gene encoding a member of the LysR family of transcriptional regulators. *Mol. Gen. Genet.* 264 (1-2): 204-211.

Heymann D.L. (2001). Strengthening global preparedness for defence against infectious disease threats. Committee on Foreign Relations, United States Senate. Hearing on *The threat of bioterrorism and the spread of infectious diseases*, 5 September.

Higa F. and Edelstein P.H. (2001). Potential virulence role of *Legionella pneumophila* ptsP ortholog. *Infect. Immun.* 69 (8): 4782-4789.

Hindahl M.S. and Iglewski B.H. (1984). Isolation and characterisation of *Legionella pneumophila* outer membrane. *J. Bacteriol.* 159 (1): 107-113.

Hindahl M.S. and Iglewski B.H. (1986). Outer membrane proteins from *Legionella pneumophila* serogroups and other *Legionella* species. *Infect. Immun.* 51 (1): 94-101.

Hindahl M.S. and Iglewski B.H. (1987). Cloning and expression of a common *Legionella* outer membrane antigen in *Escherichia coli*. *Microb. Pathog.* 2 (2): 91-99.

Hobbie J.E., Daley R.J. and Jasper S. (1977). Use of nucleopore filters for counting bacteria by fluorescence microscopy. *Appl. Environ. Microbiol.* 33 (5): 1225-1228.

Hoffman P. (1984). Bacterial physiology. In: *Legionella, Proceedings of the 2nd International Symposium* ed. Thornsberry C, Balows A, Feeley JC and Jakubowski W: pp 61-67. Washington DC: American Society for Microbiology.

Hoffman P.S. and Garduno R.A. (1999). Surface-associated heat shock proteins of *Legionella pneumophila* and *Helicobacter pylori*: roles in pathogenesis and immunity. *Infect. Dis. Obstet. Gynecol.* 7 (1-2): 58-63 (abstract only)

Hoffman P.S., Pine L. and Bell S. (1983). Production of superoxide and hydrogen peroxide in medium used to culture *Legionella pneumophila*: catalytic decomposition of charcoal. *Appl. Env. Microbiol.* 45 (3): 784-791.

Hoffman P.S., Ripley M. and Weeratna R. (1992a). Cloning and nucleotide sequence of a gene (OmpS) encoding the major outer membrane protein of *Legionella pneumophila*. *J. Bacteriol.* 174 (3): 914-920.

Hoffman P.S., Seyer J.H. and Butler C.A. (1992b). Molecular characterisation of the 28- and 31-kilodalton subunits of the *Legionella pneumophila* major outer membrane protein. *J. Bacteriol.* 174 (3): 908-913.

Hübner I., Obst U., Steinmetz I. and Bitter-Suermann D. (1992). Microbiological-immunological rapid method to detect legionellae in water. *Acta Hydrochim. Hydrobiol.* 20: 205-207.

Hughes M.A. and Steele T.W. (1994). Occurrence and distribution of *Legionella* species in composted plant materials. *Appl. Environ. Microbiol.* 60 (6): 2003-2005.

Hurley M.C., Balazovich K., Albano M., Engleberg N.C. and Eisenstein B.I. (1993). *Legionella pneumophila* Mip inhibits protein kinase C. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp 69-70.

Joy J.R., McKinney K.M., Train J.O., Blue W.F., Watkins J.D. and Ramsay D. (1988). Recovery of *Legionella* species from water samples by four different methods. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp. 200-201

Hussong D., Colwell R.R., O'Brien M., Weiss E., Pearson A.D., Weiner R.M. and Burge W.D. (1987). Viable *Legionella pneumophila* not detectable by culture on agar media. *Biotechnology* 5: 947-950.

*(1980). Legionnaires' disease in Johannesburg. *South African Med-J* 52: 13-17.

*Ikedo M. and Yabuuchi E. (1986). Ecological studies of *Legionella* species. I. Viable counts of *Legionella pneumophila* in coolant tower water. *Microbiol. Immunol.* 30 (5): 413-423.

*(1986). Serogroup 1 infection with the Direct Enzyme Immunosorbent Assay (DEIA) test. *J. Clin. Microbiol.* 25 (4): 854-856.

Inglis (2000).

Kassir C.W. (2002). *Legionella and Legionella*. Presented at the 37th International

Ishimatsu S., Miyamoto H., Hori H., Tanakas I. and Yoshida S. (2001). Sampling and detection of *Legionella pneumophila* aerosols generated from an industrial cooling tower. *Ann. Occup. Hyg.* 45 (6): 421-427.

*(1996). *Legionella and the environment*. *Proc. 9th Int. Conf. on Legionnaires' Disease* 17 (2): 133-181.

Jantzen E., Sonesson A., Tangen T. and Eng J. (1993). Hydroxy-fatty acid profiles of *Legionella* species: diagnostic usefulness assessed by principal component analysis. *J. Clin. Microbiol.* 31 (6): 1413-1419.

*(1996). Community outbreak of Legionnaires' disease: an investigation confirming the potential for cooling towers to

*Johnson W., Elliot J.A., Helms C.M. and Renner E.D. (1979). A high molecular weight antigen in the Legionnaires' disease bacterium: isolation and partial characterisation. *Ann. Intern. Med.* 90 (4): 638-641.

*(1986). Survival of *Legionella pneumophila* within cysts of *Acanthamoeba polyphaga* following chlorine treatment. *J. Appl. Bacteriol.* 60 (5): 11-20.

Joly J.R., Alary M. (1993). Occurrence of nosocomial Legionnaires' disease in hospitals with contaminated potable water supply. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC.

Kinoshita Y.A., Mol H. and Zähringer U. (1995). Structural study of a highly O-acetylated

Joly J.R., McKinney R.M., Tobin J.O., Bibb W.F., Watkins I.D. and Ramsay D. (1986). Development of a standardised subgrouping scheme for *Legionella pneumophila* serogroup 1 using monoclonal antibodies. *J. Clin. Microbiol.* 23: 768-771.

Kinoshita C., High A.S. and Rogers J.G. (1993). Role of the 25kDa major outer membrane

Jousimies-Somer H.R., Waarala S. and Väisänen M.L. (1993). Recovery of *Legionella* species from water samples by four different methods. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp. 200-201.

Krudson G.B. and Milsted F. (1973). A plasmid in *Legionella pneumophila*. *J. Bacteriol.* 25 (1): 1062-1063.

Kaplan C., Zwi S., Kallenbach J., Singer J., Koornhof H.J., Mauff A.C. and Dubery B. (1980). Legionnaires' disease in Johannesburg. *South African Med. J.* 58: 13-17.

Kazandjian D., Chiew R. and Gilbert G.L. (1997). Rapid diagnosis of *Legionella pneumophila* Serogroup 1 infection with the Binax Enzyme Immunoassay urinary antigen test. *J. Clin. Microbiol.* 35 (4): 954-956.

Keevil C.W. (2000). Biofilms and Legionella. Presented at the 5th International Conference on Legionella, September 26-29, Neu-Ulm, Germany.

Keleti G. and Shapiro M.A. (1987). Legionella and the environment. *CRC Crit. Rev. Env. Control* 17 (2): 133-181.

Keller D.W., Hajjeh R., DeMaria A., Fields B.S., Pruckler J.M., Benson R.S., Kludt P.E., Lett S.M., Mermel L.A., Giorgio C. and Breiman R.F. (1996). Community outbreak of Legionnaires' disease: an investigation confirming the potential for cooling towers to transmit Legionella species. *Clin. Infect. Dis.* 22 (2): 257-261.

Kilvington S. and Price J. (1990). Survival of *Legionella pneumophila* within cysts of *Acanthamoeba polyphaga* following chlorine exposure. *J. Appl. Bacteriol.* 68 (5): 19-25.

Kirby J.E., Vogel J.P., Andrews H.L. and Isberg R.R. (1998). Evidence of pore-forming ability by *Legionella pneumophila*. *Mol. Microbiol.* 27 (2): 323-336.

Knirel Y.A., Moll H. and Zähringer U. (1996). Structural study of a highly O-acetylated core of *Legionella pneumophila* serogroup 1 lipopolysaccharide. *Carbohydr. Res.* 293 (2): 223-234.

Kniros C., High A.S. and Rodgers F.G. (1999). Role of the 25kDa major outer membrane protein of *Legionella pneumophila* in attachment to U-937 cells and its potential as a virulence factor for chick embryos. *J. Appl. Microbiol.* 86 (2): 237-244.

Knudson G.B. and Mikesell P. (1980). A plasmid in *Legionella pneumophila*. *Infect. Immun.* 29 (1): 1092-1095.

Kool J.L., Bergmire-Sweat D., Butler J.C., Brown E.W., Peabody D.J., Massi D.S., Carpenter J.C., Pruckler J.M., Benson R.F. and Fields B.S. (1999). Hospital characteristics associated with colonisation of water systems by Legionella and risk of nosocomial Legionnaires' disease: a cohort study of 15 hospitals. *Infect. Control Hosp. Epidemiol.* 20 (12): 798-805.

Kuchta J.M., States S.J., McGlauchlin J.E., Overmeyer J.H., Wadowski R.M., McNamara A.M., Wolford R.S. and Yee R.B. (1985). Enhanced chlorine resistance of tap water-adapted *Legionella pneumophila* as compared with agar medium-passaged strains. *Appl. Env. Microbiol.* 50 (1): 21-26.

Kusnetsov J.M., Martikainen P.J., Jousimies-Somer H.R., Väisänen M.L., Tulkki A.I., Ahonen H.E. and Nevalainen A.I. (1993). Physical, chemical and microbiological water characteristics associated with the occurrence of Legionella in cooling tower systems. *Wat. Res.* 27 (1): 85-90.

Kusnetsov J.M., Jousimies-Somer H.R., Nevalainen A.I. and Martikainen P.J. (1994). Isolation of Legionella from water samples using various culture methods. *J. Appl. Bacteriol.* 76: 155-162.

Kusnetsov J.M., Ottoila E. and Martikainen P.J. (1996). Growth, respiration and survival of Legionella pneumophila at high temperatures. *J. Appl. Bacteriol.* 81: 341-347.

Lambert M.A. and Moss W. (1989). Cellular fatty acid compositions and isoprenoid quinone contents of 23 Legionella species. *J. Clin. Microbiol.* 27 (3): 465-473.

La Scola B., Mezi L., Weiller P.J. and Raoult D. (2001). Isolation of *Legionella anisa* using an amoebic coculture procedure. *J. Clin. Microbiol.* 39 (1): 365-366.

LeChevalier M.W., Cawthon C.D. and Lee R.G. (1988). Inactivation of biofilm bacteria. *Appl. Env. Microbiol.* 54 (10): 2492-2499.

Leoni E., Legnani P.P. (2001). Comparison of selective procedures for isolation and enumeration of *Legionella* species from hot water systems. *J. Appl. Microbiol.* 90 (1): 27-33.

Lever M.S. (1993). Production and characterisation of a *Legionella pneumophila* specific monoclonal antibody. *FEMS Microbiol. Lett.* 107 (1): 5-9.

Lin Y.E., Vidic R.D., Stout J.E. and Yu V.L. (1998). *Legionella* in water distribution systems. *J. Am. Wat. Works. Assoc.* 90 (9): 112-121.

Ludwig B., Schmid A., Marre R. and Hacker J. (1991). Cloning, genetic analysis, and nucleotide sequence of a determinant coding for a 19-kDa peptidoglycan-associated protein (Ppl) of *Legionella pneumophila*. *Infect. Immun.* 59 (8): 2515-2521.

Lo Presti F., Riffard S., Meugnier H., Reyrolle M., Lasne Y., Grimont P.A., Grimont F., Vandenesch F., Etienne J., Fleurette J. and Freney J. (1999). *Legionella taurinensis* sp. nov., a new species antigenically similar to *Legionella spiritensis*. *Int. J. Syst. Bacteriol.* 49 Pt 2: 397-403.

Lück P.C. (2000). Standardisation of typing of *Legionella*. Presented at the 5th International Conference on *Legionella*, September 26-29, Neu-Ulm, Germany.

Lundholm M., Jacobson K., Zettersten U. and Bernander S. (2000). Variations in *Legionella pneumophila* non-1 strains in nosocomial outbreaks (1993-1999) and environment at a Swedish University hospital. PFGE and AFLP analysis. Presented at the 5th International Conference on *Legionella*, September 26-29, Neu-Ulm, Germany.

Maartens G., Lewis S.J., de Gouveia C., Bartie C. *et al.* (1994). Atypical bacteria are a common cause of community-acquired pneumonia in hospitalised patients. *South African Med. J.* 84 (10): 678-682.

Mahbubani M.H., Bej A.K., Miller R., Haff L., DiCesare J. and Atlas R.M. (1990). Detection of *Legionella* with polymerase chain reaction and gene probe methods. *Mol. Cell. Probes* 4: 175-187.

* Maher W.E., Plouffe J.F. and Para M.F. (1982). Plasmid profiles of clinical and environmental isolates of *Legionella pneumophila* SG1. *J. Clin. Microbiol.* 18 (6): 1422-1423.

Maiwald M., Schill M., Stockinger C., Helbig J.H., Luck P.C., Witzleb W. and Sonntag H.G. (1995). Detection of *Legionella* DNA in human and guinea pig urine samples by the polymerase chain reaction. *Eur. J. Clin. Microbiol. Infect. Dis.* 14: 25-33.

Makin T. and Hart C.A. (1989). Detection of *Legionella pneumophila* in environmental water samples using a fluorescein conjugated monoclonal antibody. *Epidem. Inf.* 103: 105-112.

Marrie T.J., Haldane D., Bezanson G. and Peppard R. (1992). Each water outlet is a unique ecological niche for *Legionella pneumophila*. *Epid. Infect.* 108: 264-270.

* Martin W.T., Fields B.S. and Hutwagner L.C. (1993). Comparison of culture and polymerase chain reaction to detect legionellae in environmental samples. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp.175.

Martinelli F., Caruso A., Moschini L., Turano A., Scarcella C. and Speziani F. (2000). A comparison of *Legionella pneumophila* occurrence in hot water tanks and instantaneous devices in domestic, nosocomial and community environments. *Curr. Microbiol.* 41 (5): 374-376.

Mauff A.C. and Koornhof H.J. (1984). Legionellosis in South Africa. In: Thornsberry C., Balows A., Feeley J.C., Jakubowski W. (eds). *Legionella*, Proceedings of the 2nd International Symposium. American Society of Microbiology, Washington, DC.

* McClain M.S., Hurley M.C., Brieland J.K. and Engleberg N.C. (1996). The *Legionella pneumophila* *hel* locus encodes intracellularly induced homologs of heavy-metal ion transporters of *Alcaligenes* species. *Infect. Immun.* 64 (5): 1532-1540.

McDade J.E., Sheppard C.C., Fraser D.W., Tsai T.S., Redus M.A. and Dowdle W.R. (1977). Legionnaires' disease. Isolation of the bacterium and demonstration of its role in other respiratory disease. *N. Engl. J. Med.* 297: 1197-1203.

McNealy T., Newsome A., Johnson R. and Berk S. (2000). Impact of amoebae, bacteria and tetrahymenae on *Legionella pneumophila* multiplication and distribution in an aquatic environment. Presented at the 5th International Conference on Legionella, September 26-29, Neu-Ulm, Germany.

Moss C.W., Weaver R.E. and Dees S.B. (1977). Cellular fatty acid composition of isolates from Legionnaires' disease. *J. Clin. Microbiol.* 6: 140-143.

Muder R.R., Yu V.L. and Woo A.H. (1986). Mode of transmission of *Legionella pneumophila*. A critical review. *Arch. Intern. Med.* 146: 1607-1612.

Muraca P.W., Stout J.E., Yu V.L. and Yee Y.C. (1988). Legionnaires' disease in the work environment: implications for environmental health. *Am. J. Hyg. Assoc.* 49 (11): 584-590.

Neblett T.R., Riddle J.M. and Dumoffi M. (1979). Surface topography and fine structure of the Legionnaires' disease bacterium: a study of six isolates from hospitalised patients. *Ann. Intern. Med.* 90 (4): 648-651.

Ng D.L.K., Koh B.B., Tay L. and Heng B.H. (1997). Comparison of the polymerase chain reaction and conventional culture for the detection of legionellae in cooling tower waters in Singapore. *Lett. Appl. Microbiol.* 24: 214-216.

Nielsen K., Bangsberg J.M. and Høiby N. (2000). Susceptibility of *Legionella* species to five antibiotics and development of resistance by exposure to erythromycin, ciprofloxacin and rifampin. *Diagn. Microbiol. Infect. Dis.* (36 (1): 43-48.

Nolte F.S., Conlin C.A., Soisin A.J.M. and Redmond S.R. (1984). Plasmids as epidemiological markers in nosocomial Legionnaires' disease. *J. Infect. Dis.* 149 (2): 251-256.

OSHA Technical Manual. Section III: Chapter 7.

Palmer C.J., Tsai Y.L., Paszko-Kolva C., Mayer C. and Sangermano L.R. (1993). Detection of Legionella species in sewage and ocean water by the polymerase chain reaction, direct fluorescent-antibody and plate culture methods. *Appl. Environ. Microbiol.* 59: 3618-3624.

Osterholm M.T., Chin T.D., Osborne D.O., Dull H.B., Dean A.G., Fraser D.W., Hayes P.S. and Hall W.N. (1983). A 1957 outbreak of Legionnaires' disease associated with a meat packing plant. *Am. J. Epidemiol.* 117 (1): 60-66.

Ott M., Bender L., Chirinos E., Ehret W. and Hacker J. (1991). Phenotype versus genotype of the 19kDa peptidoglycan associated protein of Legionella (PplA), among Legionellae and other gram-negative bacteria. *Microb. Pathog.* 11 (5): 357-365.

Ott M., Bender L., Chirinos E., Ehret W. and Hacker J. (1991). Phenotype versus genotype of the 19kDa peptidoglycan associated protein of Legionella (PplA), among Legionellae and other gram-negative bacteria. *Microb. Pathog.* 11 (5): 357-365.

Otten S., Iyer S., Johnson W. and Montgomery R. (1986). Serospecific antigens of *Legionella pneumophila*. *J. Bacteriol.* 167 (3): 893-904.

Palmer C.J., Tsai Y.L., Paszko-Kolva C., Mayer C. and Sangermano L.R. (1993). Detection of Legionella species in sewage and ocean water by the polymerase chain reaction, direct fluorescent-antibody and plate culture methods. *Appl. Environ. Microbiol.* 59: 3618-3624.

Palmer C.J., Tsai Y.L., Paszko-Kolva C., Mayer C. and Sangermano L.R. (1993). Detection of Legionella species in sewage and ocean water by the polymerase chain reaction, direct fluorescent-antibody and plate culture methods. *Appl. Environ. Microbiol.* 59: 3618-3624.

Paszko-Kolva C., Hacker P.A., Stewart M.H. and Wolfe R.L. (1993). Inhibitory effect of heterotrophic bacteria on cultivation of *Legionella dumoffii*. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp. 203-205.

Paszko-Kolva C., Hacker P.A., Stewart M.H. and Wolfe R.L. (1993). Inhibitory effect of heterotrophic bacteria on cultivation of *Legionella dumoffii*. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp. 203-205.

Paszko-Kolva C., Yamamoto H., Shahamat M., Sawyer T.K., Morris G. and Colwell R.R. (1991). Isolation of amoebae and Pseudomonas and Legionella from eyewash stations. *Appl. Environ. Microbiol.* 57 (1): 163-167.

Paszko-Kolva C., Yamamoto H., Shahamat M., Sawyer T.K., Morris G. and Colwell R.R. (1991). Isolation of amoebae and Pseudomonas and Legionella from eyewash stations. *Appl. Environ. Microbiol.* 57 (1): 163-167.

Paszko-Kolva C., Yamamoto H., Shahamat M. and Colwell R.R. (1993). Polymerase chain reaction, gene probe and direct fluorescent antibody staining of *Legionella pneumophila* SG1 in drinking water and environmental samples. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC. Pp. 181-183.

Pearlman E., Engleberg N.C. and Eisenstein B.I. (1985). Identification of protein antigens of *Legionella pneumophila* serogroup 1. *Inf. Immun.* 47 (1): 74-79.

Pearlman E., Engleberg N.C. and Eisenstein B.I. (1985). Identification of protein antigens of *Legionella pneumophila* serogroup 1. *Inf. Immun.* 47 (1): 74-79.

Petersen L.R. and Catchpole M. (2001). Surveillance for infectious diseases in the European Union. A small European centre may have an important coordinating role (editorial). *British Med. J.* 323: 818-819.

Petitjean F., Dournon E., Strosberg A.D. and Hoebeke J. (1990). Isolation, purification and partial analysis of the lipopolysaccharide antigenic determinant recognised by a monoclonal antibody to *Legionella pneumophila* SG1. *Res. Microbiol.* 141 (9): 1077-1094.

Pine L., Franzus M.J. and Malcolm G.B. (1986). Guanine is a growth factor for *Legionella* species. *J. Clin. Microbiol.* 23 (1): 163-169.

Pine L., Hoffman P.S., Malcolm G.B., Benson R.F. and Franzus M.J. (1986). Role of keto-acids and reduced-oxygen-scavenging enzymes in the growth of *Legionella* species. *J. Clin. Microbiol.* 23 (1): 33-42.

Pine L., Hoffman P.S., Malcolm G.B., Benson R.F. and Keen M.G. (1984). Determination of catalase, peroxidase, and superoxide dismutase within the genus *Legionella*. *J. Clin. Microbiol.* 20 (3): 421-429.

Randall T.W., Naidoo P., Newton K.A., Botha P.W.A., Koornhof H.J. and Dubery B. (1980). Legionnaires' disease in Port Elizabeth. *South African Med. J.* 58 (17):17-23.

Reeves M.W., Pine L., Hutner S.H., George J.R. and Harrell W.K. (1981). Metal requirements of *Legionella pneumophila*. *J. Clin. Microbiol.* 13 (4): 688-695.

Reinthal F.F., Sattler J., Schaffler-Dulling K., Weinmayr B. and Marth E. (1993). Comparative study of procedures for isolation and cultivation of *Legionella pneumophila* from tap water in hospitals. *J. Clin. Microbiol.* 31 (5): 1213-1216.

Rietschel E.T., Brade H., Brade L., Kaca W., Kawahara K., Lindner B., Luderitz T., Tomita T., Schade U. and Seydel U. (1985). Newer aspects of the chemical structure and biological activity of bacterial endotoxins. *Prog. Clin. Biol. Res.* 189: 31-51.

- Riffard S., Douglas S., Brooks T., Springthorpe S., Fillion L.G. and Sattar S.A. (2001). Occurrence of Legionella in groundwater: an ecological study. *Water Sci. Tech.* 43 (12): 99-102.
- Ristroph J.D., Hedlund K.W. and Gowda S. (1981). Chemically defined medium for *Legionella pneumophila* growth. *J. Clin. Microbiol.* 13 (1): 115-119.
- Roberts K.P., August C.M. and Nelson J.D. (1987). Relative sensitivities of environmental legionellae to selective isolation procedures. *Appl. Environ. Microbiol.* 53 (12): 2704-2707.
- Robinson P.N., Heidrich B., Tiecke F., Fehrenbach F.J. and Rolfs A. (1996). Species-specific detection of Legionella using polymerase chain reaction and reverse dot-blotting. *FEMS Microbiol. Lett.* 140: 111-119.
- Roig J., Aguilar X., Ruiz J., Domingo C., Mesalles E., Manterola J. and Morera J. (1991). Comparative study of *Legionella pneumophila* and other nosocomially acquired pneumonias. *Chest* 99: 344-350.
- Rowbotham T.J. (1980). Preliminary report on the pathogenicity of *Legionella pneumophila* for freshwater and soil amoebae. *J. Clin. Pathol.* 33: 1179-1183.
- Rowbotham T.J. (1983). Isolation of *Legionella pneumophila* from clinical specimens via amoebae, and the interaction of those and other isolates with amoebae. *J. Clin. Pathol.* 36: 978-986.
- Ruef C. (1998). Nosocomial Legionnaires' disease – strategies for prevention. *J. Microbiol. Methods* 33: 81-91.
- Sadoski A.B., Wiater L.A. and Shuman H.A. (1993). Identification of *Legionella pneumophila* genes required for growth within and killing of human macrophages. *Infect. Immun.* 61 (12): 5361-5373.
- Saint C.P. and Ho L. (1999). A PCR test for the identification and discrimination of *Legionella longbeachae* serogroups 1 and 2. *J. Microbiol. Methods* 37 (3): 245-253.

Sampson J.S., Plikaytis B.B. and Wilkinson H.W. (1986). Immunologic response of patients with legionellosis against major protein-containing antigens of *Legionella pneumophila* SG1 as shown by immunoblot analysis. *J. Clin. Microbiol.* 23 (1): 92-99.

Selander R.K., McKinney R.M., Whittam T.S., Bibb W.F., Brenner D.J., Nolte F.S. and Pattison P.E. (1985). Genetic structure of populations of *Legionella pneumophila*. *J. Bacteriol.* 163: 1021-1037.

Segal G. and Shuman H.A. (1998). Intracellular multiplication and human macrophage killing by *Legionella pneumophila* are inhibited by conjugal components of IncQ plasmid RSF1010. *Mol. Microbiol.* 30 (1): 197-208.

Shelton B.G., Kerbel W., Witherell L. and Millar J.D. (2000). Review of Legionnaires' disease. *Am. Ind. Hyg. Assoc. J.* (61): 738-742.

Smith L., Carroll K. and Mottice S. (1993). Comparison of membrane filters for recovery of legionellae from water samples. *Appl. Env. Microbiol.* 59 (1): 344-346.

Sonesson A., Jantzen E., Bryn K., Larsson L. and Eng J. (1989). Chemical composition of a lipopolysaccharide from *Legionella pneumophila*. *Arch. Microbiol.* 153 (1): 72-78.

Sonesson A., Jantzen E., Bryn K., Tangen T., Eng J. and Zähringer U. (1994). Composition of 2,3-dihydroxy fatty acid-containing lipopolysaccharides from *Legionella israelensis*, *Legionella maceachernii* and *Legionella micdadei*. *Microbiology* 140 (6): 1261-1271.

Sonesson A., Jantzen E., Tangen T. and Zähringer U. (1994). Lipopolysaccharides of *Legionella erythra* and *Legionella oakridgensis*. *Can. J. Microbiol.* 40 (8): 666-671.

Sonesson A., Jantzen E., Tangen T. and Zähringer U. (1994). Chemical characterisation of lipopolysaccharides from *Legionella feeleeii*, *Legionella hackeliae* and *Legionella jordanis*. *Microbiology* 140 (10): 2663-2671.

Sonesson A., Jantzen E., Tangen T. and Zähringer U. (1994). Chemical composition of lipopolysaccharides from *Legionella bozemanii* and *Legionella longbeacheae*. *Arch. Microbiol.* 162 (4): 215-221.

Soni R., Seale J.P. and Young I.H. (1999). Fulminant psittacosis requiring mechanical ventilation and demonstrating serological cross-reactivity between *Legionella longbeacheae* and *Chlamydia psittaci*. *Respirology* 4 (2): 203-205 (abstract).

* Sopena N., Sabria M., Pedro-Botet M.L., Manterola J.M., Dominguez J., Modol J.M., Tudela P., Ausina V. and Foz M. (1999). Prospective study of community-acquired pneumonia of bacterial etiology in adults. *Eur. J. Clin. Microbiol. Infect. Dis.* 18 (12): 852-858.

Springston J. (2000). *Legionella lives*. *Engineering Systems*.
http://www.esmagazine.com/es/cda/articleinformation/features/bnp_fetures_item/0,2503,3615,00.html

Standards Australia AS 3896 (1991). Australian Standard: Waters – examination for legionella.

* States S.J., Conly L.F., Kuchta J.M., Oleck B.M., Lipovich M.J., Wolford R.S., Wadowski R.M., McNamara A.M., Sykora J.L. and Keleti G. (1987). Survival and multiplication of *Legionella pneumophila* in municipal drinking water systems. *Appl. Environ. Microbiol.* 53 (5): 979-986.

Steele T.W., Lanser J. and Sangster N. (1990). Isolation of *Legionella longbeacheae* serogroup 1 from potting mixes. *Appl. Environ. Microbiol.* 56 (1): 49-53.

Steinert M., Engelhart H., Flügel M., Wintermeyer E. and Hacker J. (1995). The Lly protein protects *Legionella pneumophila* from light but does not directly influence its intracellular survival in *Hartmanella vermiformis*. *Appl. Environ. Microbiol.* 61 (6): 2428-2430.

Susa M., Hacker J. and Mayer R. (1996). De novo synthesis of *Legionella pneumophila* antigens during intracellular growth in phagocytic cells. *Infect. Immun.* 64 (5): 1679-1684.

* Steinert M., Emödy L., Amann R. and Hacker J. (1997). Resuscitation of viable but non-culturable *Legionella pneumophila* Philadelphia JR32 by *Acanthamoeba castellanii*. *Appl. Environ. Microbiol.* 63 (5): 2047-2053.

Steinmetz I., Rheinheimer C. and Bitter-Suermann (1992). Rapid identification of legionellae by a colony blot assay based on a genus-specific monoclonal antibody. *J. Clin. Microbiol.* 30 (4): 1016-1018.

Ta A.C., Shakh-JS, Y. Y.L. and Rapoport M.M. (1990). Comparison of culture methods

St John G. and Steinman H.M. (1996). Periplasmic copper-zinc superoxide dismutase of *Legionella pneumophila*: role in stationary-phase survival. *J. Bacteriol.* 178 (6): 1578-1584.

Stone B.J. and Abu Kwaik Y. (1998). Expression of multiple pili by *Legionella pneumophila*: identification and characterisation of a type IV pilin gene and its role in adherence to mammalian and protozoan cells. *Infect. Immun.* 66 (4): 1768-1775.

* Technical Memoranda: Minimising the risk of Legionnaires' disease. (TM13:1987). The

Stout J.E. (1998). Culture methodology for *Legionella* species. HC Special Report 308. HC Information Resources, Fallbrook. CA.

Stout J.E., Lin Y.S., Goetz A.M. and Muder R.R. (1998). Controlling *Legionella* in hospital water systems: experience with the superheat-and-flush method and copper-silver ionisation. *Infect. Control Hosp. Epidemiol.* 19 (12): 911-914.

Stout M.J., Morse S.A. and Miller R.D. (1980). Intermediate metabolism in *Legionella*

Strebel P.M., Ramos J.M., Eidelman I.J., Tobiansky L., Koornhof H.J. and Küstner H.G.V. (1988). Legionnaires' disease in a Johannesburg teaching hospital. Investigation and control of an outbreak. *South African. Med. J.* 73: 329-333.

Tracter W.L., Benson R.P., Hawes L., Gidding R., Dwyer B., Molyneux W.R. and Grimont

Struelens M.J., Maes N., Deplano A., Bornstein N. and Grimont F. (1993). Comparison of methods for subtyping *Legionella pneumophila*. In: Barbaree J.M., Breiman R.F. and

* Dufour A.P. (eds). *Legionella. Current status and emerging perspectives*. American Society for Microbiology, Washington, DC.

cellular enzymes of *Legionella pneumophila*. *Infect. Immun.* 33 (2): 632-635.

Susa M., Hacker J. and Marre R. (1996). De novo synthesis of *Legionella pneumophila* antigens during intracellular growth in phagocytic cells. *Infect. Immun.* 64 (5): 1679-1684.

Isabel Roberts, in: Carlhorst J.M., Probst R.R. and Dubour A.P. (eds). *Legionella*

Szewzyk R., Allestam G. and Stenström T. (1991). Improved recovery of *Legionella* from water samples by use of black membrane filters. *Zbl. Hyg.* 192: 258-263,

Szewzyk U., Szewzyk R., Manz W. and Schleifer K. (2000). Microbiological safety of drinking water. *Ann. Rev. Microbiol.* 54: 81-127.

Ta A.C., Stout J.E., Yu V.L. and Wagener M.M. (1995). Comparison of culture methods for monitoring *Legionella* species in hospital potable water systems and recommendations for standardisation of such methods. *J. Clin. Microbiol.* 33 (8): 2118-2123.

Tarp B., Jensen J.S., Ostergaard L. and Andersen P.L. (1999). Search for agents causing atypical pneumonia in HIV-positive patients by inhibitor-controlled PCR assays.

Technical Memoranda: Minimising the risk of Legionnaires' disease. (TM13:1991). The Chartered Institution of Building Services Engineers, Delta House, 222 Balham High Road, London SW12 9BS.

Terranova W., Cohen M.L. and Fraser D.W. (1978). 1974 Outbreak of Legionnaires' disease diagnosed in 1977. *Lancet* Vol II, July 15: 122-124.

Tesh M.J., Morse S.A. and Miller R.D. (1983). Intermediary metabolism in *Legionella pneumophila*: utilisation of amino acids and other compounds as energy sources. *J. Bacteriol.* 154 (3): 1104-1109.

Thacker W.L., Benson R.F., Hawes L., Gidding H., Dwyer B., Mayberry W.R. and Brenner D.J. (1991). *Legionella fairfieldensis* sp. nov. isolated from cooling tower waters in Australia. *J. Clin. Microbiol.* 29 (3): 475-478.

Thorpe T.C. and Miller R.D. (1981). Extracellular enzymes of *Legionella pneumophila*. *Inf. Imm.* 33 (2): 632-635.

Tiefenbrunner F., Arnold A., Taraboi E., Cernek U. and Emde K. (1993). Comparison of different detection methods for isolation of *L. pneumophila* from water supplied of Alpine

hotel resorts. In: Barbaree J.M., Breiman R.F. and Dufour A.P. (eds). *Legionella. Current status and emerging perspectives*. American Society for Microbiology, Washington, DC.

Tobiansky L., Drath A., Dubery B. and Koornhof H.J. (1986). Seasonality of *Legionella* isolated from environmental sources. *Isr. J. Med. Sci.* 22 (10): 640-643.

Tobin J.O., Watkins I.D., Woodhead S. and Mitchell R.G. (1986). Epidemiological studies using monoclonal antibodies to *Legionella pneumophila* SG 1. *Isr. J. Med. Sci.* 22: 711-714.

Tompkins L.S. and Loutit J.S. (1993). Detection of *Legionella* by molecular methods. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella: Current Status and Emerging Perspectives*. American Society for Microbiology, Washington, DC. Pp 163-168.

Tyndall R.L. and Domingue E.L. (1982). Cocultivation of *Legionella pneumophila* and free-living amoebae. *Appl. Env. Microbiol.* 44 (4): 954-959.

Vandenesch F., Surgot M., Bornstein N., Paucod J.C., Marmet D., Isoard P. and Fleurette J. (1990). Relationship between free amoeba and *Legionella*: studies in vitro and in vivo. *Zbl. Bakt.* 272: 265-275.

Vesey G., Dennis P.J., Lee J.V. and West A.A. (1988). Further development of simple tests to differentiate the legionellas. *J. Appl. Bacteriol.* 65: 339-345.

Viswanathan V.K., Edelstein P.H., Pope C.D. and Cianciotto N.P. (2000). The *Legionella pneumophila* iraAB locus is required for iron assimilation, intracellular infection and virulence. *Infect. Immun.* 68 (3): 1069-1079.

Wadowski R.M. and Yee R.M. (1985). Effect of non-Legionellaceae bacteria on the multiplication of *Legionella pneumophila* in potable water. *Appl. Env. Microbiol.* 49 (5): 1206-1210.

Wiater L.A., Dunn K., Maxfield F.R. and Shuman H.A. (1998). Early events in phagosome establishment are required for intracellular survival of *Legionella pneumophila*. *Infect. Immun.* 66 (9): 4450-4460.

* Wiese A., Brandenburg K., Ulmer A.J., Seydel U. and Muller-Loennies S. (1999). The dual role of lipopolysaccharide as effector and target molecule. *Biol. Chem.* 380 (7-8): 767-784.

Wilkinson H.W., Drasar V., Thacker W.L., Benson R.F., Schindler J., Potuznikova B., Mayberry W.R. and Brenner D.J. (1988). *Legionella moravica* sp. nov. and *Legionella brunensis* sp. nov. isolated from cooling tower water. *Ann. Inst. Pasteur Microbiol.* 139 (4): 393-402.

Williams H.N., Paszko-Kolva C., Shahamat M., Palmer C., Pettis C. and Kelley J. (1996). Molecular techniques reveal high prevalence of *Legionella* in dental units. *J. Am. Dent. Assoc.* 127 (8): 1188-1193.

Winn W.C. (1988). Legionnaires' disease: historical perspective. *Clin. Microbiol. Rev.* 1 (1): 60-81.

* Wintermeyer E., Flügel M., Ott M., Steinert M., Rdest U., Mann K.H. and Hacker J. (1994). Sequence determination and nutational analysis of the lly locus of *Legionella pneumophila*. *Infect. Immun.* 62 (3): 1109-1117.

Wireman J.W., Schmidt A., Scavo C.R. and Hutchins D.T. (1993). Biofilm formation by *Legionella pneumophila* in a model domestic hot water system. In: Barbaree J.M., Breiman R.F. and DuFour A.P. (eds). *Legionella, Current Status and Emerging Perspectives*. American Society for Microbiology. Washington, DC.

Worksafe Australia (1989). Legionnaires' disease and related illnesses. National Occupational Health and Safety Commission. Australian Government Publishing Service, Canberra.

Wright J.B., Ruseska I., Athar M.A., Corbett S. and Costerton J.W. (1989). *Legionella pneumophila* grows adherent to surfaces in vitro and in situ. *Infect. Control Hosp. Epidemiol.* 10 (9): 408-415.

Yamamoto H., Klein T.W., Shinomiya H., Nakano M. and Friedman H. (1992). Infection of macrophages with *Legionella pneumophila* induces phosphorylation of a 76-kDa protein. *Infect. Immun.* 60 (8): 3452-3455.

Yamamoto H., Sugiura M., Kusunoki S., Ezaki T., Ikedo M. and Yabuuchi E. (1992). Factors stimulating propagation of legionellae in cooling tower water. *Appl. Environ. Microbiol.* 58 (4): 1394-1397.

Yu V.L. (1990). *Legionella pneumophila* (Legionnaires' disease). In: Mandell, Douglas and Bennet (eds). Principles and Practice of Infectious Diseases. Third Edition. Churchill Livingstone Inc.

Yu V.L. (1998). Resolving the controversy on environmental cultures for *Legionella*: a modest proposal. *Inf. Control Hosp. Epid.* 19 (12): 983-897.

Zähringer U., Knirel Y.A., Lindner B., Helbig J.H., Sonesson A., Marre R. and Rietschel E.T. (1995). The lipopolysaccharide of *Legionella pneumophila* serogroup 1 (strain Philadelphia 1): chemical structure and biological significance. *Prog. Clin. Biol. Res.* 392: 113-139.