

Supplementary Table 1: Site of collection of most frequently identified genotypes (N=202)

Genotypes	Number of strains detected per genotype (%)				
	Site 1	Site 2	Site 5	Site 6	Site 7
	n=68 (%)	n=22 (%)	n=11 (%)	n=61 (%)	n=40 (%)
HAstV-1	11 (16.2)	-	2 (18.2)	12 (19.7)	6 (15.0)
HAstV-2	2 (2.9)	2 (9.1)	-	-	1 (2.5)
HAstV-3	-	-	-	3 (4.9)	-
HAstV-4	-	2 (9.1)	-	-	-
HAstV-5	3 (4.4)	1 (4.5)	3 (27.3)	5 (8.2)	6 (15.0)
HAstV-6	-	1 (4.5)	-	1 (1.6)	-
HAstV-8	1 (1.5)	-	-	4 (6.6)	2 (5.0)
T8/T1	11 (16.2)	8 (36.4)	3 (27.3)	10 (16.4)	9 (22.5)
T8/T4	6 (8.8)	1 (4.5)	1 (9.1)	4 (6.6)	-

KEY:

Site 1: Chris Hani Baragwanath Academic Hospital, Gauteng

Site 2: Mapulaneng Hospital and Matikwane Hospital, Mpumalanga

Site 5: Edendale Hospital, Kwa-Zulu Natal

Site 6: Red-Cross Hospital, Western Cape

Site 7: Ngwelezane Hospital, Kwa-Zulu Natal