

The physical demands of wheelchair tennis match play: a systematic review with meta-analysis

Running heading: Physical demands of wheelchair tennis

Corresponding Author:

Babette M. Pluim

b.pluim@knltb.nl

Samuel Williamson, ORCID iD 0000-0002-3533-0603

International Tennis Federation, London, UK

UK Sports Institute, United Kingdom

Clare L. Ardern, ORCID iD 0000-0001-8102-3631

Department of Physical Therapy, University of British Columbia, Vancouver, Canada

Sport and Exercise Medicine Research Centre, La Trobe University, Melbourne, Australia

Cain Berry

International Tennis Federation, London, United Kingdom

Neil Heron, ORCID iD 0000-0002-4123-9806

Centre for Public Health, Queen's University Belfast, Northern Ireland

School of Medicine, Keele University, Staffordshire, England.

Dina C. Janse van Rensburg, ORCID iD 0000-0003-1058-6992

Section Sports Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

Medical Committee, World Netball, Manchester, United Kingdom

Marleen G.T. Jansen, ORCID iD 0000-0002-8441-7748

Toptennis Department, Royal Netherlands Lawn Tennis Association (KNLTB), Amstelveen, the Netherlands.

Centre for Human Movement Sciences, University Medical Centre Groningen, University of Groningen, Groningen, the Netherlands

Sam McCormick, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Northern Ireland

Machar Reid, ORCID iD 0000-0002-7196-3731

Tennis Australia, Melbourne, Australia

Alejandro Sánchez Pay, ORCID iD 0000-0003-1600-4172

Faculty of Sport Science, University of Murcia, Spain

Tobias Saueressig, ORCID iD 0000-0002-3332-3856

Physio Meets Science GmbH, Leimen, Germany

Linda J. Schoonmade, ORCID iD 0000-0002-2407-5977
University Library, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

Robert B. Shaw
Spinal Cord Injury British Columbia, Vancouver, Canada

Rienk van der Slikke, ORCID iD 0000-0002-3630-3519
Human Kinetic Technology, The Hague University of Applied Sciences, the Hague, Netherlands

Nick Webborn, ORCID iD 0000-0003-3636-5557
IPC Medical Committee, Bonn, Germany
Centre for Sport and Exercise Science and Medicine (SESAME), School of Sport and Service
Management, University of Brighton, Brighton, United Kingdom
School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, United
Kingdom

Babette M. Pluim, ORCID iD 0000-0003-0655-8980
1. Section Sports Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa
2. Amsterdam Collaboration on Health & Safety in Sports (ACHSS), IOC Research Center of
Excellence, Amsterdam, the Netherlands
3. Department of Orthopaedic Surgery and Sports Medicine, Amsterdam UMC location University of
Amsterdam, Amsterdam, the Netherlands
4. Academic Center for Evidence-based Sports medicine (ACES), Amsterdam, the Netherlands
5. Medical Department, Royal Netherlands Lawn Tennis Association (KNLTB), Amstelveen, the
Netherlands

ABSTRACT

Background

Wheelchair tennis, a globally popular sport, features a professional tour spanning 40 countries and over 160 tournaments. Despite its widespread appeal, information about the physical demands of wheelchair tennis is scattered across various studies, necessitating a comprehensive systematic review to synthesise available data.

Objective

To provide a detailed synthesis of the physical demands associated with wheelchair tennis, encompassing diverse factors such as court surfaces, performance levels, sport classes, and sexes.

Methods

We conducted comprehensive searches in the PubMed, Embase, CINAHL, and SPORTDiscus databases, covering articles from inception to March 1, 2023. Forward and backward citation tracking from the included articles was carried out using Scopus, and we established eligibility criteria following the PECOS framework. Our study focused on wheelchair tennis players participating at regional, national, or international levels, including both juniors and adults, and open and quad players. We analysed singles and doubles matches and considered sex (male, female), sport class (open, quad) and court surface type (hard, clay, grass) as key comparative points. The outcomes of interest encompassed play duration, on-court movement, stroke performance, and physiological match variables. The selected study designs included observational cross-sectional, longitudinal, and intervention studies (baseline data only). We calculated pooled means or mean differences with 95% confidence intervals and employed a random-effects meta-analysis with robust variance estimation. We assessed heterogeneity using Cochrane Q and 95% prediction intervals.

Results

Our literature search retrieved 643 records, with 24 articles meeting our inclusion criteria. Most available information focused on international male wheelchair tennis players in the open division, primarily competing in singles on hard courts. Key findings (mean [95%CI]) for these players on hard courts were: match duration 65.9 minutes [55.0, 78.8], set duration 35.0 minutes [28.2, 43.5], game duration 4.6 minutes [0.92, 23.3], rally duration 6.1 seconds [3.7, 10.2], effective playing time 19.8% [18.9, 20.7], and work-to-rest ratio 1:4.1 [1:3.7, 1:4.4]. Insufficient data were available to analyse play duration for female players. However, for available data on hard court matches, the average set duration was 34.8 minutes [32.5, 37.2]. International male players on hard court covered an average distance per match of 3859 metres [1917, 7768], with mean and peak average forward speeds of 1.06 m/s [0.85, 1.32] and 3.55 m/s [2.92, 4.31] respectively. These players executed an average of 365.9 [317.2, 422.1] strokes per match, 200.6 [134.7, 299.0] per set, 25.4 [16.7, 38.7] per game, and 3.4 [2.6, 4.6] per rally. Insufficient data were available for a meta-analysis of female players' on-court movement and stroke performance. The average and peak heart rates of international male players on hard court were 134.3 [124.2, 145.1] and 166.0 [132.7, 207.6] beats per minute, and the average match heart rate expressed as a percentage of peak heart rate was 74.7% [46.4, 100]. We found no studies concerning regional players or juniors, and only one study on doubles match play.

Conclusions

While we present a comprehensive overview of the physical demands of wheelchair tennis, our understanding predominantly centers around international male players competing on hard courts in the open division. To attain a more comprehensive insight into the sport's physical requirements,

future research should prioritise the inclusion of data on female and quad players, juniors, doubles, and matches played on clay and grass court surfaces. Such endeavours will facilitate the development of more tailored and effective training programs for wheelchair tennis players and coaches.

The protocol for this systematic review was registered a priori at the International Platform of Registered Systematic Review and Meta-analysis Protocols (Registration doi: 10.37766/inplasy2023.3.0060).

Key Points

The mean match duration for open class international male and female wheelchair tennis players in singles on hard court was just over an hour.

Mean game duration for male players was around 5 minutes, with a mean rally duration of 6 seconds.

Their effective playing time was 20% of the actual playing time, with a work to rest ratio of 1:4.

On average, open class international male players covered almost 4 km per match on hard court, with mean and peak average forward speeds of 3.8 and 12.8 km/hour.

The average and peak match heart rates of international male players on hard court were 134 and 166 beats per minute, respectively. Average match heart rate expressed as a percentage of peak heart rate was 75%.

These players hit an average of 365 strokes per match, 200 strokes per set, 25 per game, and 3 to 4 per rally.

1.0 Introduction

Wheelchair tennis, the Para sport version of tennis [1], caters to individuals with physical impairments and was first played in 1976 [2]. The sport involves playing tennis while seated in a wheelchair and follows the same rules as standing tennis, except that the ball can bounce twice. Matches typically follow a best-of-three-sets format, with tie-breaks used to determine set outcomes when necessary. Wheelchair tennis promotes inclusive participation and psychosocial wellbeing[3], accommodating players at both recreational and professional levels. In recreational play, there is even the flexibility to mix standing and seated players. The sport is played on various court surfaces, including hard courts, clay, and grass.

Wheelchair tennis has two sport classes: the Open division, for players with a permanent impairment in one or both of their lower extremities, and the Quad division, for players with additional permanent impairments in their upper extremities that limit their ability to handle the racket, play shots and manoeuvre the wheelchair. Athlete eligibility for participation and class determination is undertaken according to the ITF classification rules[4]. Current estimates suggest there are around 10,000 recreational wheelchair tennis players globally, with fewer than 1,000 participating in regular international competition[5]. The sport is played in over 100 countries, and the ITF wheelchair tennis tour features more than 160 tournaments across 40 countries worldwide[5].

To excel in wheelchair tennis, players must possess various tactical, technical, physical, and psychological skills[6-8]. As the sport evolves, physical attributes have become increasingly important for success at the elite level. The physical demands of wheelchair tennis are extensive, including strength[9], power[10], balance[11], coordination[12, 13], agility[12, 13], and aerobic endurance[14]. The court surface,[15, 16] player ranking[17], division[18], age[13] and sex[19, 20] can all influence the demands of match-play in wheelchair tennis.

A thorough understanding of the physical demands of wheelchair tennis is central to developing players effectively and monitoring the progression of this relatively new professional sport. While the physical demands of standing tennis have been documented in recent literature [21], there is no published review exploring the physical demands of wheelchair tennis.

Therefore, we aimed to summarise the physical demands of wheelchair tennis, focusing on singles and doubles in the open and quad divisions, across all age groups, sexes, performance levels, and court surfaces, by reviewing the available scientific literature on the subject.

2.0 Methods

The reporting of this systematic review was guided by the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) Statement [22]. We prospectively registered the review protocol at the International Platform of Registered Systematic Review and Meta-analysis Protocols to ensure transparency and adherence to a predetermined plan (Registration doi: 10.37766/inplasy2023.3.0060).

2.1 Eligibility Criteria

The PECOS (Population, Exposure, Comparison, Outcome, and Study design) framework was used to define the eligibility criteria.

The inclusion criteria were:

- i) Population: Male and female wheelchair tennis players of regional, national, or international playing level, including both juniors (≤ 18 years) and adults;
- ii) Exposure: Singles and doubles wheelchair tennis matches played on hard court, clay, or grass, following the rules set forth by the International Tennis Federation (ITF);
- iii) Comparison: Sex (male/female), court surface (hard, clay, grass), sport class (open/quad);
- iv) Outcome: Studies must report at least one parameter related to the duration of play (e.g., strokes, rallies, games, sets, and matches), on-court movement (e.g., accelerations, decelerations, changes of direction, distance covered, and speed), stroke performance (count, speed, and scoring), or physiological variables (e.g., heart rate, oxygen uptake, energy expenditure, and blood lactate);
- v) Study designs: descriptive cross-sectional studies, analytic observational prospective and retrospective cohort studies, as well as intervention studies (only baseline data included)

The exclusion criteria were:

- i) Editorials, notes, letters, case reports, and reviews;
- ii) Studies of wheelchair tennis with modified match rules (e.g., time-capped matches);
- iii) Studies of standing (able-bodied) tennis;
- iv) Studies reporting only on biomechanical variables.

2.2 Data Sources and Searches

A comprehensive search to identify relevant studies was performed in four bibliographic databases: PubMed, Embase.com, CINAHL (via Ebsco) and SPORTDiscus (via Ebsco), covering the period from inception to March 1, 2023. The search was developed with a medical information specialist (LS). It used both controlled search terms (MeSH in PubMed and Emtree in Embase, CINAHL Headings in CINAHL, and Thesaurus terms in SPORTDiscus) and free-text terms. The search strategy focused on the keywords 'wheelchair' and 'tennis' as the index terms or free-text words, along with their synonyms and closely related terms. No language or date restrictions were applied. Duplicate articles were removed by the medical information specialist (LS) using Endnote X20.5 (Clarivate), following the Amsterdam Efficient Deduplication (AED) method [23] and the Bramer method [24]. A backward citation search was also conducted for all included articles using Scopus. The comprehensive database search strategy can be found in **Online Resource 1**.

2.3 Selection Process

Two reviewers (SW and BMP) independently screened all titles and abstracts for eligibility using Rayyan [25]. The full-text articles were assessed as required. The inter-rater reliability was assessed using Cohen's Kappa for both the initial and the final screening. Any discrepancies in the assessment were resolved through discussion until an agreement was reached. If the reviewers could not reach a consensus, an independent reviewer (MGTJ) was available for the final decision. Reasons for exclusion were documented at each stage of the screening process.

2.4 Data extraction

Two reviewers (SW and BMP) independently extracted the following information regarding the characteristics of the included studies: name of the first author; year of publication; country in which the study was conducted; study design; population; sample size (number of participants and matches); age of participants; percentage of male participants; playing level; court surface; sport

class (Open or Quad); assessment tool(s) used; comparison; and a summary of the outcome parameters of each study. Data related to the outcome were extracted for analysis. When there was overlap between studies, we removed the duplicate data. Authors were contacted if data were missing or required clarification.

The playing level of the participants was determined based on the level of the tournaments the player participated in, such as regional, national, or international, or by their ranking. The sport class (Open or Quad) was determined by the classification status described in the article or the reported impairment. The court surfaces were classified as hard, clay, or grass, based on the playing surface of the matches. Differences were resolved through discussion, and a third reviewer (MGJJ) was available if consensus could not be reached.

Data related to outcomes in the following areas were extracted (Table 1):

- Time characteristics: matches, sets, games, points (rallies), and strokes
- On-court movement: accelerations, decelerations, turns, distance covered, average and peak wheelchair speed
- Physiological variables: heart rate, oxygen uptake, energy expenditure, blood lactate levels
- Stroke performance: speed, number, and scoring

Table 1. Outcome parameters for time characteristics, on-court movement, stroke performance, and physiological variables

Time characteristics	Stroke performance (speed)
Match/set/game duration (min)	First serve speed ($\text{km}\cdot\text{h}^{-1}$)
Rally/stroke duration (s), Rally pace ($\text{strokes}\cdot\text{s}^{-1}$)	Second serve speed ($\text{km}\cdot\text{h}^{-1}$)
Effective playing time (%)	Groundstroke speed ($\text{km}\cdot\text{h}^{-1}$)
Work-to-rest ratio	Forehand speed ($\text{km}\cdot\text{h}^{-1}$)
	Backhand speed ($\text{km}\cdot\text{h}^{-1}$)
On-court movement	
Accelerations/decelerations per minute/speed zone/match (#)	Stroke performance (number)
Turns per match/minute (#)	First serve percentage (%)
Distance covered per match/set/game/point/stroke (m)	Strokes per match/set/game/rally/second (#)
Distance covered per speed zone/minute (m)	Serves per match/set/game (#)
Average/peak movement speed ($\text{m}\cdot\text{s}^{-1}$)	First serves per match/set/game (#)
	Second serves per match/set/game (#)
Physiological match variables	Forehands per match/set/game (#)
Average/peak heart rate (min^{-1}), percentage peak heart rate (%)	Backhands per match/set/game (#)
Oxygen uptake, relative ($\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$)	
Oxygen uptake, absolute ($\text{L}\cdot\text{min}^{-1}$)	Scoring
Percentage peak oxygen uptake (%)	Points per match/set/game (#)
Energy expenditure, relative ($\text{kcal}/\text{kg}/\text{match}$)	Games per match/set (#), sets per match (#)
Energy expenditure, absolute (kcal/match)	Bounces per point (#)
Mean blood lactate ($\text{mmol}\cdot\text{L}^{-1}$)	Shots hit off one bounce (%)

2.5 Methodological Quality Assessment

Two reviewers (SW and BMP) used the Joanna Briggs Institute checklist for analytical cross-sectional studies to assess the methodological quality of the included studies [26]. The checklist comprised eight items that included questions on study inclusion criteria, participants, setting, exposure, condition,

confounding factors, validity and reliability of the measurement technique, and appropriate statistical analysis. Each question was rated as 'yes', 'no', or 'unclear'. Any differences in rating were resolved through discussion until a consensus was reached. The quality assessment outcomes were not used to determine study inclusion or for sub-group analysis based on methodological quality or risk of bias.

2.6 Data Synthesis

The analysis examined the outcomes across two broad categories: "male" and "female".

Quantitative statistical analysis was conducted by calculating pooled means or mean differences, along with 95% confidence intervals (CI) for movement variables reported by three or more studies. In cases where median values were provided, the quantile estimation method was employed to estimate the mean and standard deviations [27]. To account for the dependence of study means, a random-effects meta-analysis was performed using robust variance estimation [28, 29]. The inverse variance method was utilised to pool the studies [30]. For analysis, measures of means were log-transformed and subsequently back-transformed to avoid implausible (i.e., negative) estimates [31].

The standard error of the log-transformed mean was calculated using the formula $\sqrt{\frac{SD^2}{n * mean^2}}$ [32].

The Cochrane Q and resulting chi-square statistic, I^2 statistic, and 95% prediction intervals (PI) were employed to assess the heterogeneity. The 95% PI estimates where the actual effects are expected to lie for 95% of similar studies that may be conducted in the future. The PI estimate may be imprecise if the number of studies is limited [33]. Therefore, we only provided the PI if we had at least five studies available.

In cases where correlation values were absent, a default correlation value of $\rho = 0.8$ was adopted for all analyses. All calculations and graphical representations were conducted using the software R [34], along with the additional packages 'metafor' [35] and 'robumeta' [36].

Qualitative Synthesis

The descriptive characteristics of each study were summarised and presented in tables and text.

Subgroup analysis

The influence of the three main court surfaces used in tennis was considered: hard, grass, and clay.

Analysis of subgroups or subsets

We planned subgroup analysis with study variables in the following subgroups: male and female; international, national, and regional level players; open and quad division; juniors and adults.

Sensitivity Analysis

As correlation values were unknown, a sensitivity analysis with a range of different correlation parameters was performed ($\rho = 0, 0.2, 0.4, 0.6, 1.0$). Missing standard deviations were imputed as the median value of the included standard deviations in the corresponding analysis [36]. Sensitivity analyses were calculated without the studies with the imputed standard deviation.

3.0 Results

3.1 Search

The literature search resulted in 538 references: 85 in PubMed, 116 in Embase, 74 in CINAHL, and 263 in SPORTDiscus. After removing duplicates, 381 papers remained. A forward and backward citation search and handsearching generated an additional 262 references, resulting in a total of 643 records for screening. Based on titles and abstracts, 33 articles that seemed to meet the inclusion criteria were selected. After a full-text screening, 9 articles were excluded, leaving 24 articles for the review (Figure 1). Cohen's Kappa for the initial screening was 0.96 and for the final screening 1.0, indicating a very high level of agreement.




Figure 1. Flow chart of the screening process.

Table II. Characteristics of the included studies

1 st author, year, country	Design	Setting	Sample size (n)	Matches (n)	Age (mean (SD)) (y)	Sex (male: n (%))	Playing level	Sporting class	Playing surface	Assessment tool	Comparison	Outcomes measures
Bernardi, 2010, Italy [37]	Analytic cross-sectional	Simulated match play (two sets)	4	2	38.5 (10.3)	4 (100)	International	Open	Hard court	HR monitor, incremental arm cranking	Five para sports	Average and peak heart rate, % peak heart rate, average VO ₂ and %VO _{2peak}
Croft, 2010, UK [38]	Analytic cross-sectional	International tournaments in Florida (USA) & UK	6	6	23 (8.2)	4 (66.7)	International	Open	NR	HR monitor, incremental submaximal exercise test	Wheelchair basketball vs wheelchair tennis	Average and peak heart rate, % peak heart rate, average VO ₂ and % VO _{2peak}
Filipic, 2009a, Slovenia [39]	Analytic cross-sectional	Simulated match play (two sets)	15	22	39.1 (8.2)	15 (100)	International (5), all minimum 5y experience	Open	Hard court	Video, SAGIT/TENNIS tracking system	Winner vs losers	Match duration, points per match, rally duration, EPT
Filipic, 2009b, Slovenia [40]	Analytic cross-sectional	Simulated match play (two sets)	15	22	39.1 (8.2)	15 (100)	International (5/15), minimum 5y experience	Open	Hard court	Video, SAGIT/TENNIS tracking system	High vs low ranked, winner vs loser	Average and peak forward speed, distance per point/game/match
Gómez, 2021 [41]	Analytic cross-sectional	XXIII Spanish National Masters in Wheelchair Tennis	8	10	39.4 (11.3)	8 (100)	International	Open	Clay court	Video, IMU	High vs low-level, high vs high-level, low vs low-level	Distance per point/match, distance during active play, average forward speed
Mason, 2020, UK [18]	Analytic cross-sectional	NEC Wheelchair Tennis Masters (best of three)	17	22	29(9), 28 (6), 35 (6)	11 (64.7)	International	Open (7 male, 6 female), quad (4 male)	Hard court	Video, Dartfish TeamPro Data 6.0, IMU, LPS	Male vs female, Open vs Quad	Set duration, distance per set, average and peak forward speed, forward/rotational acceleration, turns per minute, strokes per rally, points per set

Ponzano, 2017, Italy [42]	Analytical cross-sectional	Simulated match play (best of three)	12	12	43,3 (13,3)	10 (83,3)	International (m), national (F)	Open	Hard court, clay	GPS, HR monitor	Hard vs clay court	Match duration, average and peak heart rate, average and peak forward speed, acceleration, deceleration, kcal per match
Roy, 2006, USA [14]	Analytic cross-sectional	Simulated match play (best of three)	6	6	40.2 (9.8)	6 (100)	International (4), national (2)	Open	Hard court	HR monitor, video	None	Average heart rate, % peak heart rate, average VO ₂ , % VO _{2peak} , kcal per match
Sánchez-Pay, 2013, Spain [43]	Analytic cross-sectional	Roland Garros, US Open (2010, 2011) (best of three)	54	27	NR	54 (100)	International	Open	Clay, hard court	Match statistics	Hard vs clay court	Match duration, 1 st serve percentage
Sánchez-Pay, 2015a, Spain [44]	Analytic cross-sectional	Spanish Wheelchair Tennis Master Cup (two sets)	8	16	NR	8 (100)	International	Open	Hard court	Video	With and without breakpoint opportunity	Match/rally duration, EPT, strokes/points per rally/game/set/match
Sánchez-Pay, 2015b, Spain [19]	Analytic cross-sectional	2014 AO, RG, and US Open (best of three)	96	84	NR	48 (50)	International	Open	Hard court, clay court	Match statistics	Hard vs clay court and wheelchair vs standing tennis	Set duration, points per game/set, 1 st serve percentage
Sánchez-Pay, 2015c, Spain [45]	Analytic cross-sectional	2012 Paralympic Games (best of three)	32	32	NR	0 (0)	International	Open	Hard court	Match statistics	Winners vs losers	1 st serve percentage
Sánchez-Pay, 2015d, Spain [46]*	Analytic cross-sectional	Simulated match play (best of three)	4	6	34.8 (11.6)	4 (100)	International	Open	Clay court	Video	None	Rally/set/match duration, WR ratio, points per set/match, strokes per rally/game/set/match, rally pace
Sánchez-Pay, 2016, Spain [47]*	Analytic cross-sectional	Simulated match play (best of three)	4	6	34.8 (11.6)	4 (100)	International	Open	Clay court	HR monitor, lactate analyser, video	Serve vs return situation	Rally/set/match duration, EPT, WR ratio, strokes per rally, average and peak heart

												rate, % peak heart rate, blood lactate
Sánchez-Pay, 2017a, Spain [48]	Analytic cross-sectional	Spanish Wheelchair Tennis Master Cup (best of three)	8	16	NR	8 (100)	International	Open	Hard court	Video	Set winner vs set loser; with vs without game point opportunity	Shots per match
Sánchez-Pay 2017b, Spain [20]	Analytic cross-sectional	International Wheelchair Tennis Tournament (best of three)	16	16	NR	8 (50)	International	Open	Hard court	Video	Male vs female; with vs without game point opportunity	Rally/set/match duration, EPT, WR ratio, strokes/points per set, strokes per rally, rally pace, bounces per point
Sánchez-Pay, 2017c, Spain [49]	Analytic cross-sectional	2012 Paralympic Games (best of three)	64	64	NR	64 (100)	International	Open	Hard court	Match statistics	Set winner vs set loser	1 st serve percentage
Sánchez-Pay, 2021a, Spain [50]	Analytic cross-sectional	2004, 2008, 2012, 2016 Paralympic Games (best of three)	NR	12	NR	12 (100)	International	Open	Hard court	Video, LINCE	Paralympic Games in 2004 vs 2008 vs 2012 vs 2016	Game/set/match duration, points per game/set/match, shots per game/set/ match
Sánchez-Pay, 2021b, Spain [15]	Analytic cross-sectional	2018 World Cup, Master Cup, Queens Club Championships (best of three)	10	13	31.9 (7.7)	10 (100)	International	Open	Hard court, clay, grass	Video	Hard vs clay vs grass court	Rally/game/set duration, points/shots per game/set, strokes per rally, 1 st serve percentage
Sánchez-Pay, 2023, Spain [51]	Analytic cross-sectional	National Master Cup (best of three)	8	16	38 (10)	8 (100)	International	Open	Clay court	LPS, IMU	Day 1 vs day 2 vs day 3 vs day 4	Distance per match
Sindall, 2013a, UK [52]	Analytic cross-sectional	2010 British Open (three sets)	15	17 (Open), 9 (Quad)	NR	11 (73,3)	International	Open, Quad	Hard court	GPS, data logging device	GPS vs Data logger	Distance per match, average and peak forward speed

Sindall 2013b, UK [17]	Analytic cross- sectional	2009 British Open (two sets)	14	14	32.0 (15.4)	14 (100)	International	Open	Hard court	Data logging device, HR monitor	High vs low- ranked, winner vs loser	Set/match duration, distance per set/match, average and peak forward speed, average and peak heart rate, % peak heart rate
Sindall, 2015, UK [53]	Analytic cross- sectional	Three ITF tournaments (three sets)	31	31	30 (12)	23 (74,2)	International	Open	Hard court	Data logger device, video	High vs low- ranked, male vs female, singles vs doubles	Time spent in speed zones
Van der Slikke, 2020, Netherland s [54]	Analytic cross- sectional	2016 NWT Championships 2017 ABN AMRO Open (best of three)	15	15	NR	15 (100)	International	Open	Hard court	Inertial sensors (incl. gyroscope)	Wheelchair basketball vs wheelchair tennis	Average and peak forward speed, peak rotational speed, forward/rotational acceleration

AO Australian Open, EPT Effective Playing Time, F Female, GPS Global Positioning System, HR Heart rate, % Percentage, IMU Inertial Measurement Unit, ITF International Tennis Federation, LPS Local Positioning System; M Male, NR Not reported, NWT National Wheelchair Tennis, RG Roland Garros, US United States, VO₂ Oxygen Consumption, % VO_{2peak} Percentage of Peak Oxygen Consumption, WR Work-to-Rest ratio. *Duplicate data included only once.

3.2 Characteristics of the Included Studies

Table II presents the characteristics of the included articles. All articles were analytical cross-sectional studies. Most of the research originated in Spain [15, 19, 20, 43-51]. The sample sizes of the included papers ranged from 4 to 64 players, with 10 out of 24 studies (42%) having 10 participants or fewer. Most papers focused on match data from international players, with three studies specifically examining data obtained from Paralympic athletes [45, 49, 50]. Only two studies included players at the national level [14, 42].

Of the 24 articles, 16 included only male participants [14, 15, 17, 37, 39-41, 43, 44, 46-51, 54], two studies had equal representation of male and female players [19, 20], and one study focused solely on female participants [45]. In the remaining five studies, male players constituted the majority [18, 38, 42, 52, 53]. The open class was more prominently represented than the quad class, with only two studies providing data from quad matches [18, 52].

3.3 Methodological Quality Assessment

Table III. Joanna Briggs Institute (JBI) checklist score (%) of the studies included in this review (n=24).

Author(s) (year)	Item number and corresponding score								Yes	No	Unclear	JBI Checklist Score
	1	2	3	4	5	6	7	8				
Bernardi et al. (2010)	N	Y	Y	Y	Y	N	Y	Y	6	2	0	6
Croft et al. (2010)	U	Y	Y	Y	Y	Y	Y	Y	7	0	1	7
Filipic et al. (2009a)	N	Y	Y	Y	N	N	Y	Y	5	3	0	5
Filipic et al. (2009b)	N	Y	Y	Y	N	N	U	Y	4	3	1	4
Gómez (2021)	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	7
Mason et al. (2020)	N	Y	Y	Y	Y	N	Y	Y	6	2	0	6
Ponzano et al. (2017)	N	Y	Y	Y	N	Y	Y	Y	6	2	0	6
Roy et al. (2006)	U	Y	Y	Y	Y	Y	Y	Y	7	0	1	7
Sánchez-Pay et al. (2013)	Y	N	Y	N	N	N	Y	Y	4	4	0	4
Sánchez-Pay et al. (2015a)	Y	N	Y	N	N	N	Y	Y	4	4	0	4
Sánchez-Pay et al. (2015b)	Y	N	Y	N	Y	Y	Y	N	5	3	0	5
Sánchez-Pay et al. (2015c)	Y	N	Y	N	Y	Y	Y	Y	6	2	0	6
Sánchez-Pay et al. (2015d)	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	8
Sánchez-Pay et al. (2016)	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	8
Sánchez-Pay et al. (2017a)	N	N	Y	N	Y	N	Y	Y	4	4	0	4
Sánchez-Pay et al. (2017b)	Y	N	Y	N	N	N	Y	Y	4	4	0	4
Sánchez-Pay et al. (2017c)	Y	N	Y	N	Y	N	Y	Y	5	3	0	5
Sánchez-Pay et al. (2021a)	U	N	Y	N	Y	N	Y	Y	4	3	1	4
Sánchez-Pay et al. (2021b)	Y	N	Y	N	Y	Y	Y	Y	6	2	0	6
Sánchez-Pay et al. (2023)	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	8
Sindall et al. (2013a)	N	N	Y	N	Y	N	Y	Y	4	4	0	4
Sindall et al. (2013b)	N	Y	Y	Y	Y	N	Y	Y	6	2	0	6
Sindall et al. (2015)	N	U	Y	Y	Y	Y	Y	Y	7	1	0	7

Vd Slikke et al. (2020)	N	U	Y	U	Y	N	Y	Y	4	2	2	4
Number of studies applying the item	11	10	24	13	18	11	23	23				

Questions from the JBI Checklist: 1. Were the criteria for inclusion in the sample clearly defined? 2. Were the study subjects and the setting described in detail? 3. Was the exposure measured in a valid and reliable way? 4. Were objective, standard criteria used for measurement of the condition? 5. Were confounding factors identified? 6. Were strategies to deal with confounding factors stated? 7. Were the outcomes measured in a valid and reliable way? 8. Was appropriate statistical analysis used? *N* No, *U* Unclear, *Y* Yes

Among the 24 studies assessed, fewer than half (11, 46%) clearly defined the criteria for inclusion (item 1), and described the participants and setting in detail (item 2). All studies measured the exposure (i.e., tennis match play) validly and reliably (24/24, 100%) (item 3). Just over half of the studies (13/24; 54%) used objective and standard criteria for measuring the condition (i.e., the underlying health condition or physical impairment) (item 4). Three-quarters of the studies adequately identified the confounding variables (18/24; 75%, item 5), but only 11 out of 24 (46%) stated the strategies used to manage them (item 6). Most studies used valid and reliable outcome measures and appropriate statistical analysis (23/24, 96%, items 7 and 8).

3.4 Quantitative Synthesis (Meta-analysis)

The data are presented as means or mean differences with 95% Confidence Intervals (95%CI), and 95% Prediction Intervals (95%PI) (if available). The datasets used for analysis are available in **Online Resource 2**, and the meta-analysis results are displayed in Tables IV to VI and **Online Resource 3**. The main findings are summarised in Figure 2. Authors Filipcic [40], Sánchez-Pay [15, 20, 50], Sindall [53], van der Slikke [54], van der Slikke (on behalf of Mason)[18], and Tolfrey (on behalf of Croft)[38] responded to our requests to provide missing data or clarification.

3.4.1 Sensitivity analysis

The sensitivity analysis with varying correlation parameters ($\rho = 0, 0.2, 0.4, 0.6, 1.0$) showed that none of the analyses influenced the results (**Online Resources 4 and 5**).

3.4.2 Time characteristics

The results of the meta-analysis of the time characteristics are presented in Table IV.

Table IV. Time characteristics

Outcome (Court type, Sex, Playing Level, Division)	Studies included in the meta-analysis (author, year)	Number of Studies (n)	Mean [95% - CI]	95% - Prediction Interval	Tau	I ²
Match Duration (Hard, Male, International, Open)	Filipic 2009a, Sánchez-Pay 2021a, Sánchez-Pay 2021b, Mason 2020, Sánchez-Pay 2017b, Sánchez-Pay 2015a, Sánchez-Pay 2013, Sindall 2013b	8	65.86 min [55.04; 78.82]	[41.55; 104.42]	0.43	85.78
Match Duration (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2016, Sánchez-Pay 2013	3	77.65 min [49.04; 122.96]		0.36	71.09
Match duration (Hard, Female, International, Open)	Mason 2020, Sánchez-Pay 2017b	2	67.66 [13.29; 345.64]		0.34	42.62
Set Duration (Hard, Male, International, Open)	Mason 2020, Sánchez-Pay 2021a, Sánchez-Pay 2021b, Sánchez-Pay 2017b, Sánchez-Pay 2015b, Sindall 2013b	6	35.04 min [28.21; 43.53]	[19.98; 61.4]	0.45	86.09
Set Duration (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2016, Sánchez-Pay 2015b	3	35.78 min [29.22; 43.80]		0.20	26.09
Set Duration (Hard, Female, International, Open)	Mason 2020, Sánchez-Pay 2017b, Sánchez-Pay 2015b	3	34.79 min [32.54; 37.21]		0.21	30.02
Game Duration (Hard, Male, International, Open)	Sánchez-Pay 2021a, Sánchez-Pay 2021b	2	4.63 min [0.92; 23.28]		0.41	67.49
Rally Duration (Hard, Male, International, Open)	Filipic 2009a, Sánchez-Pay 2021b, Sánchez-Pay 2017b, Sánchez-Pay 2015a	4	6.14 s [3.68; 10.24]		0.57	99.83

Outcome (Court type, Sex, Playing Level, Division)	Studies included in the meta-analysis (author, year)	Number of Studies (n)	Mean [95% - CI]	95% - Prediction Interval	Tau	I ²
Effective Playing Time (%) (Hard, Male, International, Open)	Filipic 2009a, Sánchez-Pay 2021b, Sánchez-Pay 2017b, Sánchez-Pay 2015a	4	19.81% [18.93; 20.74]		0.00	0.00
Effective Playing Time (%) (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2016	2	17.65% [17.54; 17.76]		0.00	0.00
Work Rest Ratio (Hard, Male, International, Open)	Sánchez-Pay 2015a, Sánchez-Pay 2021b, Sánchez-Pay 2017b	3	1: 4.05 [1:3.71; 1:4.41]		0.00	0.00

95% - CI 95% Confidence Interval, I² I-square statistic, Min Minutes, 95%-PI 95% Prediction Interval, Tau Kendal's Tau

3.4.2.1 Match duration

The average duration of matches played by international male players on hard court was 65.9 minutes (min), 95%CI [55.0, 78.8], and a 95% PI of [41.6, 104.4] (Table IV). The average match duration on clay court was 77.7 min, 95%CI [49.0, 123.0]. The average duration of matches of female players on hard court was 67.8 min, 95%CI [13.3, 345.6] (two studies). There was not enough information to conduct a meta-analysis for match duration on grass courts for male players, clay or grass courts for female players, or quad players.

3.4.2.2 Set duration

The average set duration of international males in the open division on hard court was 35.0 min with a 95%CI of [28.2, 43.5] and a 95% PI of [20.0, 61.4], and on clay court, it was 35.8 min with a 95% CI of [29.2, 43.8]. For female players, the average set duration on hard court was 34.8 min with a 95% CI of [32.5, 37.2]. There was not enough data to conduct a meta-analysis for set duration for male players on grass courts, female players on clay or grass courts, or quad players.

3.4.2.3 Game duration

The average game duration of international male players in the open division on hard court was 4.6 min 95%CI [0.92, 23.3]. There was insufficient information to conduct a meta-analysis for game durations on clay and grass courts, or for female and quad players.

3.4.2.4 Rally duration

The average rally duration of international male players in the open division on hard court was 6.1 seconds, with a 95%CI of [3.7, 10.2]. There was insufficient information to conduct a meta-analysis for rally durations on clay and grass courts, or for female and quad players.

3.4.2.5 Effective playing time and work-to-rest ratio

Effective playing time for international male players on hard court was 19.8% with a 95%CI of [18.9, 20.7], and on clay court 17.7% with a 95% CI of [17.5, 17.8] (two studies). The work-to-rest ratio was 1:4.1 with a 95%CI of [1:3.7, 1:4.4] on hard court and 4.1 with a 95%CI [3.6, 5.6] on clay court (two studies). Insufficient information was available for a meta-analysis of effective playing time of male players on grass court or for female and quad players.

3.4.3 On-court movement

The results of the meta-analysis of the on-court movement are presented in Table V.

Table V. On-court movement

Outcome (court type, sex, playing level, division)	Studies included in the meta-analysis	Number of studies (n)	Mean [95% - CI]	95% - PI	Tau	I ²
Distance per match (Hard, Male, International, Open)	Mason 2020, Sindall 2013a, Sindall 2013b	3	3859.22 m [1917.19; 7768.45]		0.34	42.62
Distance per set (Hard, Male, International, Open)	Mason 2020, Sindall 2013b	2	1900.48 m [229.82; 15715.86]		0.47	82.46
Average Forward Speed (Hard, Male, International, Open)	Filipic 2009b, Mason 2020, Sindall 2013a, Sindall 2013b, vd Slikke 2020	5	1.06 m·s ⁻¹ [0.85; 1.32]	[0.62; 1.81]	0.42	96.49
Peak Forward Speed (Hard, Male, International, Open)	Filipic 2009b, Mason 2020, Sindall 2013a, Sindall 2013b, vd Slikke 2020	5	3.55 m·s ⁻¹ [2.92; 4.31]	[2.31; 5.45]	0.37	95.75

95%-CI 95% Confidence Interval, I² I-square statistic, 95%-PI 95% Prediction Interval, *Tau* Kendal's Tau

3.4.3.1 Accelerations, decelerations, and turns

Insufficient information was available to conduct a meta-analysis of the number of accelerations, decelerations, and turns during wheelchair tennis matches.

3.4.3.2 Distance covered

The average distance per match by international male players in the open division on hard court was 3859 m, with a 95% CI of [1917, 7768]. Insufficient information was available for a meta-analysis of distance covered per match on clay and grass courts, or for female and quad players.

The average distance covered per set by international males on hard court in the open division was 1900 m, with a 95% CI of [230, 15716] (two studies). A meta-analysis was not possible for distance covered per set on clay and grass courts, or for female and quad players, due to insufficient data.

There was also insufficient information for a meta-analysis of distance covered per stroke, point, game, set, or speedzone/minute.

3.4.3.3 Movement speed

The average forward speed of international male wheelchair tennis players on hard court was $1.06 \text{ m}\cdot\text{s}^{-1}$, with a 95% CI of [0.85, 1.32], and a 95% PI of [0.62, 1.81]. The mean peak forward speed was $3.55 \text{ m}\cdot\text{s}^{-1}$ with a 95%CI of [2.92, 4.31], and a 95% PI of [2.31, 5.45]. There was insufficient information to conduct a meta-analysis of the peak and average forward speeds for female and quad players.

3.4.4 Physiological match variables

3.4.4.1 Heart rate

The average and peak heart rates of international male players on hard court were 134.3 [124.2, 1451.1] and 166.0 [132.7, 207.6] beats per minute. Expressed as a percentage of peak heart rate, the average match heart rate was 75% with a 95% CI of [46.4, 100]. A meta-analysis was not possible for the average heart rates of male players on clay and grass courts, for female and quad players, and for peak heart rates due to insufficient information.

3.4.4.2 Oxygen uptake and energy expenditure

A meta-analysis of oxygen uptake and energy expenditure of wheelchair tennis players during matchplay could not be performed due to insufficient data.

3.4.5 Stroke performance (speed)

The lack of available information prevented a meta-analysis on the stroke speed of wheelchair tennis players during matchplay.

3.4.6 Stroke performance (number)

The results of the meta-analysis of the stroke performance are presented in Table VI.

Table VI. Stroke performance (number) and scoring

Outcome (Court type, Sex, Playing Level, Division)	Studies included in the meta-analysis (author, year)	Number of Studies	Mean [95% - CI]	95% - Prediction Interval	Tau	I ²
First Serve Percentage (Hard, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2015b, Sánchez-Pay 2013, Sánchez-Pay 2017c	4	64.56% [57.77; 72.15]		0.12	5.80
First Serve Percentage (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2015b, Sánchez-Pay 2013	3	66.28% [61.34; 71.61]		0.00	0.00
First Serve Percentage (Hard, Female, International, Open)	Sánchez-Pay 2015b, Sánchez-Pay 2015c	2	59.3 % [57.90; 60.72]		0.18	35.77
Strokes per Match (Hard, Male, International, Open)	Sánchez-Pay 2021a, Sánchez-Pay 2021b, Sánchez-Pay 2017a, Sánchez-Pay 2015a	4	365.91 [317.21; 422.07]		0.00	0.00
Strokes per Match (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2015d	2	405.01 [24.51; 6691.3]		0.54	86.65
Strokes per Set (Hard, Male, International, Open)	Sánchez-Pay 2021a, Sánchez-Pay 2021b, Sánchez-Pay 2017b, Sánchez-Pay 2015a	4	200.64 [134.65; 298.97]		0.46	89.33
Strokes per Game (Hard, Male, International, Open)	Sánchez-Pay 2021a, Sánchez-Pay 2021b, Sánchez-Pay 2015a	3	25.4 [16.68; 38.68]		0.45	93.74
Strokes per Rally (Hard, Male, International, Open)	Mason 2020, Sánchez-Pay 2015a, Sánchez-Pay 2021b, Sánchez-Pay 2017b	4	3.43 [2.56; 4.60]		0.32	98.54
Strokes per Rally (Hard, Female, International, Open)	Mason 2020, Sánchez-Pay 2017b	2	3.10 [3.10; 3.10]		0.00	0.00

Outcome (Court type, Sex, Playing Level, Division)	Studies included in the meta-analysis (author, year)	Number of Studies	Mean [95% - CI]	95% - Prediction Interval	Tau	I ²
Points per Match (Hard, Male, International, Open)	Filipic 2009a, Sánchez-Pay 2021a, Sánchez-Pay 2015a, Sánchez-Pay 2021b	4	126.19, [93.2; 170.85]		0.44	85.94
Points per Set (Hard, Male, International, Open)	Mason 2020, Sánchez-Pay 2021a, Sánchez-Pay 2021b, Sánchez-Pay 2017b, Sánchez-Pay 2015a, Sánchez-Pay 2015b	6	57.95 [51.66; 65.01]	[45.36; 74.05]	0.29	60.38
Points per Set (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2015b, Sánchez-Pay 2015d	3	58.85 [42.87; 80.79]		0.33	71.77
Points per Set (Hard, Female, International, Open)	Mason 2020, Sánchez-Pay 2015b, Sánchez-Pay 2017b	3	57.64 [52.02; 63.86]		0.00	0.00
Points per Game (Hard, Male, International, Open)	Sánchez-Pay 2021a, Sánchez-Pay 2021b, Sánchez-Pay 2015a, Sánchez-Pay 2015b	4	6.56 [6.26; 6.87]		0.29	90.48
Points per Game (Clay, Male, International, Open)	Sánchez-Pay 2021b, Sánchez-Pay 2015b, Sánchez-Pay 2015d	3	7.07 [5.04; 9.93]		0.24	67.68

95%-CI 95% Confidence Interval, I² I-square statistic, *Tau* Kendal's Tau

3.4.6.1 First serve percentage

The percentage of accurate 1st serves for international male players on hard court was 64.6% with a 95% CI of [57.8, 72.2]. It was 66.3% on clay court with a 95% CI of [61.3, 71.6]. For female players on hard court, the average percentage of 1st serves was 59.3% with a 95% CI of [57.9, 60.7].

3.4.6.2 Strokes per match, set, game, and rally

The average number of strokes per match of international male players on hard court was 365.9, with a 95% CI of [317.2, 422.1], and on clay court, it was 405.0 with a 95% CI of [24.5, 6691.3]. Their average number of strokes per set on hard court was 200.6, with a 95% CI of [134.7, 299.0], per game 25.4, with a 95% CI of [16.7, 38.7], and per rally 3.4, with a 95% CI of [2.6, 4.6]. For female players, the average number of strokes per rally was 3.1, with a 95% CI of [3.1, 3.1].

There was insufficient information available to conduct a meta-analysis for the average number of strokes per rally, game, and set on clay and grass courts, as well as for female and quad players.

3.4.7 Scoring

3.4.7.1 Points per match, set, and game

The average number of points per match by international male players on hard court was 126.2, with a 95% CI of [93.2, 170.9]. Their average number of points per set on hard court was 58.0, with a 95% CI of [51.7, 65.0], and on clay court 58.9, with a 95% CI of [42.9, 80.8]. Their average number of points per game on hard court was 6.6, with a 95% CI of [6.3, 6.9], and on clay court, it was 7.1 with a 95% CI of [5.0, 9.9].

For international female players on hard court, the average number of points per set was 57.6, with a 95% CI of [52.0, 63.9].

There was insufficient information for a meta-analysis of the average number of points per game, set, and match on all surfaces for female and quad players, except for the number of points per set for females on hard court. Moreover, there was insufficient information for meta-analysis of the number of games per match and set, and for the number of sets per match.

3.5 Qualitative (descriptive) synthesis

Several outcome parameters, such as rotational speed, turns per minute, accelerations, decelerations, heart rate, oxygen consumption, energy expenditure, and blood lactate, had to be excluded from the meta-analysis due to a lack of insufficient data. The primary findings are outlined below (for additional details, please refer to Online Resource 2). Additionally, results from studies directly comparing open class men, women and quads in a single study, or those examining court surface variations, could not undergo meta-analysis due to insufficient available data, and are presented separately below.

3.5.1 Time characteristics

3.5.1.1 Rally, set and match duration

Sánchez-Pay et al.[20] compared rally, set, and match duration of eight male and eight female wheelchair tennis players in an international hard court tournament in the open division. The average match duration, expressed as mean (SD), was 64.7 (13.4) min for males, and 77.6 (31.7) min for females. The mean set duration was 30.4 (8.0) min for males, and 34.5 (9.1) min for females. These differences were not statistically significant. There was a statistically significant difference in

rally duration between male and female players, with an average rally duration of 5.58s (3.65) in male players and 6.82s (4.83) in female players ($p=.001$).

Mason et al.[2020] compared set duration between open male and female players, and quad players during the 2017 NEC Wheelchair Tennis Masters on hard court. The study included the best internationally ranked players: seven male players who played 21 sets, six female players who played 17 sets, and four quad players who played 11 sets. Set duration in male players was 39 min:18s (12:22), in female players 34:01 (10:43), and quad players 28:50 (08:07), showing statistically significant differences between open male vs quad, and open female vs quad.

3.5.1.2 Rally pace

Sánchez-Pay et al.[20] compared rally pace between male and female matches on hard court. Male players averaged a rally pace of one stroke every 1.9 (0.38) s, while female players averaged one stroke every 2.15 (0.41) s [20]. In another study, Sánchez Pay et al.[46] measured rally pace in four male players on clay court, and found that they averaged one stroke every 2.2 s.

3.5.1.3 Effective playing time and work-to-rest ratio

Sánchez-Pay et al.[20] compared effective playing time and work-to-rest ratio in male and female players on hard court. They reported an effective playing time of 20.8 (3.4) % in male players, and 22.3 (5.2) % in female players. For the work-to-rest ratio, the numbers were 1:3.95 (1:0.86) in male players, and 1:3.72 (1:1.11) in female players. There was no data on effective playing time and work-to-rest ratio in quad players.

3.5.2 On-court movement

3.5.2.1 Accelerations and decelerations (males vs females vs quads; clay vs hard court)

Two studies examined acceleration in wheelchair tennis. Mason et al.[18] investigated average forward and rotational acceleration on hard courts in international male and female open class and quad players. The mean forward acceleration, measured in $\text{m}\cdot\text{s}^{-2}$, was 1.08 (0.18) for male players, 0.91 (0.15) for female players, and 0.72 (0.08) for quad players. Similarly, the mean rotational acceleration, measured in $\text{deg}\cdot\text{s}^{-2}$, was 205 (28) for male players, 211 (29) for female players, and 144 (31) for quad players. The differences between the open division and the quad division were statistically significant ($p<0.0005$).

In the study by Ponzano & Gollin[42], the mean maximum acceleration and deceleration were compared on clay and hard courts. They studied a group consisting of ten male and two female wheelchair tennis players. The mean maximum acceleration on clay and hard courts was 1.5 (0.2) $\text{m}\cdot\text{s}^{-2}$. The mean maximum deceleration in the same group of players was -1.68 (0.27) $\text{m}\cdot\text{s}^{-2}$ on clay court and -1.63 (0.32) $\text{m}\cdot\text{s}^{-2}$ on hard court.

3.5.2.2 Turns

In wheelchair tennis, turns serve the same purpose as direction changes in standing tennis. The study conducted by Mason et al.[18] examined the frequency of turns per minute and categorised them into turns towards the racket hand and turns towards the non-racket hand. The results showed that male players had an average of 5.6 (0.5) turns towards the racket hand, while female players had 6.2 (1.1) turns per minute. Quad players averaged 4.9 (0.8) turns per minute towards the racket hand. Similarly, the number of turns per minute towards the non-racket hand was 7.5 (0.7) for male players, 7.2 (1.0) for female players, and 6.8 (0.6) for quad players. The differences between the open division and the quad division were statistically significant ($p<0.0005$ for turns to the racket hand, and $p=0.016$ for turns to the non-racket hand).

3.5.2.3 Distance covered during a set

Mason et al.[18] used an indoor tracking system and inertial measurement units to measure the distance covered during an average set of open class male and female players and quads on hard court. This system has been investigated previously and been shown to provide valid and reliable data [55]. They found that male players covered the longest distance (2220 m), followed by female players (1840 m), with quad players covering the shortest distance (1275m).

Sindall et al.[17] measured the distance covered per match on hard court of international male tennis players in the open and quad class by comparing the use of a General Positioning System (GPS) and a data logging device. Distance covered by open class players measured by GPS was 2891 (1000) m vs 3963 (1340) m measured by a data logging device. This difference was statistically significant ($p=0.0001$). In quad players, these numbers were 2675 (438) m and 3931 (505) m respectively ($p=0.0001$).

Filipic et al.[40] reported on distance covered per point, game and match of 15 male players who played 22 singles matches on hard court, using video-analysis with the SAGIT/TENNIS tracking system. They found a distance of 6.1 (2.0) m per point (rally), 26.2 (26.9) m per game and 613.0 (273.2) m per match. It should be noted that they only measured distance covered during the active phase of play. Gómez [41] measure the distance per point and per match of 8 international male tennis players in the open division during 10 matches on clay court, and compared the distance covered during the active phase of the match and during the total match. Distance per point was 7.8 (8.4) m, distance during active play 633.7 (266.8) m, and during the whole match 3372.8 m.

No studies reported on distance per game or per point (rally) in female or quad players.

3.5.2.4 Movement speed

Sindall et al. [17] compared forward movement speed using GPS and a data logging device of open and quad class players on hard court. The average and peak speed of open class players using GPS was 0.8 (0.1) $\text{m}\cdot\text{s}^{-1}$ and 3.5 (0.4) $\text{m}\cdot\text{s}^{-1}$, and 1.0 (0.2) and 3.3 (0.6) $\text{m}\cdot\text{s}^{-1}$ using data logging. In quad players, these numbers were 0.6 (0.1) and 3.1 (0.2) $\text{m}\cdot\text{s}^{-1}$ for GPS, and 0.9 (0.0) and 2.8 (0.4) $\text{m}\cdot\text{s}^{-1}$ for data logging.

Van der Slikke et al.[54] measured average and peak rotational speed in international male wheelchair tennis players on hard court (open class). They recorded a mean peak rotational speed of 369 (79) $\text{deg}\cdot\text{s}^{-1}$. They did not measure female or quad players.

3.5.3 Physiological match variables

3.5.3.1. Heart rate

Sindall et al.[17] measured a peak heart rate of 167 (22) beats per minute in 12 male players on hard court. Sánchez-Pay et al.[47] studied four male players on clay court during simulated match play, and their average and peak heart rates were 124.3 (24.7) and 163.3 (11.6) beats per minute. Croft et al.[38] reported an average and peak heart rate of 146 (16) and 180 (18) beats per minute in four males and two females on hard court. Ponzano et al.[42] compared average and peak heart rate on clay and hard court in a group of ten males and two females. Average and peak heart rates on hard court were 123 (14) and 159 (16), and on clay court, 123.3 (18.4) and 158.6 (18.4) beats per minute.

3.5.3.2 Oxygen uptake

Three studies examined oxygen uptake in wheelchair tennis. Bernardi et al.[37] calculated average VO_2 in four male paralympic wheelchair tennis players during two tennis matches. This calculation

was based on their peak VO_2 achieved during an incremental arm cranking exercise and heart rate monitoring during simulated match play. The average VO_2 during play was recorded as $24.2 (2.6) \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ and $1.7 (0.18) \text{ L}\cdot\text{min}^{-1}$, equivalent to 73% of their peak VO_2 .

Croft et al.[38] investigated the peak VO_2 in four male and two female international wheelchair tennis players on a hard court. The players performed submaximal and VO_2 peak tests in their sport-specific wheelchairs on the treadmill. Heart rate was monitored during competition, and the researchers calculated the average and peak VO_2 during play using linear regression equations. The average VO_2 was determined to be $1.36 (0.42) \text{ L}\cdot\text{min}^{-1}$, equalling 68% of their peak VO_2 .

In a study conducted by Roy et al.[14], the average VO_2 of six skilled wheelchair tennis players was calculated. This estimation was based on their VO_2 levels during an arm crank ergometer test and heart rate monitoring during simulated match play. Each player participated in two matches on hard court. The average estimated VO_2 during match play was $27.1 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$, equalling 49.9% of the peak VO_2 achieved during the ergometer test.

3.5.3.3 Energy expenditure

Two studies investigated energy expenditure during a wheelchair tennis match. Ponzano & Gollin[42] compared the energy expenditure of ten male and two female tennis players on clay and hard courts. Energy expenditure was $3.7 (0.9) \text{ kcal}\cdot\text{kg}^{-1}$ per match on clay courts and $3.3 (0.8) \text{ kcal}\cdot\text{kg}^{-1}$ per match on hard courts. This corresponds to approximately 238 (68) and 217 (78) kcal per match.

Roy et al.[14] studied the physiological responses of six skilled male wheelchair tennis players during two tennis matches on hard court. The participants completed an arm crank ergometer test that measured heart rate via a telemetry device and VO_2 peak via open circuit spirometry. He estimated their energy expenditure based on their heart rate during the match and found a mean value of 365 kcal per match.

3.5.3.4 Blood lactate levels

Sánchez-Pay et al.[47] studied blood lactate levels during simulated match play of four wheelchair tennis players (three matches each). Capillary blood was extracted from the earlobe during the changeovers after each of the unequal number of games, i.e., 1st, 3rd, and 5th game and so on, until the end of each match. The mean blood lactate concentration was $1.41 \text{ mmol}\cdot\text{L}^{-1} (0.43)$, based on 55 samples.

3.5.4 Stroke performance (number)

3.5.4.1 Strokes per rally (males vs females vs quads)

Two studies compared the number of strokes in different player groups. Sánchez-Pay et al.[20] compared the number of strokes per rally between male and female players on hard court. They found that male players had an average of $3.9 (1.7)$ strokes per rally, and female players had an average of $3.1 (2.0)$ strokes per rally. Mason et al.[18] also investigated this aspect and reported that male players had an average of $3.1 (0.5)$ strokes per rally, while female players had a similar average of $3.1 (0.8)$ strokes per rally. For quad players, the average was $2.5 (0.5)$ strokes per rally.

3.5.4.2 Strokes per set (males vs females, hard vs clay vs grass court)

Sánchez-Pay et al.[20] compared the number of strokes per set between male and female players, and found an average of $157.2 (31.1)$ strokes per set for male players and $180.5 (77.8)$ for female players.

Sánchez-Pay et al.[15] compared the number of strokes per set for male players on different court surfaces. The study found that male players had an average of 272.9 (92.1) strokes per set on hard court, 206.0 (28.6) on clay court, and 184.2 (47.1) on grass court.

3.5.5 Scoring

3.5.5.1 Games per set

Sánchez-Pay et al.[20] presented findings on the average number of games per set played by male and female international wheelchair tennis players during the 2014 Australian Open (hard court), Roland Garros (clay court), and US Open tournaments (hard court). The results indicated that male players averaged 8.8 (2.4) games per set at the Australian Open, 8.9 (2.1) at Roland Garros, and 9.8 (2.4) at the US Open. Female players averaged 8.3 (1.3) games per set at the Australian Open, 8.1 (1.1) at Roland Garros, and 9.6 (2.2) at the US Open.

3.5.5.2 Bounces per point

In this same study conducted by Sánchez-Pay et al.[20], the researchers also examined the number of bounces per point. The average number of bounces per point was reported as 2.46 (1.85) for male players and 2.97 (2.53) for female players.

3.5.5.3 Shots off one bounce

Mason et al.[18] studied the percentage of shots played off a single bounce. Their results revealed that male players hit 85.7% (6.8) of their shots after a single bounce, while female players hit 82.1% (7.6) off one bounce. Similarly, quad players hit 81.5% (6.2) after one bounce.

Sánchez-Pay et al.[15] studied the percentage of shots played after one, two or no bounce on different court surfaces. There was a statistically significant difference between court surfaces for shots played without bounce, with higher values on grass court (4.6%) than on hard (3.0%) or clay court (2.2%). No statistical significant differences were found between court surfaces for shots played after one or two bounces.



Figure 2. A summary of the physical demands of wheelchair tennis singles in open class international male players on hard court.

4.0 Discussion

This systematic review aimed to assess the physical demands of open and quad class male and female wheelchair tennis players, playing on various surfaces, by analysing existing literature and examining key factors such as play duration, on-court movement, stroke performance, and physiological variables during matches. We provide insights into the duration of matches, sets, and rallies, effective playing time, and work-to-rest ratio. We describe stroke performance, including the first-serve percentage, and on-court wheelchair movement, including distance covered, forward speed, accelerations, decelerations, and turns. Finally, we consider physiological variables like heart rate, oxygen uptake, and energy expenditure during matches. The compiled data can guide players, coaches and support staff when customising training programs and devising effective match strategies. Our analysis provides a baseline of knowledge for future work as more data becomes available on different court surfaces.

4.1. Time characteristics

Matches in wheelchair tennis, particularly for international male players in the open class on hard court, were generally shorter (approximately 65 mins) than matches in standing tennis (approximately 90 min).[21] The opponents' skill and competition depth play a crucial role in match duration. Evenly matched players are more likely to engage in extended points, games, and sets, resulting in longer matches. At the moment, wheelchair tennis does not possess the same level of competition depth as standing tennis, and matches are shorter, particularly in the early rounds of a tournament.

However, given that only nine studies analysed the duration of wheelchair tennis matches, as well as the variety of playing conditions (indoor vs outdoor, official vs simulate, two sets plus supertiebreak vs best of three sets, court pace or type of ball among others), it is difficult to draw more detailed conclusions. There was a lack of research focusing on female players, quad players, juniors, and lower-level players, which limits a comprehensive understanding of match duration across different categories.

On average, male and female wheelchair tennis players had similar set duration, around 35 minutes. These durations were generally shorter than those typically seen in ATP and WTA players during Grand Slam events.[19] Variability in set duration seemed to be a function of the number of games per set, a relationship that has also been observed in standing tennis [21].

Wheelchair tennis players had a slower pace of play than standing tennis players. Despite having slightly longer rally durations, the number of strokes per rally was lower in wheelchair tennis, pointing to an overall slower pace of the game. Mason et al.[18] found that male wheelchair tennis players hit 86% of their shots after the first bounce, in contrast to 82% by female and quad players. Most winners were hit after one bounce (91%, 92%, and 88% respectively), indicating that the second bounce is primarily used in defensive situations. This finding seems to contrast with a prior study that found more frequent and arguably tactical use of the second bounce by higher-ranked players compared to their lower-ranked counterparts [56]. Further dissecting the strategic interplay between ball bounce and other game context such as the players' court position, use of forehand/backhand and the trajectory of the shots, which are currently missing from the literature, will provide even more practical insight to interpret rally dynamics. Regardless, it is interesting that the effective playing time was similar in wheelchair and standing tennis, hovering around 20%. Nonetheless, the shorter rally duration, the slower rally pace as well as the use of the second bounce

show that wheelchair tennis is tactically different from standing tennis, so it is important to design training programs specifically for this sport.

Comparing the time and game characteristics regarding playing surface in several studies showed relatively small differences between both sexes[15, 19, 43].

4.2 Stroke performance

At the Grand Slams, the first-serve percentages (the proportion of accurate first serves) of wheelchair tennis players were similar to standing tennis players[19]. However, the percentage of points won on first serves was approximately 10-15% lower in wheelchair tennis players. Serve speed and placement of the ball are important factors that determine the success rate of the stroke. The lower success rate of first serves in wheelchair tennis can be attributed to three primary factors. Firstly, wheelchair tennis players have a lower hitting plane compared to standing players [57, 58], meaning their serves cannot feature the same horizontal velocity profile. Secondly, wheelchair tennis players generate less force due to the absence of leg drive, resulting in lower serve speed [59]. Thirdly, wheelchair tennis players serve in a static position, making it difficult to overcome the inertia to move and hit the next ball from an optimal position [60].

Regrettably, there was limited additional information regarding stroke performance in wheelchair tennis, including metrics such as stroke speed and the frequency of different stroke types, and further studies are eagerly anticipated.

4.3 On-court movement

On-court movement plays a pivotal role in a wheelchair tennis, and this review showed some interesting findings. Within the open division, male players covered the most ground per set, followed by their female counterparts, while quads covered the shortest distance[18]. This discrepancy in distance covered can be attributed, at least in part, to the greater forward acceleration and speed observed in male players within the open division. These differences may be influenced by the inherent upper body strength advantage of male open-class players compared to females and quad players, who, by definition, have at least one upper limb affected which may well impact their ability to manoeuvre the chair.

The extent of ground covered is related to the match's scoring dynamics. A closely contested match often leads to more and longer points, contributing to an overall increase in the distance covered by the players on the court.

Average speed and maximum speed, acceleration, and deceleration did not differ between hard and clay court in a study with 12 nationally ranked male tennis players and a counterbalanced design [42]. Playing surface did not affect the players' performance. Further studies, including on grass, are needed.

It is also important to consider the measurement methods employed in the study. Filipcic et al. [40] reported the lowest distance covered during a tennis match, which was 613 meters, using video analysis with the SAGIT/ TENNIS tracking software. These lower values can be attributed to the researchers' decision to solely consider movement during points played, unlike other studies [17, 51, 52] that encompassed the entirety of the match.

This hypothesis gains support from research by Gómez et al.[41], who examined ten matches from the XXIII National male wheelchair tennis masters in Spain. Gómez compared the distance covered during the active phase of play to the total match time, using wireless inertial movement units

(WIMU PROTM, RealTrack Systems, Spain). The distance covered during the active phase of play measured 634 meters, while during the total match time, it extended to 3373 meters.

Sindall et al.[52] compared the criterion validity and accuracy of 1 Hz GPS and a data logging device attached to the wheels. GPS values for distance and speed were consistently lower than equivalent values obtained by data logging. GPS with higher sampling rates than 1 HZ were recommended by the authors for wheelchair tennis, because the measurement device used should be valid and reliable.

There is room to more comprehensively examine the movement demands of wheelchair tennis. Validated technologies and standardisation of measurement are critical in this regards, and the increased use of optical tracking systems in Grand Slam tennis holds considerable promise to further develop an understanding of the sport.

4.4 Physiological variables

The studies conducted [14, 37, 38, 42, 47] reported an average heart rate ranging from 121 to 146 bpm, with peak heart rates spanning 158 to 180 bpm. The average match heart rate expressed as a percentage of peak heart rate ranged from 66-78%. Heart rate monitoring offers a means to infer the intensity of play, monitor training programs, and design effective training programs. However, while heart rate monitoring serves as a valuable tool to assess play intensity, there are a range of additional factors that can influence heart rate among wheelchair tennis players. These factors encompass age (with lower heart rates in older individuals) [61], the level of spinal cord injury (with lower heart rates in cases of higher spinal cord injury) [62], training level (linked to lower heart rates in individuals that are better trained) [63], and the extent of body muscle mass used (tied to lower heart rates when less body muscle mass is involved) [64].

Ponzano & Gollin[42] compared mean and peak heart rates as well as energy expenditure of 12 nationally ranked wheelchair tennis players on clay and hard courts, and found closely matched values on both playing surfaces. The estimated energy expenditure figures (approximately 220 kcal per match on hard court and 240 kcal on clay) were slightly lower than those reported by Roy et al.[14] for six competitive wheelchair tennis players on hard courts (350 kcal). Notably, Ponzano et al. derived their energy expenditure calculations based on GPS data and metabolic power, while Roy et al. based their estimations on oxygen uptake during an arm ergometer test and heart rate recordings during matchplay. These different methodological approaches complicate the direct comparison of findings between the two studies.

4.5 Comparison to other sports

Elite male wheelchair padel players covered shorter distances per set (around 320 m) than elite male wheelchair tennis players (around 1900 m)[65]. Wheelchair padel players also moved slower (around 0.54 m/s versus 1.1 m/s), and displayed a lower percentage of peak heart rate (65-68%) than wheelchair tennis players (75%), indicating a lower intensity of play in wheelchair padel. When comparing the activity demands of wheelchair tennis to wheelchair basketball, a systematic review showed that basketball players covered longer distances (5-6 km vs 4 km) in shorter time periods (40 vs 65 mins) at a higher percentage of peak heart rate (85 vs 75%), indicating that wheelchair basketball is played at a higher intensity than wheelchair tennis [66]. In a comparative analysis of wheelchair mobility performance among wheelchair basketball, tennis, and rugby players in a single study, the results indicated that wheelchair basketball players showed the highest average wheelchair mobility performance levels, while rugby displayed the lowest. Wheelchair tennis fell between the two sports for most outcomes[54].

4.5 Training recommendations

The compiled data can serve as a valuable resource for players and coaches in customising training programs and devising effective match strategies. Indeed, quantifying the demands of the game often represents an important initial step in best actioning the principles of specificity and overload in training settings. In a practical sense, by synthesising representative internal load such as expected heart rates in wheelchair tennis, off-court training can be shaped more specifically and so too can exercise testing [67]. Monitoring heart rate provides crucial insights into the player's response to training load and helps assess fatigue levels, aiding in early injury prevention and detection [68]. During practice, players should be able to easily cover a distance of at least 4 km in intervals. For individuals with a spinal cord injury below T5, it is recommended to strive to regularly achieve an intensity of at least 75% of peak heart rate. For those with a spinal cord lesion above T5, a preferred method of intensity monitoring would be RPE (rate of perceived exertion), and players should be able to maintain an intensity of 8/10.

Statistics related to match duration, on-court movement, and stroke performance are also valuable for simulating match play during training sessions and formulating effective tactics during tournaments. Training programs should prioritise maximising cardiovascular fitness, building strength, and incorporating wheelchair manoeuvring drills. Players should also commit training time to developing skills like quick turns and improving stroke efficiency, particularly in serving. Markers of external load like stroke count can provide an important guide for trunk and upper limb conditioning. With technological advance, it is likely that these markers will become more joint or limb specific in the future to further aid player preparation. By combining physiological data and performance statistics, players can tailor their physical, technical tactical, and mental programs to excel in wheelchair tennis.

4.6 Strengths and limitations

We thoroughly reviewed the existing literature regarding the physical demands of wheelchair tennis. One notable limitation is the scarcity of studies investigating the physical demands on clay and grass courts. Similarly, there is a lack of research focusing on female players, quad players, lower-level players, juniors, and doubles play.

Most existing studies have concentrated on the physical demands of international male wheelchair tennis players participating in the open class, specifically on hard courts. The limited number of studies available for each specific outcome variable often hindered the possibility of conducting meta-analyses, and resulted in wide confidence and prediction intervals. Additionally, we have concerns about the reliability and validity of certain measurement techniques used for on-court movement analysis within the included studies. This may impact the accuracy of the data collected and, in turn, influence the reliability of our conclusions relating to on-court movement.

5.0 Conclusion

Our understanding of the physical demands of wheelchair tennis is largely centered around international male players on hard courts in the open division. To provide a truly comprehensive understanding of the sport's physical requirements, future research should prioritise the inclusion of data on female and quad players, and matches played on clay and grass court surfaces. Such endeavours will be instrumental in facilitating more tailored and effective training programs for athletes and coaches within the wheelchair tennis community.

Similarly, there is the need for further research that details the movement strategies and anatomical contributions from body segments to produce the physical demands of wheelchair tennis, in a population of athletes with a wide range of impairments and physical function.

Supplementary information The online version contains supplementary material available at ...

DECLARATIONS

Funding TS obtained financial support from the Royal Netherlands Lawn Tennis Association to conduct the statistical analysis. The remaining authors did not receive any financial assistance for the research, authorship, or publication of this article.

CONFLICTS OF INTEREST

BMP and MGTJ are employed by the Royal Netherlands Lawn Tennis Association (KNLTB). CB is employed by the International Tennis Federation (ITF). BMP and SW are Heads of Classification of the ITF. CLA, NH, DCJvR, SM, MR, AJSP, TS, LJS, RBS, RvdS, and NW have no conflicts of interest that are directly relevant to the content of this article.

Ethics approval: not applicable.

Consent to participate: not applicable

Consent for publication: not applicable.

Availability of data and material All data generated or analysed during this study are included in this published article and its supplementary information files.

Author's contributions SW, BMP, MGTJ, SM and NH conceived the idea for the study, while CLA, CB, MR, AJSP, RBS, RvdS, and NW made substantial contributions to the study's design. LJS conducted the literature search with input from SW, MJ, and BMP. SW, BMP, and MGTJ screened the articles and extracted the data. TS analysed the data, conducted the statistical analysis and meta-analysis. SW, BMP, MGTJ, SM, and NH prepared the original draft. All authors helped critically revise the work for important intellectual content, approved the final version of the work, and agreed to be accountable for all aspects of the work.

REFERENCES

1. Committee IP. IPC guide to Para and IPC terminology. International Paralympic Committee; 2021. p. 16.
2. Harnett J. History of wheelchair tennis New York, NY: USTA; 2017 [Available from: <https://www.usta.com/en/home/stay-current/national/history-of-wheelchair-tennis.html>].
3. Richardson EV, Papathomas A, Smith B, Goosey-Tolfrey VL. The psychosocial impact of wheelchair tennis on participants from developing countries. *Disabil Rehabil*. 2017;39(2):193-200.
4. ITF. Wheelchair Tennis Classification Rules. London: International Tennis Federation 2023.
5. ITF. ITF Global Tennis Report 2021. London: International Tennis Federation; 2021.
6. Sánchez-Pay A. Determinant physical factors in the wheelchair tennis player. *ITF Coach Sport Sci Rev*. 2021;29(83):35-7.
7. Rietveld T. Wheeling performance in wheelchair tennis: understanding and improving a complex skill. Groningen: University of Groningen; 2022.
8. Sánchez-Pay A, Torres-Luque G, Sanz-Rivas D. Analysis of competitive wheelchair tennis. *ITF Coach Sport Sci Rev* 2014;22(63):22-4.
9. Moon HB, Park SJ, Kim AC, Jang JH. Characteristics of upper limb muscular strength in male wheelchair tennis players. *J Exerc Rehabil*. 2013;9(3):375-80.
10. Sánchez-Pay A, Martínez-Gallego R, Crespo M, Sanz-Rivas D. Key physical factors in the serve velocity of male professional wheelchair tennis players. *Int J Environ Res Public Health*. 2021;18(4).
11. Altmann VC, Hart AL, Vanlandewijck YC, van Limbeek J, van Hooff ML. The impact of trunk impairment on performance of wheelchair activities with a focus on wheelchair court sports: a systematic review. *Sports Med Open*. 2015;1(1):22.
12. Sindall P, Lenton JP, Mason BS, Tolfrey K, Cooper RA, Martin Ginis KA, et al. Practice improves court mobility and self-efficacy in tennis-specific wheelchair propulsion. *Disabil Rehabil Assist Technol*. 2021;16(4):398-406.
13. Rietveld T, Vegter RJK, van der Slikke RMA, Hoekstra AE, van der Woude LHV, de Groot S. Wheelchair mobility performance of elite wheelchair tennis players during four field tests: Inter-trial reliability and construct validity. *PLoS ONE*. 2019;14(6):e0217514.
14. Roy JL, Menear KS, Schmid MM, Hunter GR, Malone LA. Physiological responses of skilled players during a competitive wheelchair tennis match. *J Strength Cond Res*. 2006;20(3):665-71.
15. Sánchez-Pay A, Sanz-Rivas D. Physical and technical demand in professional wheelchair tennis on hard, clay and grass surfaces: implication for training. *Int J Perform Anal Sport*. 2021b;21(4):463-76.
16. Sánchez-Pay A, Sanz-Rivas D. Influence of hard, artificial grass and clay court surfaces on wheelchair tennis players' mobility-a case study. *Eur J Adapt Phys Activ*. 2021;14(2):10.
17. Sindall P, Lenton JP, Tolfrey K, Cooper RA, Oyster M, Goosey-Tolfrey VL. Wheelchair tennis match-play demands: effect of player rank and result. *Int J Sports Physiol Perform*. 2013b;8(1):28-37.
18. Mason BS, van der Slikke RMA, Hutchinson MJ, Goosey-Tolfrey VL. Division, result and score margin alter the physical and technical performance of elite wheelchair tennis players. *J Sports Sci*. 2020;38(8):937-44.
19. Sánchez-Pay A, Palao MJ, Torres-Luque G, Sanz-Rivas D. Differences in set statistics between wheelchair and conventional tennis on different types of surfaces and by gender. *Int J Perform Anal Sport*. 2015b;15(3):1177-88.
20. Sánchez-Pay A, Torres-Luque G, Sanz-Rivas D. Activity patterns in male and female wheelchair tennis matches. *Kinesiology* 2017b;49(1).
21. Pluim BM, Jansen MGT, Williamson S, Berry C, Camporesi S, Fagher K, et al. Physical demands of tennis across the different court surfaces, performance levels and sexes: a systematic review with meta-analysis. *Sports Med*. 2023;53(4):807-36.

22. Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160.
23. Otten R, de Vries R, Schoonmade L. Amsterdam Efficient Deduplication (AED) method (Version 1). Zenodo. 2019.
24. Bramer WM, Giustini D, de Jonge GB, Holland L, Bekhuis T. De-duplication of database search results for systematic reviews in EndNote. *J Med Libr Assoc*. 2016;104(3):240-3.
25. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016;5(1):210.
26. JBI. Critical appraisal tools Adelaide: JBI Collaboration; 2023 [Available from: <https://jbi.global/critical-appraisal-tools>].
27. McGrath S, Zhao X, Steele R, Thombs BD, Benedetti A. Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. *Stat Methods Med Res*. 2020;29(9):2520-37.
28. Tanner-Smith EE, Tipton E, Polanin JR. Handling complex meta-analytic data structures using robust variance estimates: a tutorial in R. *J Dev Life Course Criminol*. 2016;2(1):85-112.
29. Fisher Z, Tipton E. Robumeta: An R-package for robust variance estimation in meta-analysis. *arXiv*. 2015;1503.02220:1-16.
30. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177-88.
31. Harrer M, Cuijpers P, Furukawa TA, Ebert DD. Doing meta-analysis with R. A hands-on guide 1st ed: Chapman and Hall/CRC 2021 13 September 2021.
32. Friedrich JO, Adhikari NK, Beyene J. The ratio of means method as an alternative to mean differences for analyzing continuous outcome variables in meta-analysis: a simulation study. *BMC Med Res Methodol*. 2008;8:32.
33. IntHout J, Ioannidis JP, Rovers MM, Goeman JJ. Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open*. 2016;6(7):e010247.
34. R Core Team. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing 2020 [Available from: <https://www.r-project.org/>].
35. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw*. 2010;36(3):1 - 48.
36. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. Cochrane Handbook for Systematic Reviews of Interventions version 6.3 (updated February 2022). 2nd ed. Chichester (UK): John Wiley & Sons; 2019.
37. Bernardi M, Guerra E, Di Giacinto B, Di Cesare A, Castellano V, Bhambhani Y. Field evaluation of paralympic athletes in selected sports: implications for training. *Med Sci Sports Exerc*. 2010;42(6):1200-8.
38. Croft L, Dybrus S, Lenton J, Goosey-Tolfrey V. A comparison of the physiological demands of wheelchair basketball and wheelchair tennis. *Int J Sports Physiol Perform*. 2010;5(3):301-15.
39. Filipčič T, Filipčič A. Time characteristics in wheelchair tennis played on hard surfaces. *Kineziologija* 2009a;41(1):67-75.
40. Filipčič T, Filipčič A. Analysis of movement velocity and distance covered in wheelchair tennis. *Kinesiolog Slov*. 2009b;15(2):25-32.
41. Gómez CA. Kinematic parameters in wheelchair tennis players competition [Masters]. Murcia: Universidad de Murcia; 2021.
42. Ponzano M, Gollin M. Physical demand of wheelchair tennis match-play on hard courts and clay courts. *Int J Perform Anal Sport*. 2017;17(4):656-65.
43. Sánchez-Pay A, Torres-Luque G, Fernández-García ÁI, Sanz-Rivas D. Analysis of the effect of playing surface on men's singles wheelchair tennis. *Cult Cienc Deporte*. 2013;24(8):217-22.
44. Sánchez-Pay A, Sanz-Rivas D, Torres-Luque G. Match analysis in a wheelchair tennis tournament. *Int J Perform Anal Sport*. 2015a;15(2):540-50.

45. Sánchez-Pay A, Torres-Luque G, Manrique DC, Sanz-Rivas D, Palao JM. Match analysis of women's wheelchair tennis matches for the Paralympic Games. *Int J Perform Anal Sport*. 2015c;15(1):69-79.
46. Sánchez-Pay A, Sanz-Rivas D, Montiel A, Zanco Z, Torres-Luque G. Wheelchair tennis match activity pattern on clay court. *Retos*. 2015d;28:50-3.
47. Sánchez-Pay A, Torres-Luque G, Sanz-Rivas D. Match activity and physiological load in wheelchair tennis players: a pilot study. *Spinal Cord*. 2016;54(3):229-33.
48. Sánchez-Pay A, Torres-Luque G, Sanz-Rivas D. Stroke performance in high-level Spanish wheelchair tennis players. *Rev Int Cienc Deporte*. 2017a;13(48):139-48.
49. Sánchez-Pay A, Torres-Luque G, Fernández-García ÁI, Sanz-Rivas D, Palao JM. Differences in game statistics between winning and losing for male wheelchair tennis players in Paralympics Games Motriz. 2017c;23(3):e101663.
50. Sánchez-Pay A, Sanz-Rivas D. Competitive evolution of professional wheelchair tennis from the Paralympic Games in Athens 2004 to Rio 2016: an observational study. *Int J Environ Res Public Health*. 2021a;18(6).
51. Sánchez-Pay A, Pino-Ortega J, Sanz-Rivas D. Influence of successive wheelchair tennis matches on handgrip strength in high-level male players. *Int J Environ Res Public Health*. 2023;20(6).
52. Sindall P, Lenton JP, Whytock K, Tolfrey K, Oyster ML, Cooper RA, et al. Criterion validity and accuracy of global positioning satellite and data logging devices for wheelchair tennis court movement. *J Spinal Cord Med*. 2013a;36(4):383-93.
53. Sindall P, Lenton J, Cooper R, Tolfrey K, Goosey-Tolfrey V. Data logger device applicability for wheelchair tennis court movement. *J Sports Sci*. 2015;33(5):527-33.
54. van der Slikke RMA, Berger MAM, Bregman DJJ, Veeger D. Wearable Wheelchair Mobility Performance Measurement in Basketball, Rugby, and Tennis: Lessons for Classification and Training. *Sensors (Basel)*. 2020;20(12).
55. Rhodes J, Mason B, Perrat B, Smith M, Goosey-Tolfrey V. The validity and reliability of a novel indoor player tracking system for use within wheelchair court sports. *J Sports Sci*. 2014;32(17):1639-47.
56. Sánchez-Pay A, Torres-Luque G, Sanz-Rivas D, Courel-Ibáñez J. The use of bounce in professional wheelchair tennis. *Int J Sports Sci Coach*. 2020;15(3):375-81.
57. Cavedon V, Zancanaro C, Milanese C. Kinematic analysis of the wheelchair tennis serve: Implications for classification. *Scand J Med Sci Sports*. 2014;24(5):e381-8.
58. Bonato M, Maggioni MA, Rossi C, Rampichini S, La Torre A, Merati G. Relationship between anthropometric or functional characteristics and maximal serve velocity in professional tennis players. *J Sports Med Phys Fitness*. 2015;55(10):1157-65.
59. Reid M, Elliott B, Alderson J. Shoulder joint kinetics of the elite wheelchair tennis serve. *Br J Sports Med*. 2007;41(11):739-44.
60. Caspall JJ, Seligsohn E, Dao PV, Sprigle S. Changes in inertia and effect on turning effort across different wheelchair configurations. *J Rehabil Res Dev*. 2013;50(10):1353-62.
61. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. *J Am Coll Cardiol*. 2001;37(1):153-6.
62. Theisen D. Cardiovascular determinants of exercise capacity in the Paralympic athlete with spinal cord injury. *Exp Physiol*. 2012;97(3):319-24.
63. Achten J, Jeukendrup AE. Heart rate monitoring: applications and limitations. *Sports Med*. 2003;33(7):517-38.
64. Hill M, Talbot C, Price M. Predicted maximal heart rate for upper body exercise testing. *Clin Physiol Funct Imaging*. 2016;36(2):155-8.
65. Navas D, Veiga S, Navarro E, Ramón-Llín J. Differences in kinematic and match-play demands between elite winning and losing wheelchair padel players. *PLoS ONE*. 2020;15(9):e0233475.

66. Stojanović E, Stojiljković N, Scanlan AT, Dalbo VJ, Berkelmans DM, Milanović Z. The activity demands and physiological responses encountered during basketball match-play: a systematic review. *Sports Med.* 2018;48(1):111-35.
67. Schneider C, Hanakam F, Wiewelhove T, Döweling A, Kellmann M, Meyer T, et al. Heart rate monitoring in team sports-a conceptual framework for contextualizing heart rate measures for training and recovery prescription. *Front Physiol.* 2018;9:639.
68. Halson SL. Monitoring training load to understand fatigue in athletes. *Sports Med.* 2014;44 Suppl 2(Suppl 2):S139-47.