

Title page

Title: Patterns of habitat selection and use by *Macaca mulatta tcheliensis* in Winter and early Spring in Temperate Forest, Jiyuan, China

Authors: Xie Dong-Ming¹ Lu Ji-Qi^{1*} Alfred M. Sichilima² Wang Bai-Shi¹

Address:

1. Institute of Biodiversity and Ecology, Zhengzhou University, Zhengzhou, 450001 China
2. Department of Zoology and Entomology, Mammal Research Institute, University of Pretoria, Pretoria 0002, South Africa.

Personal E-mail: Lu Ji-Qi (corresponding author): lujq@zzu.edu.cn

Running title: Habitat selection and use in winter

Abstract

The chosen habitat of any animal species comprises a range of environmental features that provide adequate resources for its continuous survival. Consequently, the criteria of habitat selection by animals, combines a wider spectrum of both environmental and extrinsic factors, with major prerequisites based on food resources, availability of shelter and suitable ethics for procreation. From this study, conducted in winter and early spring, at Mt. Wangwushan area, located on N 35°05'-35°15', E 112°12'-112°22', in Taihangshan Macaque National Nature Reserve (TMNNR), Jiyuan, Henan Province, we show by elaborative results that *Macaca mulatta tcheliensis* was specifically associated with the following habitat characteristics: 1) the average tree DBH (diameter at breast height) with over 15 cm; 2) the distance from human disturbance of less than 2000 m; 3) the distance to water with less than 1000 m; 4) the gradient of mountain slope of 15°~40°; 5) which was markedly tarrying with the altitude ranges from 1000 m to 1300 m; and 6) where the canopy coverage was less than 60%. In addition, during this study, rhesus macaque mostly inhabited the sunny slopes of mountains than the shady areas. Results of the first five principal components analysis (PCA) accounted for the total variance of 68.88%, while the other factors showed insignificant effects on habitat selection by rhesus macaque in the temperate forest. In conclusion, these new results increase our understanding on the living status, under the harshest condition, in winter and early spring of *Macaca mulatta tcheliensis*, the sub-species of rhesus macaques as may be linked to its habitat selection and utilization, in the temperate forest.

Keywords: Rhesus macaque; habitat selection; ecological factors; winter; temperate forest.

Introduction

Ecology provides the impetus to understanding factors and processes in determining the distribution and abundance of species in space and time (Brown et al. 1995, Greenwood et al. 1996). Habitat-selection behavior as the identification procedure of preferred habitats by an animal, is a fundamental topic in wildlife ecology, and best explains techniques on how mobile organisms adapt to changing conditions (Railsback and Harvey, 2002). The following four criteria, in the classification of habitat selection as proposed by Johnson (1980) include: 1) the microhabitat, 2) the components of the home range, 3) the home range itself, and 4) the geographical distribution or regional level. The home range is more-or-less a restricted area where an animal moves during its normal activities (Harris et al., 1990). The third level features the particular habitat's attributes in a home range, influencing the animal to select the area. Habitat attributes may vary depending on the composition, spatial arrangement proportions and the animal species, including both environmental and extrinsic factors. In the biogeographic or continental spheres, it is well established that extrinsic factors of energetic constraints and concepts of seasonality in locating food are major determinants on the abundance of animal distribution and life historic traits of populations (Lindstedt and Boyce, 1985; Jarvis et al., 1994; Ferguson, 2002; Wiegand et al., 2008; Wilbert et al., 2000; Sichilima et al., 2008). Therefore, the patterns of food availability in the habitat and the seasonal energetic expense in locating food by animals, may not only influence the levels of species' distribution in the habitat but also the quality and the criterion of selecting the home range, suitable for the population (Sichilima et al., 2008).

In the previous decades, ecologists devoted more efforts and concerns to understanding the forces behind both the typical habitat selection and the related implications associated with predator avoidance. It has been observed that some prey species benefit from adopting multiple environmental habitats in order to decrease vulnerability from predators (Lima, 1998). For prey species therefore, predation pressure is one of the most important extrinsic factors influencing the spatial behavior of individuals and habitat selection (Bos and Carthew, 2003; Kotler, 1997; Lima and Dill, 1990).

Rhesus macaque which is listed as second class of key protected wildlife of China, is the only

non-human primate with a broader geographic distribution (Fooden, 2000; Southwick et al. 1996), and their populations are found ubiquitously throughout mainland Asia; from Afghanistan to India and Thailand to southern China (Rowe 1996; Smith and McDonough 2005). The *Macaca mulatta tcheliensis* is a subspecies of rhesus macaque endemic to China (Jiang et al., 1991; Zhang et al., 1991), and the majority of populations of this subspecies were confined mainly to the southern Mt. Taihangshan area (N 34°54'~35°16', E 112°02'~112°52') which is near the boundary of Henan and Shanxi Provinces of China, and where a National Nature Reserve was established in 1998 (Lu et al., 2007). Climate in this region is hot, rainy, and humid in summers, while cold and dry in winter with the annual temperature ranging from -20°C (-4° F) to 40°C (104° F). Average annual precipitation was 641 mm (2.10 ft), with the heaviest rain recorded from June to August (Qu et al. 1993; Hu et al., 2004). This is similar to the climate of the Pakistan region where rhesus macaques are found, but the habitat in Pakistan is dominated by mixed evergreen and deciduous forests that are highly disturbed (Goldstein and Richard, 1989), and which are also in proximity with habitat conditions of the Japanese monkeys, *Macaca fuscata* (Wada and Ichiki, 1980; Nakagawa, 1989).

The long winter with poor food supply, in Mt. Taihangshan area, is recorded from November to March of every year and is predictably a crucial period for the survival of macaques in their life history (Song and Qu, 1996; Lu et al., 2007). Zoo-geographical habitat fragmentation and degradation by deforestation, together with the highway construction are major environmental factors impairing the macaque conservation (Lu et al., 2007). Unfortunately, the ecological knowledge in winter, on the spatial and foraging behaviors, habitat selection and utilization for this rarely studied sub-species is unknown.

The main aim of this research work was to provide information on habitat selection and use by this sub-species of rhesus macaque in the temperate forest during winter and early spring, and to get its possible conservation implications. Consequently, a wide range of environmental factors associated with this animal were investigated as follows; terrain, altitude of the area, slope of site, slope of face, gradient of the slope, the distance from water sources, human disturbance, size of canopy, concealment, and average tree DBH (diameter at breast height). The two major and specific questions to be addressed in this study were: 1) what are preferred habitats for macaque in

winter? and 2) what are major factors influencing habitat use?

1 Methods

1.1 Study Area

The field work was conducted in Tiantan Station of Mt. Wangwushan area, located at N 35°05'-35°15', E 112°12'-112°22', in Taihangshan Macaque National Nature Reserve (TMNRR) which is about 40 km northwest from Jiyuan, Henan, China. The Reserve is about 56,600 ha in area, bordering the Yellow River in the south, and reaching the boundary of Shanxi Province. Continental and marine wind currents, with a prevalence of a northwest wind in winter and southeast wind in summer, ultimately influenced the monsoon type of climate in this study area. The region has four distinct seasons i.e. Summer, Winter, Spring and Autumn, with the variation in sunlight whilst heat and precipitation mostly occurring simultaneously. The mean annual sunshine is 2367.7 hr, with a total solar radiation of 118.17 kcal / cm², while the mean temperature is 14.3°C in the plain areas, 14.9°C in the foothills along the Southern Mt. Taihangshan, 13.2~14.0°C in the lower hilly area, and 8.6~11.1°C in the mountainous area. The mean annual precipitation for this area is 695 mm, and is 70 mm higher than in the plains, though precipitation exhibits both temporal and spatial variation. There are 1698 species of vascular plants, belonging to 734 genera and 163 families in the Reserve and most of them belonged to seed plants (Song and Qu, 1996; Lü et al., 2002; Lu et al., 2007).

The aim of this research was to study the habitats of *Macaca mulatta tcheliensis* inhabiting within the Tiantan Station of Mt. Wangwushan area in TMNRR. In the study area, there were 2 group species of macaque found and were differentiated by the codes of WW-1 and WW-2, respectively. Thirty nine (39) individuals were identified in the group of WW-1, while only eight (8) were identified in the group of WW-2. Since fewer individuals within WW-2 and it was seldomly seen in studied area, we took WW-1 as focal troop. Our research protocol was permitted by the Administration of TMRRN, and had complied with animal care regulations and applicable national laws of China. In the course of field work and investigations, we exerted non invasive methods on the macaques.

1.2 Quadrat Establishment and Survey

Field studies in the sites were conducted from November 2007 to February 2008, and from November 2008 to February 2009. Habitats used by macaque were located based on the activity evidence of its foraging, calling, playing and struggling etc. Using a portable GPS, every located site was geographically identified and parameters of latitude, longitude and altitude were recorded. A sampling quadrat of 10 m × 10 m (100 m²) was subsequently marked to establish the area of a particular site. During the study period, total 296 quadrats (180 of 2007-2008 and 116 of 2008-2009, respectively) were identified and data collection was carried out in all sites. A similarly marked site, located 50 m away from the rest of study area was established for the purpose of a control quadrat (Young et al., 1991). In our study area, the early spring only accounted for few weeks after winter before the abundance of food resources were recorded and most other weather conditions in the habitat were still very similar to those of winter season. To this effect therefore, there was no major seasonal fluctuations worth the separate analyses in the habitats.

Though many ecological factors would exert influences on animal habitat selection and use (Bond et al., 2002; Bos and Carthew, 2003), we mainly concerned those important factors for winter habitat use (Homan et al., 2000). In each located quadrat, necessary parametric measurements were calculated and recorded from the following environmental factors; topography, altitude of the site, slope position, slope direction, slope gradient, the distance from water sources, human disturbance, size of canopy, concealment, and the average DBH (diameter at breast height). The definition and categories of these ecological factors are as follows:

Topography: This referred to local topographical characteristics of the habitat, and was classified into hillside, ridge and valley of mountain.

Altitude: This referred to the a.s.l. height of the site where the activity traces were detected, and was categorised into three levels: ≤ 1000 m, 1000~1300 m, and ≥ 1300 m.

Slope position: This referred to a level of slanting of an angle on the surface of the mountain, to the horizontal surface, with different positions including upper, middle, and lower positions,

Slope-direction: This referred to which direction and level was the slope facing i.e. against or towards the sun, including; sunny slope (S 67.5° E~S22.5° W), half sunny slope (N 22.5° E~S 67.5° E and S 22.5° W~N 67.5° W) and shady slope (N 67.5° W~N 22.5° E).

Slope-gradient: This referred to the extent of calculated slope in degrees, and was determined in three levels: gentle slope ($\leq 15^\circ$); slight slope ($15^\circ \sim 40^\circ$); steep slope ($\geq 40^\circ$).

Distance from water source: This factor was classified into two levels, near (≤ 1000 m), and far (> 1000 m), based on the distance from the center of quadrat to water source.

Human disturbance: This factor reflects the extent of human disturbance including roads, buildings, other human activities, etc. and was divided into three levels: ≤ 1000 m; $1000 \sim 2000$ m; ≥ 2000 m.

Canopy: This referred to the coverage degree of tree crown, and was defined in three categories, $\leq 30\%$, $30\% \sim 60\%$, and $\geq 60\%$.

Concealment: This referred to the extent at which the selected habitat, sheltering macaques was hidden from the clear visibility. This factor was determined by the horizontal front view of the selected position within a visible distance. Concealments were expressed at three specific levels, with high (≥ 50 m), moderate (20-50 m), and lower concealment (≤ 20 m).

Average DBH: The DBH was referred to the "diameter at breast height", with the collected data of average DBH categorized at ≤ 15 cm, ≥ 30 cm, and $15 \sim 30$ cm.

The Vanderploeg and Scavia's index (E_i) (Lechowicz 1982) was used to identify habitat use by macaque in study area:

$$E_i = [W_i - (1/n)] / [W_i + (1/n)]$$

Where W_i is the selectivity coefficient (Chesson 1978), and n is the number of quadrant. The index W_i was calculated using the following formula:

$$W_i = (r_i / p_i) / \sum (r_i / p_i)$$

Where p_i is the proportion of samplings in the i th level, and r_i is the proportion of habitats in the i th class. E_i is scaled between -1 and +1. When $E_i = -1$, macaque did not select this habitat (N); $-1 < E_i < 0$, macaque avoided this habitat (NP), when $E_i = 0$, macaque selected and used this habitat randomly (R), and $0 < E_i < 1$ indicates preferred this habitat (P), $E_i = 1$, macaque specially preferred the habitat (SP).

1.3 Data Analysis

SPSS for windows (version 13.0) was adopted in data analysis. In order to determine how the

effects of different factors influenced the macaque's habitat selection, the principal component analysis (PCA) was employed. Under the condition of eigenvalue > 1 , the principal component was considered a crucial determinant in the habitat use by macaques.

2 Results

2.1 Preferred habitat of macaques

From this study, results showed that macaques usually chose habitats where the following characteristics were evident: 1) the average tree DBH was over 15 cm; 2) the distance from human disturbance was not less than 2000 m; 3) the distance to the water source less than 1000 m; 4) the slope gradient was $15^{\circ}\sim 40^{\circ}$; 5) with the altitude ranging from 1000 m to 1300 m; and 6) the canopy cover less than 60%. Furthermore, macaques that occurred at Mt. Wangwushan area preferred sunny slopes in winter and early spring (Table 1).

[Insert Table 1 here]

2.2 Factors affecting habitat use of macaques

Results from the principal component analysis (PCA) revealed that 1 - 5 components, accounted for 68.88% on the total variance (Table 2). The first principal axis reflected effects derived from altitude and slope-direction, which represented the geographical traits of the selected habitat by macaques and was thus named as a geographical factor. The distance from the water source contributed more to the second principal axis and was regarded as water factor. A combination of slope gradient, concealment, and the average tree DBH, constituted the third and single factor known as foraging on the principal axis while the human disturbance factor, and the resting factor, were fourth and fifth principal components, respectively (Table 3).

[Insert Table 2 here]

[Insert Table 3 here]

3 Discussion

Rhesus macaques are regarded as habitat generalists, and their adaptive ability to marginal or degraded ecological conditions has been a key factor in allowing the species to survive in fragmented landscapes and highly populated areas (Lu et al., 2007; Song and Qu, 1996; Wada and Ichiki, 1980). In temperate regions therefore, cold and snow is prevalent in winter and this

adversely limits foraging and other activities to a greater extent. This harsh period is thus regarded as a pivotal stage for mammals in their life history and compels them to adapt to new habitat use for survival (Homan et al., 2000). The habitat selection by this animal also takes into consideration the predation risks, abundance and quality of food resources which changed seasonally (Rueda et al., 2008).

Across all habitat types, the macaques sub-species on which this study was conducted, in winter and early spring, invested much of its time on the activities of foraging and resting, in addition to other characteristic behaviors of grooming, playing, moving around etc. (Seth & Seth 1986). A variety of biological and abiotic factors might have influenced the geographical distribution pattern of animals but food, water, and concealment were the three major components influencing the habitat selection, while the food resource factor was ranked highly essential (Ma et al., 2004). Mammals endemic to the temperate area easily adapted to the severe conditions during winter period and subsequently exhibit various adaptive patterns during habitat use and selection (Wilbert et al., 2000; Young et al., 1991). Similarly, during this field study, the activity traces of macaque were often found in ridges, with gentle and sunny slopes, suggesting two energy saving points that, in ridges, the animals easily moved from one place to another in order to physically reduce the energetic costs and also gained more energy for body temperature maintenance directly from the sun.

The *Macaca mulatta tcheliensis* is a phytophagous animal, feeding on plants and leaves of some trees, which are abundantly available during the summer period (Wada 1984; Ren et al., 2001; Hu et al., 2004). In winter, however, these food resources become scarce and the animal exhibits adaptive characteristic patterns of phytodiversity and increases the probability of finding alternative food items including scattered seeds such as acorns of Cork oak (*Quercus variabilis*), seeds of Oriental Bittersweet (*Celastrus orbiculatus*), fruits of Dteplum Persimmon (*Diospyros lotus*), tubers of some plants, barks of some trees and so on, which are usually found in the valleys or foot-hills (Strier, 2000; Hanya, 2004; Guo et al., 2007, 2008). In this research, we also found that macaques preferred the habitats with sunny slopes and canopy cover of less than 60%, intimating that incumbent tree species in these areas possibly had produced fruits, seeds, nuts, and

corns, which might have dropped and are still available on the ground during the winter period for macaques to eat. It is also suggested that the reason for a choice of habitats closer to human activities in winter by macaques might be probably targeted on food crop remains like; *Zea mays*, *Daucus carota*, *Brassica campestris* together with root crops like cassava and sweetpotato which farmers grow. Nevertheless, from a security perspective, the macaques would still select those taller trees, with stronger branches far away from humans as night-sleep sites (Guo et al, 2010).

Water was also another important factor affecting habitat use and selection by macaques in winter. The mean annual precipitation was 690 mm in study area, and was insufficient in winter (Song and Qu, 1996; Hu et al., 2004). Our result showed that macaques were commonly seen searching and wandering in the valley and foot-hills, because there were some temporary water resources in such sites (Song and Qu, 1996), and for several times, we observed macaque individuals drinking water *in situ* in the valley. It was, however, reported on two contrasts that *Rhinopithecus roxellana* and *R. bieti* always take some little quantities of water (Li et al, 2006; Su et al, 1998), while it is evident that despite the Japanese macaque (*M. fuscata*) may be found very close to the hot-springs, they preferred gaining heat to drinking hot water (Suzuki, 1965; Wada, 1964).

Because the macaques are found in such a broad geographic area, it is difficult to specifically summarize their habitats precisely. Depending on the type of habitat the animals may be found, they are quadrupedal, predominantly arboreal or terrestrial (Seth et al. 2001). The result from this study indicated that in the northernmost part of Mt. Taihangshan area in China, rhesus macaques inhabit a secondary deciduous forest between 300 and 1200 m a.s.l. (984 and 3937 ft) in a temperate climate with cold, snowy winters similar to the climate of the central, Midwestern United States (Qu et al. 1993; Lu et al., 2007). To make the macaque conservation compatible with our current policies and the local economic development, there is need for continuous protection of similar characteristic habitats as mentioned above. There is need therefore, for more ecological work to be carried out in the macaques foraging behavior, in temperate habitats in order to further our understand in the varying adaptation characteristics of the animal

From the current study, we got a primary understanding on ecological factors affecting

habitat selection and use in winter of rhesus macaques in its northernmost distributed area. It was surprisingly that the result from PCA explained 68.880% of the variance of examined factors, which is partly because the examined factors had no close relationship between each other. This result also indicated that many other ecological factors possibly exert role on habitat selection and use of rhesus macaques. We should pay more attention on analyzing influences of the other ecological factors in the future researches.

Acknowledgments We thank Dr. LÜ Jiu-Quan, Mr. WANG Hao-Feng, TANG Fa-You, KONG Mao-Cai, HOU Jia-Fu, Guo Xiang-Bao, and QU Lei for their assistance in the field investigation. This research was financially supported by National Natural Science Foundation of China (No. 30770381 and 30970378).

References

- Bond BT, Burger LW Jr, Leopold BD, Jones JC, Godwin KD, 2002. Habitat use by cottontail rabbits across multiple spatial scales in Mississippi. *Journal of Wildlife Management*, 66: 1171-1178
- Bos DG, Carthew SM, 2003. The influence of behaviour and season on habitat selection by a small mammal. *Ecography*, 26: 810-820
- Brown JH, Mehlman DW, Stevens GC, 1995. Spatial variation in abundance. *Ecology*, 76: 2028-2043
- Chesson J. 1978: Measuring preference in selective predation. *Ecology*, 59: 211-215
- Ferguson SH. 2002. The effects of productivity and seasonality on life history: comparing age at maturity among moose (*Alces alces*) populations. *Global Ecology and Biogeography*, 11: 303-312
- Fooden J. 2000. Systematic review of the rhesus macaque, *Macaca mulatta* (Zimmermann, 1780). *Fieldiana Zoology*, 96: 1-180
- Goldstein SJ, Richard AF. 1989. Ecology of rhesus macaques (*Macaca mulatta*) in northwest Pakistan. *International Journal of Primatology*, 10(6): 531-567
- Greenwood JJD, Gregory RD, Harris S, Morris PA, Yalden DW. 1996. Relations between

- abundance, body-size and species number in British birds and mammals. *Philosophical Transactions of the Royal Society of London B*, 351: 265-278
- Guo ST, Ji WH, Li BG, Li M. 2008. Response of a group of Sichuan snub-nosed monkeys to commercial logging in the Qinling Mountains, China. *Conservation Biology*, 22(4): 1055-1064
- Guo ST, Li BG, Watanabe K. 2007. Diet and activity budget of *Rhinopithecus roxellana* in the Qinling Mountains, China. *Primates*, 48(4): 268-276
- Guo XB, Wang ZL, Tian JD, Lu JQ, Liu JD. 2010. Sleeping site selection by rhesus macaques (*Macaca mulatta tcheliensis*) in Taihangshan National Nature Reserve, Henan, China. *Sichuan Journal of Zoology*, 29 (6): 849-856
- Hanya G. 2004. Diet of a Japanese macaque troop in the coniferous forest of Yakushima. *International Journal of Primatology*, 25(1): 55-71
- Harris SW, Cresswell J, Forde PG, Trehela WJ, Woollard T, Wray S. 1990. Home-range analysis using radio-tracking data: A review of problems and techniques particularly as applied to mammals. *Mammal Review*, 20: 97-123
- Homan HJ, Linz GM, Bleier WJ. 2000. Winter habitat use and survival of female ring-necked pheasant (*Phasianus colchicus*) in southeastern north Dakota. *The American Midland Naturalist*, 143(2): 463-480
- Hu YM, Zhang YL, Wei FY, Wang SX. 2004. The analysis of climatic resources in Taihangshan Mountains National Nature Reserve, Jiyuan, China. *Henan Meteorology*, 4: 27
- Jarvis JUM, Oriain MJ, Bennett NC, Sherman PW. 1994. Mammalian eusociality- a family affair. *Trends in Ecology and Evolution*, 9: 47-51
- Jiang XL, Wang YX, Ma SL. 1991. Taxonomic and distribution of subspecies of rhesus monkey (*Macaca mulatta*) in China. *Zoological Research*, 12: 241-246
- Johnson DH. 1980. The composition of usage and availability measurements for evaluating resource preference. *Ecology*, 61: 65-71
- Kotler BP. 1997. Patch use by gerbils in a risky environment: manipulating food and safety to test four models. *Oikos*, 78: 274-282
- Lechowicz MJ. 1982. The sampling characteristics of selectivity indices. *Ecology*, 52 (1): 22-30

- Li DY, Peng ZS, Ren BP, Grüter CC, Zhou QH, Wei FW. 2006. Early autumn habitat selection by the Yunnan snub-nosed monkey (*Rhinopithecus bieti*) in Tacheng, Yunnan. *Journal of China West Normal University (Natural Sciences)*, 27(3): 233-238
- Lima SL, Dill LM. 1990. Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology*, 68: 619-640
- Lima SL. 1998. Stress and decision making under the risk of predation: recent developments from behavioral, reproductive and ecological perspectives. *Advances in the Study of Behavior*, 27: 215-290
- Lindstedt SL, Boyce MS. 1985. Seasonality, fasting endurance, and body size in mammals. *American Naturalist* 125: 873-878
- Lu JQ, Hou JH, Wang HF, Qu WY. 2007. Current status of *Macaca mulatta* in Taihangshan mountain areas, Jiyuan, Henan, China. *International Journal of Primatology*, 28 (5): 1085-1091
- Lü JQ, Lu JQ, Li JY, Zhang JY, Qu WY. 2002. Feeding habits of *Macaca mulatta* in Taihang Mountains. *Chinese Journal of Ecology*, 21: 29-31
- Ma JZ, Zou HF, Jia JB. 2004. *Wildlife Management*. Harbin: Northeast Forestry University Press, 43-62
- Nakagawa N. 1989. Bioenergetics of Japanese monkeys (*Macaca fuscata*) on Kinkazan island during winter. *Primates*, 30(4): 441-460
- Qu WY, Zhang RZ, David M, Southwick CH. 1993. Rhesus monkeys (*Macaca mulatta*) in the Taihang Mountains, Jiyuan county, Henan, China. *International Journal of Primatology*, 14, 607-621
- Railsback SF, Harvey BC. 2002. Analysis of habitat-selection rules using an individual-based model. *Ecology*, 83(7): 1817-1830
- Ren BP, Zhang SY, Wang LX, Liang B, Li BG. 2001. Vertical distribution of different age-sex classes in a foraging group of Sichuan golden monkeys (*Rhinopithecus roxellana*). *Folia Primatologica*, 72: 96-99
- Rowe N. 1996. *The pictorial guide to the living primates*. East Hampton, New York: Pogonias Press, 263
- Rueda M, Rebollo S, Gálvez-Bravo L, Escudero A. 2008. Habitat use by large and small

- herbivores in a fluctuating Mediterranean ecosystem: Implications of seasonal changes. *Journal of Arid Environments*, 72 (9): 1698-1708
- Seth PK, Chopra PK, Seth S. 2001. Indian rhesus macaque: habitat, ecology and activity patterns of naturally occurring populations. *Envis Bulletin: Wildlife and Protected Areas*, 1(1): 68-80
- Seth PK, Seth S. 1986. Ecology and behavior of rhesus monkeys in India. In: *Primate Ecology and Conservation*. Else JG, Lee PC (eds.), Cambridge University Press, Cambridge. 2, 192-305
- Sichlimma AM, Bennett NC, Faulkes CG, Le Comber SC. 2008. Evolution of African mole-rat sociality: burrow architecture, rainfall and foraging in colonies of the cooperatively breeding *Fukomys mechowii*. *Journal of Zoology*, 275: 276-282
- Smith DG, McDonough J. 2005. Mitochondrial DNA variation in Chinese and Indian rhesus macaques (*Macaca mulatta*). *American Journal of Primatology*, 65(1): 1-25
- Song CS, Qu WY. 1996. *Scientific Survey of Taihangshan Macaque Nature Reserve*. Beijing: China Forestry Publishing House, 56-105
- Southwick CH, Zhang YZ, Jiang HS, Liu ZH, Qu WY. 1996. Population ecology of rhesus macaques in tropical and temperate habitats in China. In: Fa JE, Lindburg DG (eds.) *Evolution and Ecology of Macaque Societies*. Cambridge University Press, New York, 95-105
- Strier KB. 2000. Population variability and regional conservation priorities for muriquis (*Brachyteles arachnoides*) in Brazil's Atlantic Forest. *Biotropica*, 32 (4b): 903-913
- Su YJ, Ren RM, Yan KH, Li JJ, Zhou ZQ, Hu ZL, Hu YF. 1998. Preliminary survey of the home range and ranging behavior of golden monkeys (*Rhinopithecus roxellena*) in Shennongjia National Natural Reserve, Hubei, China. In: Jablonski NG (ed.) *The Natural History of the Doucs and Snub nosed Monkeys*. Singapore: World Scientific Publishing, 255-268
- Suzuki A. 1965. An ecological study of wild Japanese monkeys in snowy areas focused on their food habits. *Primates*, 6(1): 31-72
- Wada K, Ichiki Y. 1980. Seasonal home range use by Japanese monkeys in the snowy Shiga heights. *Primates*, 21(4) : 468-483
- Wada K. 1964. Some observations on the life of monkeys in a snowy district of Japan. *Physiology and Ecology*, 12: 151-174

- Wada K. 1983. Ecological adaptation in rhesus monkeys at the Kumaon Himalaya. *The Journal of the Bombay Natural History Society*, 80(3): 469-498
- Wiegand T, Naves J, Garbulsky M, Fernández N. 2008. Animal habitat quality and ecosystem functioning: Exploring seasonal patterns using NDVI. *Ecological Monographs*, 78(1) : 87-103
- Wilbert CJ, Buskirk SW, Gerow KG. 2000. Effects of weather and snow on habitat selection by American martens (*Martes americana*). *Canadian Journal of Zoology*, 78(10) : 1691-1696
- Young L, Zhang GM, Zhang ZW. 1991. Winter movements and habitat use by Cabot's Tragopans *Tragopan caboti* in southeastern China. *Ibis*, 133(2): 121-126
- Zhang YZ, Quan GQ, Zhao TG, Southwick CH. 1991. Distribution of macaques (*Macaca*) in China. *Acta Theriologica Sinica*, 11: 171-185

Table and figure legends

Table 1 Habitat selection by rhesus macaque at Mt. Wangwushan area in winter and early spring

Table 2 The eigenvalue of habitat selection by rhesus macaque at Mt. Wangwushan area in winter and early spring

Table 3 The rotated matrix of the eigenvector of habitat selection by rhesus macaque in Mt. Wangwushan area in winter and early spring

Table 1 Habitat selection by rhesus macaque at Mt. Wangwushan area in winter and early spring

Item	i	p_i	W_i	E_i	Habitat selection	Item	i	p_i	W_i	E_i	Habitat selection
Topography	Hillside	194	0.66	0.98	P	Distance from water source	≤1000m	285	0.96	0.99	SP
	Ridge	28	0.09	0.45	P		>1000m	11	0.04	-0.42	NP
	Gully	74	0.25	0.90	P						
Altitude	≤1000m	26	0.09	0.39	P	Human disturbance	≤1000m	227	0.77	0.99	SP
	1000-1300m	237	0.80	0.99	SP		1000-2000m	66	0.22	0.87	P
	≥1300m	33	0.11	0.57	P		≥2000m	3	0.01	-0.94	NP
Slope position	Upper	84	0.28	0.92	P	Canopy (%)	≤30%	188	0.64	0.98	P
	Middle	104	0.35	0.95	P		30%-60%	105	0.35	0.95	P
	Lower	108	0.37	0.95	P		≥60%	3	0.01	-0.94	NP
Slope direction	Sunny slope	209	0.71	0.99	SP		≥50m	174	0.59	0.98	P
	Half sunny slope	77	0.26	0.90	P	Concealment	20-50m	103	0.35	0.95	P
	Shady slope	10	0.03	-0.49	NP		≤20m	19	0.06	0.02	R
Slope gradient	≤15°	21	0.07	0.03	R		≤15cm	9	0.03	-0.57	NP
	15°-40°	195	0.66	0.98	P	Average DBH	15-30cm	235	0.79	0.99	SP
	≥40°	80	0.27	0.91	P		≥30cm	52	0.18	0.80	P

Note: i -Eigenvalue; p_i - Quadrat number with i eigenvalue; W_i -Selection coefficient; E_i -Selection index; P-Preference; SP-special preference; N-No Selection; R-Random selection; NP-Non Preference.

Table 2 The eigenvalue of habitat selection by rhesus macaque at Mt. Wangwushan area in winter and early spring

Main component No.	Eigenvalue	Contribution rate (%)	Accumulative contribution rate (%)
1	1.930	19.300	19.300
2	1.475	14.751	34.051
3	1.265	12.651	46.702
4	1.216	12.160	58.862
5	1.002	10.018	68.880
6	0.856	8.562	77.442
7	0.727	7.270	84.712
8	0.666	6.655	91.367
9	0.521	5.210	96.577
10	0.342	3.423	100.000

Note: Number 1 to 10 refers to the first up to the tenth principal components.

Table 3 The rotated matrix of the eigenvector of habitat selection by rhesus macaque in Mt. Wangwushan area in winter and early spring

Variables	Principal components				
	1	2	3	4	5
Topography	0.433	-0.600	-0.023	0.419	0.128
Altitude	-0.726	-0.018	0.301	0.244	-0.057
Slope site	0.421	0.326	-0.063	-0.472	0.182
Slope face	0.710	-0.471	-0.065	-0.255	0.059
Slope gradient	0.232	0.384	0.639	-0.218	0.147
Distance from water resource	0.355	0.600	-0.341	0.180	-0.120
Human disturbance	0.312	0.348	-0.278	0.671	0.009
Canopy	0.281	0.356	0.411	0.233	0.076
Concealment	0.377	-0.173	0.627	0.300	-0.129
Average tree DBH	-0.192	0.008	-0.050	0.138	0.942