

Cryphonectriaceae: Biodiverse and Threatening Tree Pathogens in the Tropics and Southern Hemisphere

Michael J. Wingfield, Nam Q. Pham,
Seonju Marinowitz, and Brenda D. Wingfield

Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural
Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa;
email: mike.wingfield@fabi.up.ac.za

**ANNUAL
REVIEWS CONNECT**

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Phytopathol. 2025. 63:553–75

First published as a Review in Advance on
May 28, 2025

The *Annual Review of Phytopathology* is online at
phyto.annualreviews.org

<https://doi.org/10.1146/annurev-phyto-121823-030316>

Copyright © 2025 by the author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information.



Keywords

Cryphonectriaceae, Eucalyptus canker, invasive fungi, forest pathogens, fungal diversity, host shift

Abstract

The chestnut blight pathogen *Cryphonectria parasitica* is well-known for the devastation it caused to North American forests. It is less well recognized that numerous other fungi in the Cryphonectriaceae are emerging as threats to native and planted forests in the tropics and Southern Hemisphere. Unlike *C. parasitica*, these fungi, such as *Chrysosporium cubensis*, initially gained attention due to a canker disease in plantations of non-native *Eucalyptus*. More than four decades of research have revealed a wide diversity of Cryphonectriaceae species that infect many other tree genera in the Myrtales. These fungi often exist as endophytes but become problematic when trees are planted outside their native range. Growing numbers of species are also undergoing host shifts from native to susceptible trees such as *Eucalyptus*, posing serious risks to both natural and planted forests. These fungi provide an example of the biodiversity of tree-infecting fungi that is understudied, despite their significant potential to harm forest ecosystems.

Stem cankers:

localized bark infection by pathogens, causing sunken or swollen areas, girdling, and death of nearby plant tissues

Eucalyptus plantation

industry: commercial plantations of Australian *Eucalyptus* species expanded globally for fiber production, reaching 20 million hectares by 2008

Myrtales: large plant order with nine families, notably Myrtaceae; many species host Cryphonectriaceae fungi, indicating many undiscovered, potentially pathogenic species

INTRODUCTION

The Cryphonectriaceae owes its notoriety to the destructive tree pathogen *Cryphonectria parasitica* (Murrill) M.E. Barr that devastated North American chestnut [*Castanea dentata* (Marsh.) Borkh.] forests beginning in the early 1900s (3). The pathogen was first discovered on infected trees in New York in 1904, although it might have existed there as early as 1893. The disease spread rapidly throughout the eastern seaboard of the United States, virtually eliminating a dominant component of the natural forests in the area (3, 68, 86, 88). By the late 1920s, most *C. dentata* trees in the native range were dying or had died, with only coppice stems remaining (51).

The chestnut blight pathogen was first described as *Diaporthe parasitica*, residing in the Diaporthales (88). Various taxonomic treatments followed (4, 29, 108), with *Cryphonectria parasitica* most recently applied to the pathogen (7). The family Cryphonectriaceae was established by Gryzenhout et al. (59) to accommodate this fungus and numerous other related species that had been described (52, 53, 56, 58, 144). All these species are pathogens infecting the cambial tissues of trees and resulting in stem cankers. By 2009, twenty-two species had been described in the Cryphonectriaceae, and these were treated in the monograph of Gryzenhout et al. (63). They were also mostly known from the Northern Hemisphere, and only *C. parasitica*, which by that time was also causing serious disease problems on *Castanea sativa* Mill. in Europe (67), was considered a serious threat to natural forests.


The discovery of a serious stem canker disease on *Eucalyptus* in Cuba added a fundamentally novel view of the Cryphonectriaceae as tree pathogens. This disease was first encountered and described by Bruner (13) as *Diaporthe cubensis* Bruner. As the taxonomy of the Diaporthales was revised over time (7), the fungus was recognized as a species typical of *Cryphonectria* and was transferred to that genus by Hodges (72) as *Cryphonectria cubensis* (Bruner) Hodges.

Prior to the description of *C. cubensis* from Cuba, *Cryphonectria* species were known only from temperate regions of the world and occurred on species in the Fagaceae (45, 88). This discovery also gave rise to a period when *C. cubensis* was recognized as a pathogen threatening the rapidly emerging *Eucalyptus* plantation industry in tropical regions of Brazil. This industry, also being established in various other parts of the Southern Hemisphere, was based on the rapid growth of *Eucalyptus* species, mostly native to Australia, where the genus is remarkably biodiverse. These trees include species with a wide range of climatic and edaphic characteristics as well as properties of commercial relevance such as the production of solid wood, pulp products, and charcoal. Consequently, when *C. cubensis* was first discovered as an important stem canker pathogen in Brazil, the fungus gained substantial new relevance (69–71).

In this review, we seek to emphasize the relevance of species in the Cryphonectriaceae occurring in the tropics and Southern Hemisphere. Owing to their particular importance on *Eucalyptus* species, the focus is mostly on species that occur on trees in the Myrtales, the plant order that accommodates *Eucalyptus* and its relatives. A point of emphasis is on the diversity of these fungi in the Southern Hemisphere, as new species are increasingly being discovered. Our aim is especially to emphasize the importance of fungal diversity in natural woody ecosystems, the fact that these fungi are unacceptably poorly studied, and that many also threaten the future health of forest trees globally.

CRYPHONECTRIA ON EUCALYPTUS: PERSPECTIVES FROM PLANTATION FORESTRY

As pathogens, the Cryphonectriaceae are best known as wound-infecting fungi that colonize cambial tissues, causing diffuse cankers that can rapidly girdle infected tree branches and stems (Figure 1). Asexual structures are flask-shaped pycnidia from which hyaline conidia exude in



(Caption for **Figure 1** appears on following page)

Disease symptoms associated with Cryphonectriaceae on Myrtaceae. (a–f) Symptoms on infected trees. (g–j) Characteristics of sexual structures. (k–m) Asexual structures.

typically yellow to orange masses. The sexual structures are perithecia, usually with long necks containing operculate asci. The ascospores are usually, but not always, two-celled, divided by a medium septum. As increasing numbers of species have been discovered, substantial diversity in these morphological characteristics has also emerged (Figure 1).

Hodges et al. (70) made the first formal discovery of Cryphonectria canker on *Eucalyptus* in Brazil, leading to the pathogen being identified as *Diaporthe cubensis*. This discovery set in motion numerous studies of the disease in Brazil, seeking to manage a problem that was potentially threatening to an emerging and important industry where significant pulp mills depended on timber from these trees (1, 42–44, 69). Various taxonomic studies sought to clarify the identification of the fungus where the generic names *Endothia*, *Diaporthe*, and *Cryphonectria* had been applied and remained confusing. Hodges (72) considered this problem in detail and determined that the fungus on *Eucalyptus* was best accommodated in *Cryphonectria*. The name *Endothia* applied to collections on *Syzygium aromaticum* (syn. *Eugenia aromatica*; also Myrtaceae) (99, 109) as *Endothia eugeniae* (Nutman & F.M. Roberts) J. Reid & C. Booth was resolved by Hodges et al. (73), who showed that the two fungi were the same and reduced *E. eugeniae* to synonymy with *C. cubensis*. The conspecificity was supported by Micales & Stipes (87) based on chemotaxonomic analyses and later confirmed based on DNA data (93).

The discovery of Cryphonectria canker on *Eucalyptus* gave rise to a heightened interest in this fungus. This is mainly due to the potential damage that it might cause to plantings of these trees. Surveys in plantations in various tropical areas of the world were undertaken, resulting in numerous new reports of *C. cubensis* [Cameroon (47), Florida, Hawaii, Puerto Rico (74), and India (117)]. Reports also included those on species of *Syzygium*, including the clove tree (*S. aromaticum*) in Brazil (73), Indonesia (73), and Tanzania (99). The severity of disease on clove was, however, much lower in Brazil and Indonesia (73). This led to the intriguing view of Hodges et al. (73) that the fungus might have had its origin on clove and been introduced into Brazil via the spice trade. By 1990, *C. cubensis* had been recorded on species of *Eucalyptus* and *Syzygium* (as *Eugenia*) in Brazil, Cameroon, Costa Rica, Cuba, India, Indonesia, Malaysia, Surinam, Puerto Rico, the United States (Florida, Hawaii), and Western Samoa (12, 71). An intriguing report of the fungus from the roots of naturally occurring *Eucalyptus marginata* in Western Australia by Davison & Coates (39) added some confusion regarding the possible origin of the fungus occurring outside the native range of these trees.

Wingfield et al. (157) published a first report of *C. cubensis* from South Africa, after stem cankers caused by the fungus were found on clonally propagated *Eucalyptus* in the subtropical coastal areas of the country in 1986. This was a time when *Eucalyptus* plantation industry was expanding rapidly in South Africa to provide fiber for multiple newly established pulp mills. The discovery of *C. cubensis* in that country was of particular concern because of the problems that were being reported from Brazil (30). Consequently, intensive studies were launched to better understand the pathogen as well as to establish management strategies to reduce its potential negative impact (129, 139–143). Taxonomic studies included not only isolates from South Africa but also those available from other parts of the world, including South American and Southeast Asian countries (91, 92, 134, 141).

An important breakthrough in understanding the possible origin of *C. cubensis* emerged when the fungus was discovered for the first time on a tree that did not reside in the Myrtaceae, thus not a close relative of *Eucalyptus*. In this case, the fungus was found in Colombia on native *Pleroma*

Clove tree: *Syzygium aromaticum*, native to Indonesia, is cultivated globally for aromatic flower buds (cloves) used as a spice since the 1700s




Figure 2

Historical timeline outlining the evolution of Cryphonectriaceae taxonomy. Waffle chart indicates the accumulation of formally recognized species ascribed to Cryphonectriaceae over time. Each square represents a single taxon, and different colors represent the plant order from which each species was first described.

(syn. *Tibouchina*) (156), a genus of trees and shrubs residing in the Melastomataceae (Myrtales) (31). Pathogenicity tests showed that isolates from *Pleroma* could infect both these trees as well as *Eucalyptus* (156). This discovery provided a basis for the hypothesis that *C. cubensis* on *Eucalyptus* in South America could have arisen from a host shift (121) from native Melastomataceae to infect *Eucalyptus*. It was also recognized at that time that infections on *Eucalyptus* in South America resulted in somewhat different symptoms from those found in South Africa (154). These observations began to hint at the probability that *C. cubensis* in different parts of the world might not be the same fungus. Until that time, all descriptions of this fungus and its close relatives had relied on the morphology of their fruiting structures.

DEEP IMPACT OF PHYLOGENETIC INFERENCE BASED ON DNA SEQUENCE DATA

As with most fungi, the application of DNA sequence data and phylogenetic inference had a fundamental impact on the taxonomy of these fungi, with many new genera and species being discovered (Figure 2). The first DNA data for the Cryphonectriaceae in the tropics and Southern Hemisphere were based only on ITS sequences (92). This study provided additional evidence that *C. cubensis* in South America and other parts of the world represented at least two different species and that these Southern Hemisphere fungi were distinct from *C. parasitica*.

Subsequent to the study of Myburg et al. (92), DNA sequence data, together with morphological characteristics, were applied to new collections of *C. cubensis* isolates from *Eucalyptus* and *Pleroma* as well as from the South African indigenous tree *Syzygium cordatum* (Myrtaceae), on which the fungus had been discovered (64, 90, 156). The occurrence of the fungus on a native tree supported the hypothesis that the pathogen known as *C. cubensis* in South Africa could have undergone a host shift from native to planted trees in the country. This and several other studies (91–93, 113) gave rise to the genus *Chrysosporthe* being established, with *Chrysosporthe cubensis*

Host shift: a situation in which a pathogen undergoes an evolutionary change to infect a new host

Table 1 Major Cryphonectriaceae species discussed in this article

Species	Previous names	Host range	Geographical distribution	Possible native range	Mode of sexual reproduction
<i>Cryphonectria parasitica</i>	<i>Diaporthe parasitica</i> <i>Endothia parasitica</i> <i>Endothia gyrosa</i> var. <i>parasitica</i> <i>Valsonectria parasitica</i>	Fagales (<i>Alnus</i> , <i>Carpinus</i> , <i>Carya</i> , <i>Castanea</i> , <i>Castanopsis</i> , <i>Fagus</i> , <i>Ostrya</i> , <i>Quercus</i>) Sapindales (<i>Acer</i> , <i>Rhus</i>) Magnoliales (<i>Liriodendron</i>) Rosales (<i>Malus</i>)	North America (USA, Canada) Europe Asia (China, India, Iran, Japan, Korea, Taiwan) Northern Africa (Tunisia) Oceania (Australia)	East Asia	Heterothallic
<i>Chrysosporthe austroafricana</i>	NA	Myrtales (<i>Corymbia</i> , <i>Eucalyptus</i> , <i>Eugenia</i> , <i>Dissotis</i> , <i>Pleroma</i> , <i>Psidium</i> , <i>Syzygium</i>)	Southern Africa (Eswatini, Malawi, Mozambique, Namibia, South Africa, Zambia)	Southern Africa	Heterothallic
<i>Chrysosporthe cubensis</i>	<i>Diaporthe cubensis</i> <i>Cryphonectria cubensis</i> <i>Cryptosporella eugeniae</i> <i>Endothia eugeniae</i> <i>Cryphonectria eugeniae</i>	Myrtales (<i>Cbaetogastra</i> , <i>Corymbia</i> , <i>Eucalyptus</i> , <i>Lagerstroemia</i> , <i>Miconia</i> , <i>Pleroma</i> , <i>Plinia</i> , <i>Psidium</i> , <i>Rhynchanthera</i> , <i>Syzygium</i>)	North America (USA: Florida) South America (Brazil, Colombia, Suriname, Venezuela) Caribbean (Cuba) Western and Central Africa (Ghana, Democratic Republic of the Congo, Cameroon)	South America	Homothallic
<i>Chrysosporthe deuterocubensis</i>	NA	Myrtales (<i>Eucalyptus</i> , <i>Eugenia</i> , <i>Melastoma</i> , <i>Metrosideros</i> , <i>Pleroma</i> , <i>Psidium</i> , <i>Syzygium</i>)	Southeast and East Asia (Indonesia, Thailand, Singapore, China, Taiwan, Malaysia, Vietnam) Pacific (USA: Hawaii) East Africa (Tanzania, Kenya, Mozambique, Malawi) Oceania (Australia)	Southeast Asia	Homothallic

Abbreviation: NA, not applicable.

(Bruner) Gryzenh. & M.J. Wingf. as the type species (56). The name *Chrysosporthe austroafricana* Gryzenh. & M.J. Wingf. was then applied to the South African fungus. Furthermore, Gryzenhout et al. (59) established the Family Cryphonectriaceae for these fungi and their phylogenetic relatives, including *C. parasitica* and other Northern Hemisphere species.

Studies on the Southern African *Chr. austroafricana* and South American *Chr. cubensis* isolates, including those from Southeast Asia, provided the first evidence that the latter collections likely represented a distinct species. van der Merwe et al. (133, 134, 137) considered this question in detail, including DNA sequence data for numerous gene regions as well as the application of microsatellite markers confirming this difference, and the name *Chrysosporthe deuterocubensis* Gryzenh. & M.J. Wingf. was erected for the Southeast Asian fungus. Interestingly, isolates from areas such as Tanzania, Kenya, Mozambique, Malawi, China, Hawaii, and Australia also resolved as *Chr. deuterocubensis* (21, 91, 93, 98, 104, 133, 149) (Table 1).

Following the discovery that *Chr. cubensis* represented more than a single species and fueled by a concern regarding these fungi as *Eucalyptus* pathogens, a period of intense collecting of the Cryphonectriaceae in the tropics and Southern Hemisphere ensued. These collections were focused on trees in the Myrtales, including but not exclusively *Eucalyptus* (2, 10, 22, 23, 25–27, 35, 36, 54, 55, 57–61, 65, 96, 98, 102, 106, 120, 126, 145, 147–150). When the first monographic treatment of these fungi by Gryzenhout et al. (63) appeared, five species of *Chrysosporthe* as well as several species in the novel genera, i.e., *Amphilogia*, *Aurapex*, *Celoportbe*, *Holocryphia*, *Microthia*, *Rostraureum*,

and *Ursicollum*, had been described (53–55, 58, 96). Descriptions of these new taxa were all supported by morphological characteristics and DNA sequence data for multiple informative gene regions. In most cases, those species occurring on non-*Eucalyptus* hosts were shown in pathogenicity trials to be able to infect *Eucalyptus* (55, 96, 111). This raised the possibility that the large number of Cryphonectriaceae on native trees in the tropics and Southern Hemisphere should be considered a serious threat to the health of the Myrtales in areas where these fungi are not native.

The Gryzenhout et al. (63) monograph provided a robust foundation for the collection and descriptions of new taxa in the Cryphonectriaceae (**Figure 2**). Most of these were on the Myrtales from the tropics and Southern Hemisphere, but they also included some species, specifically of those associated with Fagales, from the Northern Hemisphere (20, 33, 75–77). As of 2024, based on DNA sequence data, 28 genera and more than 70 species are known in the Cryphonectriaceae (**Figure 2**, **Figure 3**). Of these, 15 genera are known specifically in the Myrtales (**Figure 3**, **Table 2**). Among the genera known from the Southern Hemisphere, *Chrysosporthe* and *Celoporthe* include the greatest number of species (**Figure 3**). A caveat here is that this representation of species abundance more likely reflects the intensity of surveys and collections rather than the actual size of the genera.

HOST SHIFTS AND AN ENDOPHYTIC LIFESTYLE

The first discovery of a species in the Cryphonectriaceae on *Pleroma* (Melastomataceae) marked a substantial change in understanding the diversity and biology of these fungi as tree pathogens (156). Where surveys could be undertaken, they initially focused on the plants in the Melastomataceae (2, 50, 55, 57, 60, 82, 102, 120, 125). These shrubs and trees, including approximately 173 genera and 5,858 species, are mostly native to the New World tropics (132). Of these, many grow in areas where plantations of eucalypts have been established, and several species such as *Miconia crenata* (syn. *Clidemia birta*) and *Melastoma malabathricum* are seriously problematic weeds in these environments (40). New or already known species began to be discovered on these plants, adding confidence that some of those occurring and causing canker disease on *Eucalyptus* had undergone host shifts from plants in the native environment (65, 96) (**Table 2**).

A realization that the Cryphonectriaceae on eucalypts established as exotics in plantations likely originated on native woody plants, not only the Melastomataceae and Myrtaceae but possibly trees in other families of the Myrtales, led to surveys being expanded. These, for example, included trees in the Combretaceae and Lythraceae, such as *Galpinia*, *Lagerstroemia*, *Sonneratia*, and *Terminalia*, that were subsequently also found to be hosts of these fungi (10, 23, 58, 118, 131, 145, 148, 149). Interestingly, some species such as *Aurifilum cerciana*, *Aurifilum terminali*, and *Chrysomorbus lagerstroemiae* found in China were shown, using artificial inoculations, to be able to infect and cause disease on *Eucalyptus* (148, 149) (**Table 2**).

Numerous genera and species of Cryphonectriaceae are found on native trees and shrubs residing in various genera and species of the Myrtales (**Table 2**). Most of these have also been shown to infect *Eucalyptus* in inoculation trials, although only a few have been found naturally infecting these trees (**Table 2**). There are few records of the Cryphonectriaceae on *Eucalyptus* or other Myrtaceae in Australia, the area of the world where most of these trees are native. Interestingly, *Cbr. deuterocubensis* was first found on *Pleroma heteromallum* (syn. *Tibouchina heteromalla*) in the Cairns Botanic Gardens, Queensland, Australia (104). Intriguingly, DNA sequence data have shown that this is the same species that was collected many years earlier by Davison & Coates (39) on the roots of *E. marginata* in Western Australia, an area with a climate atypical for *Cbr. deuterocubensis*. The only other species commonly found on *Eucalyptus* in Australia is a *Holocryphia* sp., which was once considered to be *Holocryphia eucalypti*, known from South Africa (26, 54, 62, 66), but likely

Eucalypts: a general term referring to the trees belonging to the Myrtaceae (myrtle family), including species of *Angophora*, *Corymbia*, and *Eucalyptus*




Figure 3
 Schematic overview of the phylogeny of Cryphonectriaceae species known in culture. Numbers in triangles represent species number in each genus. Genera occurring on Myrtales in the tropics and Southern Hemisphere are indicated in bold. Where genome sequences are available, the list reflects the number of species sequenced but not strains, which in some cases, such as *Cryphonectria parasitica*, is at least 90.

represents a cryptic and distinct species (26). This fungus appears to have a relatively wide host range and to occur on trees in regions that have temperate or subtropical environments (16, 17, 54). It is thought to be native to Australia (16, 17, 39, 62, 100, 151, 158) and has also been found on *Pleroma urvilleanum* (syn. *Tibouchina urvilleana*), presumably having undergone a host shift to that non-native tree, commonly planted as an ornamental (66).

A survey of native *Metrosideros angustifolia* in the Western Cape Province of South Africa, which has a Mediterranean climate, led to the discovery of numerous new species of *Holocryphia* and a single species in a new genus *Diversimorbus*, causing cankers on these trees (26). A population

Table 2 Species of Cryphonectriaceae found on native *Myrtales* hosts and *Eucalyptus* hosts in the tropics and Southern Hemisphere

Genus	Species	Distribution	<i>Eucalyptus</i> hosts	Other <i>Myrtales</i> hosts	Reference(s)
<i>Aurantiosacculus</i>	<i>Aurantiosacculus acutatus</i>	Australia	<i>Eucalyptus viminalis</i>	NA	36
	<i>Aurantiosacculus eucalyptorum</i>	Australia	<i>Eucalyptus globulus</i>	NA	36
<i>Aurapex</i>	<i>Aurapex penicillata</i>	Colombia	<i>Eucalyptus grandis</i>	<i>Miconia theaezans</i> , <i>Miconia lineatum</i> <i>Andesantbus lepidotus</i> (syn. <i>Tibouchina lepidota</i>) <i>Pleroma urvilleanum</i> (syn. <i>Tibouchina urvilleana</i>)	50, 55
<i>Aurifilum</i>	<i>Aurifilum cerciana</i>	China	<i>E. grandis</i> hybrids ^a	<i>Terminalia neotaliala</i>	148
	<i>Aurifilum marmelostoma</i>	Cameroon Taiwan	NA	<i>Terminalia mantaly</i> , <i>Terminalia ivorensis</i>	10, 118
	<i>Aurifilum terminali</i>	China	<i>E. grandis</i> hybrids ^a	<i>T. neotaliala</i>	149
<i>Celoportha</i>	<i>Celoportha borbonica</i>	La Réunion	<i>E. grandis</i> ^a	<i>Pleroma heteromallum</i> (syn. <i>Tibouchina grandifolia</i>)	2
	<i>Celoportha cerciana</i>	China	<i>E. grandis</i> hybrids	NA	150
	<i>Celoportha dispersa</i>	South Africa Zambia	<i>E. grandis</i> ^a	<i>Syzygium cordatum</i> , <i>Syzygium guineense</i> , <i>Syzygium legatii</i> <i>Pleroma granulosum</i> (syn. <i>Tibouchina granulosa</i>) <i>Heteropyxis canescens</i>	96, 145
	<i>Celoportha eucalypti</i>	China	<i>Eucalyptus</i> sp.	<i>Syzygium jambos</i>	22, 149
	<i>Celoportha foliorum</i>	Indonesia	NA	<i>Syzygium</i> sp.	20
	<i>Celoportha fontana</i>	Zambia	<i>E. grandis</i> ^a	<i>S. guineense</i>	147
	<i>Celoportha guangdongensis</i>	China	<i>Eucalyptus</i> sp.	<i>S. jambos</i> <i>Sonneratia apetala</i>	22, 131
	<i>Celoportha hauoliensis</i>	USA (Hawaii)	NA	<i>Psidium cattleianum</i> <i>S. jambos</i>	114
	<i>Celoportha hawaiiensis</i>	USA (Hawaii)	NA	<i>P. cattleianum</i> <i>Syzygium</i> sp., <i>S. jambos</i>	114
	<i>Celoportha indonesiensis</i>	Indonesia	NA	<i>Syzygium aromaticum</i>	22
	<i>Celoportha paradisiaca</i>	USA (Hawaii)	NA	<i>P. cattleianum</i> <i>S. jambos</i>	114

(Continued)

Table 2 (Continued)

Genus	Species	Distribution	<i>Eucalyptus</i> hosts	Other Myrtales hosts	Reference(s)
	<i>Celoportbe syzygii</i>	China	<i>E. grandis</i> hybrids ^a , <i>Eucalyptus urophylla</i> hybrids	<i>Psidium guajava</i> <i>Syzygium cumini</i> , <i>Syzygium hancei</i> , <i>S. jambos</i> , <i>Syzygium samarangense</i>	22, 149
	<i>Celoportbe tibouchinae</i>	La Réunion	<i>E. grandis</i> ^a	<i>P. heteromallum</i> (syn. <i>T. grandifolia</i>)	2
	<i>Celoportbe woodiana</i>	South Africa Mozambique	<i>E. grandis</i> ^a	<i>P. granulosum</i> (syn. <i>T. granulosa</i>) <i>S. cordatum</i>	82, 147
<i>Chrysomorbis</i>	<i>Chrysomorbis lagerstroemiae</i>	China	<i>E. urophylla</i> × <i>E. grandis</i> ^a	<i>Lagerstroemia speciosa</i>	23
<i>Chrysoportha</i>	<i>Chrysoportha austroafricana</i>	South Africa Mozambique Malawi Zambia Namibia Eswatini	<i>E. grandis</i>	<i>Corymbia henryi</i> <i>S. cordatum</i> , <i>S. guineense</i> , <i>Syzygium apiculatum</i> <i>P. granulosum</i> (syn. <i>T. granulosa</i>) <i>Dissotis</i> sp. <i>Eugenia capensis</i> <i>P. guajava</i>	65, 82, 90, 98, 112, 127, 145, 157
	<i>Chrysoportha brasiliensis</i>	Brazil	NA	<i>Rhynchantbera grandiflora</i> <i>Miconia theaezans</i> , <i>Miconia ibaguensis</i>	120
	<i>Chrysoportha colombiana</i>	Colombia	<i>E. grandis</i> ^a	<i>Henriettea seemannii</i>	125
	<i>Chrysoportha cubensis</i>	USA (Florida) Colombia Brazil Venezuela Cuba Suriname Ghana Democratic Republic of Congo Congo Cameroon	<i>Eucalyptus angulosa</i> , <i>Eucalyptus botryoides</i> , <i>Eucalyptus camaldulensis</i> , <i>Eucalyptus dagambae</i> , <i>Eucalyptus deglupta</i> , <i>E. grandis</i> , <i>Eucalyptus microcorys</i> , <i>Eucalyptus paniculata</i> , <i>Eucalyptus pilularis</i> , <i>Eucalyptus propinqua</i> , <i>Eucalyptus robusta</i> , <i>Eucalyptus saligna</i> , <i>Eucalyptus tereticornis</i> , <i>Eucalyptus trabutii</i> , <i>E. urophylla</i>	<i>S. aromaticum</i> <i>M. theaezans</i> , <i>Miconia rubiginosa</i> , <i>Miconia sericea</i> (syn. <i>Clidemia sericea</i>) <i>Lagerstroemia indica</i> <i>Rhynchantbera mexicana</i> <i>Corymbia citriodora</i> , <i>Corymbia maculata</i> <i>P. granulosum</i> (syn. <i>T. granulosa</i>), <i>P. heteromallum</i> (syn. <i>Tibouchina heteromalla</i>), <i>P. urvilleanum</i> (syn. <i>T. urvilleana</i>) <i>Chaetogastra mollis</i> (syn. <i>Tibouchina mollis</i>) <i>Plinia edulis</i> (syn. <i>Marlierea edulis</i>) <i>P. cattleianum</i>	6, 8, 50, 56, 60, 63, 98, 101, 111, 115, 122, 133

(Continued)

Table 2 (Continued)

Genus	Species	Distribution	<i>Eucalyptus</i> hosts	Other Myrtales hosts	Reference(s)
	<i>Chrysoporthe deuterocubensis</i>	Indonesia Thailand Singapore Australia China Taiwan Malaysia Vietnam USA (Hawaii) Tanzania Kenya Mozambique Malawi	<i>Eucalyptus marginata</i> , <i>E. grandis</i> , <i>E. grandis</i> × <i>Eucalyptus pellita</i> , <i>E. urophylla</i> × <i>E. grandis</i> , <i>E. camaldulensis</i>	<i>Melastoma malabatricum</i> , <i>Melastoma candidum</i> , <i>Melastoma sanguineum</i> <i>S. aromaticum</i> , <i>S. cumini</i> , <i>S. jambos</i> , <i>S. samarangense</i> <i>P. urvilleanum</i> (syn. <i>T. urvilleana</i>), <i>P. heteromallum</i> (syn. <i>T. heteromalla</i>) <i>Eugenia</i> sp. <i>P. guajava</i> , <i>P. cattleianum</i> <i>Metrosideros polymorpha</i>	21, 41, 56, 60, 63, 98, 104, 106, 114, 128, 130, 133, 149
	<i>Chrysoporthe doradensis</i>	Ecuador Colombia Brazil	<i>E. grandis</i> , <i>E. deglupta</i> , <i>E. grandis</i> × <i>E. urophylla</i>	<i>R. grandiflora</i> <i>P. granulosum</i> (syn. <i>T. granulosa</i>) <i>H. seemannii</i>	57, 119, 122, 125
	<i>Chrysoporthe hodgesiana</i>	Colombia	<i>E. grandis</i> ^a	<i>Pleroma semidecandrum</i> (syn. <i>Tibouchina semidecandra</i>), <i>P. urvilleanum</i> (syn. <i>T. urvilleana</i>) <i>A. lepidotus</i> (syn. <i>T. lepidota</i>) <i>M. theaezans</i>	56, 111
	<i>Chrysoporthe inopina</i>	Colombia	<i>E. grandis</i> ^a	<i>P. semidecandrum</i> (syn. <i>T. semidecandra</i>), <i>P. urvilleanum</i> (syn. <i>T. urvilleana</i>) <i>A. lepidotus</i> (syn. <i>T. lepidota</i>)	50, 60
	<i>Chrysoporthe puriensis</i>	Brazil	<i>E. grandis</i> × <i>E. urophylla</i> ^a	<i>Pleroma candolleianum</i> (syn. <i>Tibouchina candolleana</i>), <i>P. granulosum</i> (syn. <i>T. granulosa</i>), <i>P. heteromallum</i> (syn. <i>T. heteromalla</i>), <i>Pleroma mutabile</i> (syn. <i>Tibouchina mutabilis</i>) <i>Tibouchina</i> sp. <i>R. grandiflora</i> <i>M. theaezans</i>	102, 107, 119
	<i>Chrysoporthe syzygiicola</i>	Zambia	<i>E. grandis</i> ^a	<i>S. guineense</i>	27
	<i>Chrysoporthe zambiensis</i>	Zambia Zimbabwe	<i>E. grandis</i>	<i>S. guineense</i>	27, 78
<i>Corticimorbus</i>	<i>Corticimorbus sinomyrti</i>	China	<i>E. urophylla</i> × <i>E. grandis</i> ^a	<i>Rhodomyrtus tomentosa</i>	25

(Continued)

Table 2 (Continued)

Genus	Species	Distribution	Eucalyptus hosts	Other Myrtales hosts	Reference(s)
Cryptometrion	Cryptometrion aestuescens	Indonesia	E. grandis	NA	61
	Cryptometrion metrosideri	New Zealand	NA	Metrosideros sp., Metrosideros excelsa	35
Diversimorbus	Diversimorbus metrosideri	South Africa	E. grandis ^a	Metrosideros angustifolia	26
Holocryphia	Holocryphia capensis	South Africa	E. grandis ^a	M. angustifolia	26
	Holocryphia eucalypti	South Africa Eswatini Uganda	Eucalyptus sp., E. grandis, E. saligna	NA	26, 145
	Holocryphia gleniana	South Africa	E. grandis ^a	M. angustifolia	26
	Holocryphia mzansi	South Africa	E. grandis ^a	M. angustifolia	26
Latruncellus	Latruncellus aurorae	Eswatini	E. grandis ^a	Galpinia transvaalica	145
Microthia	Microthia bavanensis	USA (Hawaii, Florida) Cuba Mexico Portugal (Azores, Madeira) Puerto Rico	Eucalyptus sp.; Eucalyptus botryoides, E. camaldulensis, E. grandis, Eucalyptus microphylla, Eucalyptus occidentalis, E. robusta, E. saligna	S. jambos, S. cumini P. cattleianum	54, 63, 114
Myrtonectria	Myrtonectria myrtacearum	South Africa	E. grandis ^a	Heteropyxis natalensis S. cordatum	2
Myrtoportbe	Myrtoportbe bodenii	Malaysia	E. grandis	NA	106
Parvomorbus	Parvomorbus eucalypti	China	E. urophylla × E. grandis, E. urophylla hybrids	NA	149
	Parvomorbus guangdongensis	China	E. urophylla hybrids, E. grandis hybrids	NA	149
Rostraureum	Rostraureum tropicale	Ecuador	NA	T. ivorensis, Terminalia superba	58
Xanthoportbe	Xanthoportbe myrticola	South Africa	E. urophylla × E. grandis, E. grandis	C. benryi	126

^aArtificial inoculation.
Abbreviation: NA, not available.

Endophytes: fungi that live in healthy plant tissues without causing visible disease for part of their life cycle

genetics study on *Holocryphia capensis* surprisingly suggested that it was likely introduced from an as yet unknown origin (24). All these fungi were shown to be able to infect and cause disease on *Eucalyptus* in a pathogenicity trial (24).

As new species of Cryphonectriaceae from many native trees and shrubs in the Myrtales have been discovered, it has become increasingly clear that they exist mostly as asymptomatic endophytes on their natural hosts. This has been most obvious for collections of these fungi on the Melastomataceae in Colombia, where damaged or fallen branches of asymptomatic plants have

been examined and found to bear fruiting structures (**Figure 1**) of these fungi (60). Mause-Sitoe et al. (82) tested this hypothesis by incubating fully asymptomatic stem tissue excised from various species of *Syzygium*, *Eugenia*, *Psidium* (Myrtaceae), and *Dissotis* (Melastomataceae) collected in Mozambique under continuously humid conditions. Fruiting structures of *Chr. austroafricana* developed on the majority of these samples after two months (82). A metagenomic study on *Eucalyptus* has also confirmed that many fungi, including those in groups such as the Diaporthales, which accommodates pathogens such as the Cryphonectriaceae, occur as asymptomatic endophytes in healthy trees (81). This is likely true for many of the woody plants in the Myrtales considered in this review.

Assuming that the Cryphonectriaceae would be present as endophytes in healthy trees and likely fruit under conditions of stress, Roux et al. (114) collected samples from the non-native Myrtaceae, showing dieback due to rust (*Austropuccinia psidii*) infection in Hawaii. This study resulted in a remarkably diverse collection of Cryphonectriaceae, including two known pathogens of *Eucalyptus*, *Chr. deuterocubensis* and *Microthia havanensis*, and three undescribed species of *Celoporthe*. The origin of these fungi is unknown, but they probably include some that have been accidentally introduced into Hawaii and others that are native. But evidence suggests that they are asymptomatic endophytes on the host trees and have probably originated from other trees in the environment (114).

Owing to their importance as pathogens of eucalypts, surveys focused on understanding the global occurrence and relevance of the Cryphonectriaceae have focused on plantations of these trees. Consequently, these have been mostly in the tropics and subtropics where non-native eucalypts have been established in intensively managed plantations. Under these situations, the three species *Chr. cubensis*, *Chr. deuterocubensis*, and *Chr. austroafricana* in South America, Southeast Asia, and Southern Africa, respectively, are most commonly associated with obvious canker diseases. However, as new areas have increasingly been investigated, several other species, known only on native Myrtales or of unknown origin, have been found causing cankers on eucalypts. For example, *Chrysoporthe doradensis* was first described on *Eucalyptus grandis* and *Eucalyptus deglupta* in Ecuador (57) and subsequently found on *Eucalyptus* and native *Rhynchanthera grandiflora* and *Pleroma granulolum* (syn. *Tibouchina granulosa*) (Melastomataceae) in Brazil (119, 122) and most recently on *Eucalyptus* as well as *Henriettea seemannii* (Melastomataceae) in Colombia (125). Likewise, various newly described genera and species have been found causing stem cankers on *Eucalyptus*, but their origins remain to be determined. Examples include *Myrtoporthe* in Malaysia (Sabah) (106), *Parvomorbus* in south China (149), and *Xanthoporthe* in South Africa (126).

POPULATION GENETICS AND PATHWAYS OF MOVEMENT

Observational evidence and patterns of occurrence of the Cryphonectriaceae on the Myrtales have provided strong evidence for many examples of these fungi moving from native trees to infect planted trees. As discussed above, most examples involve these fungi becoming pathogens of eucalypts in established plantations (65, 98, 111). However, there are a few cases in which the Cryphonectriaceae have apparently moved from eucalypts to non-native ornamental trees such as *Pleroma* (syn. *Tibouchina*) (115, 136).

Genetic diversity studies have been conducted on various species of Cryphonectriaceae in an effort to determine the areas of origin of some of these fungi. The first of these studies was conducted on populations of *Chr. cubensis* in Brazil (143) and *Chr. austroafricana* in Southern Africa (141, 146), utilizing vegetative compatibility groups as markers. They provided evidence that populations of these fungi were relatively diverse, but they were not sufficiently informative to provide evidence of origin. Once it became recognized that these fungi occur on native plants, and when

Heterothallic:

fungi that require the presence of a compatible individual for sexual reproduction

MAT1-1 and**MAT1-2 idiomorphs:**

mating in ascomycete fungi is controlled by the *MAT1* locus, with *MAT1-1* and *MAT1-2* variants determining sexual reproduction mode

Homothallic:

fungi species able to undergo sexual reproduction in the absence of a partner of opposite mating type

powerful tools such as microsatellite markers became more readily available, various studies were undertaken to consider the population diversity of other species (24, 94, 95, 97, 101, 136).

To better understand the relationships between Cryphonectriaceae on native and closely associated planted trees, population genetic studies have continued to be undertaken. The most important constraint underpinning these studies remains the difficulty involved in establishing collections of these fungi on native hosts. These trees and shrubs mostly occur naturally in inaccessible areas, such as tropical jungles, mountainous areas, and regions of the world where security issues can be problematic. Yet some such collections have been made, notably for *Chrysosporthe cubensis* and *Cbr. puriensis* in South America (101, 136), *Cbr. austroafricana* (127) and *Holocryphia* spp. in Southern Africa (24, 94), and *Cbr. deuteroecubensis* in Southeast Asia (128). Results of all these studies have provided robust evidence of the Cryphonectriaceae occurring naturally on native Myrtales and undergoing host shifts to related planted trees.

The increasing availability of whole-genome sequences for fungi and, in this case, the Cryphonectriaceae from the tropics and Southern Hemisphere will provide many opportunities to better understand their biology, native origins, and pathways of movement. Genomes are currently available for most species of *Chrysosporthe* (80, 135, 138, 152, 153), and projects are well advanced to sequence species in all the genera (**Figure 3**). From those sequence data already available, it has been possible to determine that *Chrysosporthe* includes heterothallic species having the *MAT1-1* and *MAT1-2* idiomorphs as well as homothallic species with both idiomorphs in single haplotypes. This knowledge of mating type distribution adds information regarding the structure of populations and whether they represent native or introduced lineages (83).

EMERGING PATHOGENS AND DISEASE THREATS

Species of the Cryphonectriaceae occurring on trees in the Myrtales include highly aggressive pathogens that have the capacity to kill trees. Owing to their importance as canker pathogens of eucalypts, the best known of these are species of *Chrysosporthe*. Most of these species have been found causing cankers on eucalypts that have been established in plantations in the tropics and Southern Hemisphere. And many of these have now been shown to have originated on native Myrtales growing in close proximity to the plantations of non-native eucalypts. Beyond natural infections, most species of the Cryphonectriaceae residing in various genera and currently known only from native Myrtales have been shown in artificial inoculations to be pathogens of eucalypts.

The only example of serious tree diseases caused by the Cryphonectriaceae arising from host shifts from native Myrtales is the canker disease on eucalypts. There is no evidence of other species of the Cryphonectriaceae having emerged as disease agents on trees in natural ecosystems. Given the fact that those trees mostly occur in remote areas, it seems reasonable to assume that any emerging pathogen would probably not be recorded until it had begun to cause serious damage and undergone substantial spread. This raises the interesting question as to how fast a new disease caused by one of these fungi as a novel pathogen on a naïve host might emerge and spread.

A fundamental difference between woody ecosystems in the Northern and Southern Hemispheres is the diversity of plants in the south. Whereas the Northern Hemisphere mostly includes only the single Boreal floral Kingdom with relatively few tree genera, the remaining five floral kingdoms occur mostly in the Southern Hemisphere and include a much greater diversity of woody plants (32). It would be reasonable to assume that rapidly emerging epidemics would occur in areas where a single species covers large areas. If this is a major defining factor, then epidemic levels of disease caused by accidentally introduced Cryphonectriaceae in the Myrtales might not

occur as dramatically as in the case with various invasive pathogens in the Northern Hemisphere, such as chestnut blight (110) and Dutch elm disease (46).

Even if extensive disease epidemics caused by accidentally introduced Cryphonectriaceae would not occur in native populations of trees in the Myrtales, serious losses to plants and associated plant diversity might still occur. An apt example is found in the case of myrtle rust caused by *Austropuccinia psidii* (9). This pathogen is believed to be native to South America (34, 48, 49), and at least two genotypes have been accidentally moved across countries of the Southern Hemisphere (38, 84, 124). After many years of anticipating its arrival in Australia, the pandemic strain of *A. psidii* reached Australia in 2010 (18, 103). This pathogen has a very wide host range, which might be similar to some species of Cryphonectriaceae, and it has rapidly emerged to infect a large number of hosts (34, 123). But there are only some cases in which species of trees are sufficiently susceptible to be considered as highly threatened (19). The example of myrtle rust might provide a reasonable view of what to expect from possible accidental introductions of species of Cryphonectriaceae across the Southern Hemisphere.

Plantation forestry using non-native eucalypts, including species of *Eucalyptus* and *Corymbia*, gave rise to the first concern that Cryphonectria canker might emerge as a serious limiting factor. From a commercial perspective, the disease has been very effectively managed by establishing planting stock known to be tolerant to infection by the Cryphonectriaceae, which for the present include mostly species of *Chrysosporthe*. One of the most important approaches in this case has been to utilize clonal hybrids between different species of *Eucalyptus* and, in many cases, test these in greenhouse and field screening trials. Such trials have been conducted in South America, South Africa, and Southeast Asia for *Chr. cubensis* (1, 79, 111, 122), *Chr. austroafricana* (139), and *Chr. deuterocubensis* (21, 106, 130). These have also included tests using different genotypes of the pathogens and considering issues such as clone-by-genotype interactions (1, 139).

The fact that the Cryphonectriaceae infecting *Eucalyptus* originate in the natural environment where these trees are being planted implies that disease-resistant clones will continuously be challenged by genetically diverse populations of the pathogens. Consequently, clones currently resistant to the pathogens might not be so after relatively short periods of time. A recent example can be found in the first example of a *Chrysosporthe* sp., in this case, *Chr. austroafricana*, infecting a species of *Corymbia* (127). This exemplifies the so-called host–pathogen arms race (28) and the need for forestry companies to introduce new genetic stock that will not be damaged by disease (14). Ultimately, technologies such as genetic modification of the planting stock will likely be required to maintain the sustainability of *Eucalyptus* forestry operations. The availability of a full sequence of the *E. grandis* genome (89), as well as of the pathogen genome, will provide the tools to ensure that disease-tolerant planting stock remains available.

Although new genotypes of known Cryphonectriaceae will continuously emerge to challenge eucalypts in plantations, new genera and species of these genera are likely to also emerge, adding greater challenges to plantation forestry. This has already been clear in recent outbreaks of Cryphonectria canker on *Eucalyptus* clonal hybrids in Indonesia, Malaysia, and South Africa. In these cases, species residing in the new genera *Cryptometrion*, *Myrtoportbe*, and *Xanthoportbe* have emerged to cause serious disease problems (61, 106, 126). Others will surely follow in the future.

FUTURE PROSPECTS

The Cryphonectriaceae on trees in the tropics and Southern Hemisphere provide a vivid example of the huge biodiversity of fungi in an area of the world that has been relatively poorly sampled. As important and emerging pathogens, they also illustrate the risks of serious tree disease problems

Host–pathogen arms race: a coevolutionary race in which plants and pathogens adapt to each other, requiring continual development of resistant cultivars and clones

emerging in the future. This is especially true as the movement of people and products continues to increase globally, thus facilitating increasing risks of accidental introductions into new areas (15, 105). This also argues for much greater attention being given to documenting and understanding the very poorly explored global fungal biodiversity.

It is relevant that many of these Cryphonectriaceae give rise to cankers on trees that are biologically similar to the disease caused by the chestnut blight fungus, *C. parasitica*. They infect their hosts easily through naturally occurring wounds and on susceptible hosts develop rapidly in the cambial tissues to girdle stems (**Figure 1**). Consequently, they have the capacity to cause diseases as serious as chestnut blight. Thus, if a species naturally occurring on a particular tree genus were to be accidentally introduced into a new area where a susceptible native congener occurs as a naïve host, a very serious epidemic could easily arise (5, 37, 155). This view is based on many similar situations in the Northern Hemisphere, and the only limiting factor might be a greater host diversity in the tropics and Southern Hemisphere.

The discovery of the Cryphonectriaceae and their importance as tree pathogens on the Myrtales first emerged from planting *Eucalyptus* spp. outside their native range. These trees have now been established in plantations over very large areas, particularly in South America, Africa, and Southeast Asia. This area is currently estimated to be approximately 22 million hectares, making eucalypts the most widely planted genus of trees in the world (116). Most of the areas where these plantations occur are also areas of a rich biodiversity of trees and shrubs in the Myrtales. Fungi in the Cryphonectriaceae and likely others represent significant future threats to planted eucalypts via host shifts that are already evident in the case of numerous species.

Species of the Cryphonectriaceae that have undergone host shifts to infect eucalypts have effectively emerged as novel pathogens of those trees. The intensive nature of plantation forestry results in the pathogen populations building up such that the chance of their being moved to new areas increases markedly. This represents a situation known as a bridgehead effect (11), where pathogens or pests, once introduced into new areas, are most likely to be moved again. In effect, planting trees that typically have very long lifespans provides land bridges for tree pathogens to move globally (155).

All currently available information suggests that tree diseases caused by the Cryphonectriaceae will become increasingly important in the future. Consequently, efforts to reduce this threat must be vigorously pursued. This should include a better understanding of the possible pathways of movement of these fungi and identifying means to reduce new invasions. Surveys of diseases on Myrtales in their natural environment should also be supported and pursued. Tools such as those relying on data from genomes to identify and understand the biology and ecology of these fungi are also becoming increasingly powerful and can be utilized, for example, in improving diagnostics and quarantine procedures (85). Importantly, companies involved in commercially exploiting trees in the Myrtales should be made aware of the risks of accidentally moving pathogens and pests such as those in the Cryphonectriaceae. This would bear advantages not only to their business ventures but also to the health of forests and forestry globally.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We are grateful to many colleagues and students who have participated in our studies on the Cryphonectriaceae, a period that has extended over approximately 40 years. These studies have

had the benefit of financial and in-kind support from companies pursuing plantation forestry, particularly with eucalypts in many countries of the tropics and Southern Hemisphere. The forestry companies that have been members of the Tree Protection Co-Operative Program in South Africa, a relationship that has lasted more than 35 years, deserve special thanks. We also thank the National Research Foundation, the Department of Science and Innovation, South Africa, and the University of Pretoria for long-term financial support.

LITERATURE CITED

1. Alfenas AC, Jeng R, Hubbes M. 1983. Virulence of *Cryphonectria cubensis* on *Eucalyptus* species differing in resistance. *Eur. J. For. Pathol.* 13:197–205
2. Ali DB, Marincowitz S, Wingfield MJ, Roux J, Crous PW, McTaggart AR. 2018. Novel *Cryphonectriaceae* from La Réunion and South Africa, and their pathogenicity on *Eucalyptus*. *Mycol. Prog.* 17:953–66
3. Anagnostakis SL. 1987. Chestnut blight: the classical problem of an introduced pathogen. *Mycologia* 79:23–37
4. Anderson PJ, Anderson HW. 1912. The chestnut blight fungus and a related saprophyte. *Phytopathology* 2:204–10
5. Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P. 2004. Emerging infectious diseases of plants: pathogen pollution, climate change and agrotechnology drivers. *Trends Ecol. Evol.* 19:535–44
6. Apetorgbor MM, Roux J. 2015. Diseases of plantation forestry trees in Southern Ghana. *Int. J. Phytopathol.* 4:5–13
7. Barr ME. 1978. The Diaporthales in North America with emphasis on *Gnomonia* and its segregates. *Mycol. Mem.* 7:1–232
8. Barreto RW, Rocha FB, Ferreira FA. 2006. First record of natural infection of *Marlierea edulis* by the eucalyptus canker fungus *Chrysosporium cubensis*. *Plant Pathol.* 55:577
9. Beenken L. 2017. *Austropuccinia*: a new genus name for the myrtle rust *Puccinia psidii* placed within the redefined family Sphaerophragmiaceae (Pucciniales). *Phytotaxa* 297:53–61
10. Begoude ADB, Gryzenhout M, Wingfield MJ, Roux J. 2010. *Aurifilum*, a new fungal genus in the Cryphonectriaceae from *Terminalia* species in Cameroon. *Antonie van Leeuwenhoek* 98:263–78
11. Bertelsmeier C, Keller L. 2018. Bridgehead effects and role of adaptive evolution in invasive populations. *Trends Ecol. Evol.* 33:527–34
12. Boerboom JHA, Maas PWT. 1970. Canker of *Eucalyptus grandis* and *E. saligna* in Surinam, caused by *Endothia havanensis*. *Turrialba* 20:94–99
13. Bruner SC. 1917. *Una Enfermedad Gangrenosa de los Eucaliptos*. Estación Central Agronómica de Cuba
14. Burdon RD. 2001. Genetic diversity and disease resistance: some considerations for research, breeding, and deployment. *Can. J. For. Res.* 31:596–606
15. Burgess TI, Wingfield MJ. 2017. Pathogens on the move: a 100-year global experiment with planted eucalypts. *Bioscience* 67:14–25
16. Carnegie AJ. 2007. Forest health condition in New South Wales, Australia, 1996–2005. I. Fungi recorded from eucalypt plantations during forest health surveys. *Australas. Plant Pathol.* 36:213–24
17. Carnegie AJ. 2007. Forest health condition in New South Wales, Australia, 1996–2005. II. Fungal damage recorded in eucalypt plantations during forest health surveys and their management. *Australas. Plant Pathol.* 36:225–39
18. Carnegie AJ, Lidbetter JR, Walker J, Horwood MA, Tesoriero L, et al. 2010. *Uredo rangellii*, a taxon in the guava rust complex, newly recorded on Myrtaceae in Australia. *Australas. Plant Pathol.* 39:463–66
19. Carnegie AJ, Pegg GS. 2018. Lessons from the incursion of myrtle rust in Australia. *Annu. Rev. Phytopathol.* 56:457–78
20. Chen Q, Bakhshi M, Balci Y, Broders KD, Cheewangkoon R, et al. 2022. Genera of phytopathogenic fungi: GOPHY 4. *Stud. Mycol.* 101:417–564

21. Chen SF, Gryzenhout M, Roux J, Xie YJ, Wingfield MJ, Zhou XD. 2010. Identification and pathogenicity of *Chrysosporthe cubensis* on *Eucalyptus* and *Syzygium* spp. in South China. *Plant Dis.* 94:1143–50
22. Chen SF, Gryzenhout M, Roux J, Xie YJ, Wingfield MJ, Zhou XD. 2011. Novel species of *Celoporthe* from *Eucalyptus* and *Syzygium* trees in China and Indonesia. *Mycologia* 103:1384–410
23. Chen SF, Liu QL, Li GQ, Wingfield MJ, Roux J. 2018. A new genus of Cryphonectriaceae isolated from *Lagerstroemia speciosa* in southern China. *Plant Pathol.* 67:107–23
24. Chen SF, van der Merwe NA, Wingfield MJ, Roux J. 2016. Population structure of *Holocryphia capensis* (Cryphonectriaceae) from *Metrosideros angustifolia* and its pathogenicity to *Eucalyptus* species. *Australas. Plant Pathol.* 45:201–7
25. Chen SF, Wingfield MJ, Li GQ, Liu FF. 2016. *Corticimorbus sinomyrti* gen. et sp. nov. (Cryphonectriaceae) pathogenic to native *Rhodomyrtus tomentosa* (Myrtaceae) in South China. *Plant Pathol.* 65:1254–66
26. Chen SF, Wingfield MJ, Roux J. 2013. *Diversimorbus metrosiderotis* gen. et sp. nov. and three new species of *Holocryphia* (Cryphonectriaceae) associated with cankers on native *Metrosideros angustifolia* trees in South Africa. *Fungal Biol.* 117:289–310
27. Chungu D, Gryzenhout M, Muimba-Kankolongo A, Wingfield MJ, Roux J. 2010. Taxonomy and pathogenicity of two novel *Chrysosporthe* species from *Eucalyptus grandis* and *Syzygium guineense* in Zambia. *Mycol. Prog.* 9:379–93
28. Clay K, Kover PX. 1996. The Red Queen hypothesis and plant/pathogen interactions. *Annu. Rev. Phytopathol.* 34:29–50
29. Clinton GP. 1908. *Chestnut bark disease*, *Diaporthe parasitica* Murr. Rep., Conn. Agric. Exp. Stn., New Haven
30. Conradie E, Swart WJ, Wingfield MJ. 1990. Cryphonectria canker of *Eucalyptus*, an important disease in plantation forestry in South Africa. *South Afr. For. J.* 152:43–49
31. Conti E, Litt A, Wilson PG, Graham SA, Briggs BG, et al. 1997. Interfamilial relationships in Myrtales: molecular phylogeny and patterns of morphological evolution. *Syst. Bot.* 22(4):629–47
32. Corlett RT. 2016. Plant diversity in a changing world: status, trends, and conservation needs. *Plant Divers.* 38:10–16
33. Cornejo C, Hauser A, Beenken L, Cech T, Rigling D. 2021. *Cryphonectria carpinicola* sp. nov. associated with hornbeam decline in Europe. *Fungal Biol.* 125:347–56
34. Coutinho TA, Wingfield MJ, Alfenas AC, Crous PW. 1998. Eucalyptus rust: a disease with the potential for serious international implications. *Plant Dis.* 82:819–25
35. Crous PW, Osieck ER, Jurjević Ž, Boers J, van Iperen AL, et al. 2021. Fungal Planet description sheets: 1284–1382. *Persoonia* 47:178–374
36. Crous PW, Summerell BA, Alfenas AC, Edwards J, Pascoe IG, et al. 2012. Genera of diaphoralean coelomycetes associated with leaf spots of tree hosts. *Persoonia* 28:66–75
37. Cunningham AA, Daszak P, Rodriguez JP. 2003. Pathogen pollution: defining a parasitological threat to biodiversity conservation. *J. Parasitol.* 89:S78–83
38. da S. Machado P, Alfenas AC, Alfenas RF, Mohammed CL, Glen M. 2015. Microsatellite analysis indicates that *Puccinia psidii* in Australia is mutating but not recombining. *Australas. Plant Pathol.* 44:455–62
39. Davison EM, Coates DJ. 1991. Identification of *Cryphonectria cubensis* and *Endothia gyrosa* from eucalypts in Western Australia using isozyme analysis. *Australas. Plant Pathol.* 20:157–60
40. DeWalt SJ, Denslow JS, Johnson MT. 2022. Ecology and control of naturalized Melastomataceae. In *Systematics, Evolution, and Ecology of Melastomataceae*, ed. R Goldenberg, FA Michelangi, F Almeda, pp. 761–89. Springer
41. Fan MC, Huang CC, Huang JS, Tsai SF, Yeh HC, Hong CF. 2013. First report of *Chrysosporthe deuterocubensis* causing canker on *Syzygium samarangense* in Taiwan. *Plant Dis.* 97:1508
42. Ferreira FA, Alfenas AC, de Freitas AL. 1978. Determinação da resistência de 16 procedências de *Eucalyptus* ao cancro causado por *Diaporthe cubensis* Bruner, no Vale do Rio Doce, Minas Gerais. *Rev. Árvore* 2:119–29
43. Ferreira FA, Reis MS, Alfenas AC, Hodges CS. 1977. Avaliação da resistência de *Eucalyptus* spp. ao cancro causado por *Diaporthe cubensis* Bruner. *Fitopatol. Bras.* 2:225–41

44. Ferreira FA, Reis MS, Alfenas AC, Silva A. 1977. Níveis de incidência natural de cancro em *Eucalyptus* spp. no Nordeste do Estado do Espírito Santo. *Rev. Árvore* 1:89–106
45. Fries EM. 1828. *Elenchus Fungorum: Sistens Commentarium in Systema Mycologicum*. Sumptibus E. Mauriti
46. Gibbs JN. 1978. Intercontinental epidemiology of Dutch elm disease. *Annu. Rev. Phytopathol.* 16:287–307
47. Gibson IAS. 1981. A canker disease of *Eucalyptus* new to Africa. *FAO For. Genet. Resour. Inf.* 10:23–24
48. Glen M, Alfenas AC, Zauza EAV, Wingfield MJ, Mohammed C. 2007. *Puccinia psidii*: a threat to the Australian environment and economy—a review. *Australas. Plant Pathol.* 36:1–16
49. Graça RN, Ross-Davis AL, Klopfenstein NB, Kim MS, Peever TL, et al. 2013. Rust disease of eucalypts, caused by *Puccinia psidii*, did not originate via host jump from guava in Brazil. *Mol. Ecol.* 22:6033–47
50. Granados GM, McTaggart AR, Rodas CA, Roux J, Wingfield MJ. 2020. Species of Cryphonectriaceae occupy an endophytic niche in the Melastomataceae and are putative latent pathogens of *Eucalyptus*. *Eur. J. Plant Pathol.* 156:273–83
51. Gravatt GF, Gill LS. 1930. *Chestnut blight*. Farmer's Bull. No. 1641, US Department of Agriculture
52. Gryzenhout M, Glen HF, Wingfield BD, Wingfield MJ. 2005. (1686) Proposal to conserve the name *Cryphonectria* (Diaporthales) with a conserved type. *Taxon* 54:539–40
53. Gryzenhout M, Glen HF, Wingfield BD, Wingfield MJ. 2005. *Amphilogia* gen. nov. for *Cryphonectria*-like fungi from *Elaeocarpus* spp. in New Zealand and Sri Lanka. *Taxon* 54:1009–21
54. Gryzenhout M, Myburg H, Hodges CS, Wingfield BD, Wingfield MJ. 2006. *Microthia*, *Holocryphia* and *Ursicollum*, three new genera on *Eucalyptus* and *Coccoloba* for fungi previously known as *Cryphonectria*. *Stud. Mycol.* 55:35–52
55. Gryzenhout M, Myburg H, Rodas CA, Wingfield BD, Wingfield MJ. 2006. *Aurapex penicillata* gen. sp. nov. from native *Miconia theaezans* and *Tibouchina* spp. in Colombia. *Mycologia* 98:105–15
56. Gryzenhout M, Myburg H, van der Merwe NA, Wingfield BD, Wingfield MJ. 2004. *Chrysosporthe*, a new genus to accommodate *Cryphonectria cubensis*. *Stud. Mycol.* 50:119–42
57. Gryzenhout M, Myburg H, Wingfield BD, Montenegro F, Wingfield MJ. 2005. *Chrysosporthe doradensis* sp. nov. pathogenic to *Eucalyptus* in Ecuador. *Fungal Divers.* 20:39–57
58. Gryzenhout M, Myburg H, Wingfield BD, Montenegro F, Wingfield MJ. 2005. *Rostrareum tropicale* gen. sp. nov. (Diaporthales) associated with dying *Terminalia ivorensis* in Ecuador. *Mycol. Res.* 109:1029–44
59. Gryzenhout M, Myburg H, Wingfield BD, Wingfield MJ. 2006. Cryphonectriaceae (Diaporthales), a new family including *Cryphonectria*, *Chrysosporthe*, *Endothia* and allied genera. *Mycologia* 98:239–49
60. Gryzenhout M, Rodas CA, Mena Portales J, Clegg P, Wingfield BD, Wingfield MJ. 2006. Novel hosts of the *Eucalyptus* canker pathogen *Chrysosporthe cubensis* and a new *Chrysosporthe* species from Colombia. *Mycol. Res.* 110:833–45
61. Gryzenhout M, Tarigan M, Clegg PA, Wingfield MJ. 2010. *Cryptometrion aestuescens* gen. sp. nov. (Cryphonectriaceae) pathogenic to *Eucalyptus* in Indonesia. *Australas. Plant Pathol.* 39:161–69
62. Gryzenhout M, Vermeulen M, Dick M, Wingfield MJ. 2010. The *Eucalyptus* canker pathogen *Holocryphia eucalypti* on *Eucalyptus* in New Zealand. *Australas. Plant Dis. Notes* 5:5–8
63. Gryzenhout M, Wingfield BD, Wingfield MJ. 2009. *Taxonomy, Phylogeny, and Ecology of Bark-Inhabiting and Tree-Pathogenic Fungi in the Cryphonectriaceae*. APS Press
64. Heath RN, Gryzenhout M, Roux J, Wingfield MJ. 2002. *The discovery of Cryphonectria cubensis on native Syzygium spp. from South Africa*. Paper presented at 7th International Mycological Congress, Oslo, Norway
65. Heath RN, Gryzenhout M, Roux J, Wingfield MJ. 2006. Discovery of the canker pathogen *Chrysosporthe austroafricana* on native *Syzygium* spp. in South Africa. *Plant Dis.* 90:433–38
66. Heath RN, Roux J, Gryzenhout M, Carnegie AJ, Smith IW, Wingfield MJ. 2007. *Holocryphia eucalypti* on *Tibouchina urvilleana* in Australia. *Australas. Plant Pathol.* 36:560–64
67. Heiniger U, Rigling D. 1994. Biological control of chestnut blight in Europe. *Annu. Rev. Phytopathol.* 32:581–99
68. Hepting GH. 1974. Death of the American chestnut. *J. For. Hist.* 18:60–67
69. Hodges C, Reis MS, Ferreira F, Henfling J. 1976. O cancro do eucalipto causado por *Diaporthe cubensis*. *Fitopatol. Bras.* 1:129–67
70. Hodges C, Reis MS, May L. 1973. Duas enfermidades em plantações de essências florestais exóticas no Brasil. *Bras. Florest.* 6:5–12

71. Hodges CS. 1980. Eucalyptus canker in the Pacific area. In *American Pacific Forestry News* (Sept-Oct), pp. 1–2. US Department of Agriculture Forest Service
72. Hodges CS. 1980. The taxonomy of *Diaporthe cubensis*. *Mycologia* 72:542–48
73. Hodges CS, Alfenas AC, Ferreira FA. 1986. The conspecificity of *Cryphonectria cubensis* and *Endothia eugeniae*. *Mycologia* 78:343–50
74. Hodges CS, Geary TF, Cordell CE. 1979. The occurrence of *Diaporthe cubensis* on *Eucalyptus* in Florida, Hawaii and Puerto Rico. *Plant Dis. Rep.* 63:216–20
75. Jiang N, Fan XL, Tian CM. 2019. Identification and pathogenicity of Cryphonectriaceae species associated with chestnut canker in China. *Plant Pathol.* 68:1132–45
76. Jiang N, Fan XL, Tian CM, Crous PW. 2020. Reevaluating Cryphonectriaceae and allied families in Diaporthales. *Mycologia* 112:267–92
77. Jiang N, Fan XL, Yang Q, Du Z, Tian CM. 2018. Two novel species of *Cryphonectria* from *Quercus* in China. *Phytotaxa* 347:243–50
78. Jimu L, Muzhinji N, Magogo C, Mureva E. 2018. *Chrysosporthe zambiensis* detected on native *Syzygium* in Zimbabwe. *Australas. Plant Dis. Notes* 13:1
79. Juárez-Palacios JC, Honorato-Salazar JA, Vázquez-Silva L, Parraguirre-Lezama JF. 2013. Pathogenicity of *Chrysosporthe cubensis* to *Eucalyptus grandis* and *E. urophylla* clones in the Southeastern of Mexico. *Madera Bosques* 19:17–36
80. Kanzi AM, Steenkamp ET, van der Merwe NA, Wingfield BD. 2019. The mating system of the *Eucalyptus* canker pathogen *Chrysosporthe austroafricana* and closely related species. *Fungal Genet. Biol.* 123:41–52
81. Kemler M, Garnas J, Wingfield MJ, Gryzenhout M, Pillay KA, Slippers B. 2013. Ion Torrent PGM as tool for fungal community analysis: a case study of endophytes in *Eucalyptus grandis* reveals high taxonomic diversity. *PLOS ONE* 8:e81718
82. Mause-Sitoe SND, Rodas CA, Wingfield MJ, Chen SF, Roux J. 2016. Endophytic Cryphonectriaceae on native Myrtales: possible origin of Chrysosporthe canker on plantation-grown *Eucalyptus*. *Fungal Biol.* 120:827–35
83. McDonald BA, Linde C. 2002. Pathogen population genetics, evolutionary potential, and durable resistance. *Annu. Rev. Phytopathol.* 40:349–79
84. McTaggart AR, Roux J, Granados GM, Gafur A, Tarrigan M, et al. 2016. Rust (*Puccinia psidii*) recorded in Indonesia poses a threat to forests and forestry in South-East Asia. *Australas. Plant Pathol.* 45:83–89
85. McTaggart AR, van der Nest MA, Steenkamp ET, Roux J, Slippers B, et al. 2016. Fungal genomics challenges the dogma of name-based biosecurity. *PLOS Pathog.* 12:e1005475
86. Merkel HW. 1905. A deadly fungus on the American chestnut. *Annu. Rep. N. Y. Zool. Soc.* 10:97–103
87. Micales JA, Stipes RJ. 1987. A reexamination of the fungal genera *Cryphonectria* and *Endothia*. *Phytopathology* 77:650–54
88. Murrill WA. 1906. A new chestnut disease. *Torreya* 6:186–89
89. Myburg AA, Grattapaglia D, Tuskan GA, Hellsten U, Hayes RD, et al. 2014. The genome of *Eucalyptus grandis*. *Nature* 510:356–62
90. Myburg H, Gryzenhout M, Heath R, Roux J, Wingfield BD, Wingfield MJ. 2002. Cryphonectria canker on *Tibouchina* in South Africa. *Mycol. Res.* 106:1299–306
91. Myburg H, Gryzenhout M, Wingfield BD, Wingfield MJ. 2002. β -Tubulin and histone H3 gene sequences distinguish *Cryphonectria cubensis* from South Africa, Asia, and South America. *Can. J. Bot.* 80:590–96
92. Myburg H, Wingfield BD, Wingfield MJ. 1999. Phylogeny of *Cryphonectria cubensis* and allied species inferred from DNA analysis. *Mycologia* 91:243–50
93. Myburg H, Wingfield BD, Wingfield MJ, Gryzenhout M. 2003. Conspecificity of *Endothia eugeniae* and *Cryphonectria cubensis*: a re-evaluation based on morphology and DNA sequence data. *Mycoscience* 44:187–96
94. Nakabonge G, Burgess T, Gryzenhout M, Wingfield B, Wingfield MJ, Roux J. 2008. Population structure of the fungal pathogen *Holocryphia eucalypti* in Australia and South Africa. *Australas. Plant Pathol.* 37:154–61

95. Nakabonge G, Cortinas MN, Roux J, Gryzenhout M, Wingfield BD, Wingfield MJ. 2005. Development of polymorphic microsatellite markers for the fungal tree pathogen *Cryphonectria eucalypti*. *Mol. Ecol. Notes* 5:558–61
96. Nakabonge G, Gryzenhout M, Roux J, Wingfield BD, Wingfield MJ. 2006. *Celoporthe dispersa* gen. et sp. nov. from native Myrtales in South Africa. *Stud. Mycol.* 55:255–67
97. Nakabonge G, Gryzenhout M, Wingfield BD, Wingfield MJ, Roux J. 2007. Genetic diversity of *Chrysoporthe cubensis* in eastern and southern Africa. *South Afr. J. Sci.* 103:261–64
98. Nakabonge G, Roux J, Gryzenhout M, Wingfield MJ. 2006. Distribution of Chrysoporthe canker pathogens on *Eucalyptus* and *Syzygium* spp. in Eastern and Southern Africa. *Plant Dis.* 90:734–40
99. Nutman FJ, Roberts FM. 1952. Acute die-back of clove trees in the Zanzibar protectorate *Ann. Appl. Biol.* 39:599–608
100. Old KM, Murray DIL, Kile GA, Simpson J, Malafant KWJ. 1986. The pathology of fungi isolated from eucalypt cankers in south-eastern Australia. *Aust. For. Res.* 16:21–36
101. Oliveira MES, Kanzi AM, van der Merwe NA, Wingfield MJ, Wingfield BD, et al. 2022. Genetic variability in populations of *Chrysoporthe cubensis* and *Chr. puriensis* in Brazil. *Australas. Plant Pathol.* 51:175–91
102. Oliveira MES, van der Merwe NA, Wingfield MJ, Wingfield BD, Soares TPF, et al. 2021. *Chrysoporthe puriensis* sp. nov. from *Tibouchina* spp. in Brazil: an emerging threat to *Eucalyptus*. *Australas. Plant Pathol.* 50:29–40
103. Pegg GS, Giblin FR, McTaggart AR, Guymer GP, Taylor H, et al. 2014. *Puccinia psidii* in Queensland, Australia: disease symptoms, distribution and impact. *Plant Pathol.* 63:1005–21
104. Pegg GS, Gryzenhout M, O'Dwyer C, Drenth A, Wingfield MJ. 2010. The *Eucalyptus* canker pathogen *Chrysoporthe* discovered in eastern Australia. *Australas. Plant Pathol.* 39:343–49
105. Raffa KF, Brockerhoff EG, Grégoire JC, Hamelin RC, Liebhold AM, et al. 2023. Approaches to forecasting damage by invasive forest insects and pathogens: a cross-assessment. *Bioscience* 73:85–111
106. Rauf MRBA, McTaggart AR, Marincowitz S, Barnes I, Japarudin Y, Wingfield MJ. 2020. Pathogenicity of *Chrysoporthe deuterocubensis* and *Myrtoporthe bodenii* gen. et sp. nov. on *Eucalyptus* in Sabah, Malaysia. *Australas. Plant Pathol.* 49:53–64
107. Rêgo GMS, de Souza IAL, Silva GA, de Oliveira MES, Ferreira MA. 2023. *Chrysoporthe puriensis* causing canker and mortality in *Pleroma mutabile* in the Atlantic Forest, Brazil. *For. Pathol.* 53:e12803
108. Rehm H. 1907. Ascomycetes exs. Fasc. 39. *Ann. Mycol.* 5:207–13
109. Reid J, Booth C. 1969. Some species segregated from the genera *Cryptospora*, *Cryptosporella*, and *Sillia*. *Can. J. Bot.* 47:1055–60
110. Rigling D, Prospero S. 2018. *Cryphonectria parasitica*, the causal agent of chestnut blight: invasion history, population biology and disease control. *Mol. Plant Pathol.* 19:7–20
111. Rodas CA, Gryzenhout M, Myburg H, Wingfield BD, Wingfield MJ. 2005. Discovery of the Eucalyptus canker pathogen *Chrysoporthe cubensis* on native *Miconia* (Melastomataceae) in Colombia. *Plant Pathol.* 54:460–70
112. Roux J, Meke G, Kanyi B, Mwangi L, Mbaga A, et al. 2005. Diseases of plantation forestry trees in eastern and southern Africa. *South Afr. J. Sci.* 101:409–13
113. Roux J, Myburg H, Wingfield BD, Wingfield MJ. 2003. Biological and phylogenetic analyses suggest that two *Cryphonectria* spp. cause cankers of *Eucalyptus* in Africa. *Plant Dis.* 87:1329–32
114. Roux J, Nkuekam GK, Marincowitz S, van der Merwe NA, Uchida J, et al. 2020. Cryphonectriaceae associated with rust-infected *Syzygium jambos* in Hawaii. *Mycologia* 176:49–79
115. Seixas CDS, Barreto RW, Alfenas AC, Ferreira FA. 2004. *Cryphonectria cubensis* on an indigenous host in Brazil: a possible origin for eucalyptus canker disease? *Mycologist* 18:39–45
116. Seng Hua L, Wei Chen L, Antov P, Kristak L, Md Tahir P. 2022. Engineering wood products from *Eucalyptus* spp. *Adv. Mater. Sci. Eng.* 2022:8000780
117. Sharma JK, Mohanan C, Florence EJM. 1985. *Disease survey in nurseries and plantations of forest tree species grown in Kerala*. Rep., Kerala Forest Research Institute, Kerala, India
118. Shen YM, Liu HL, Huang TC, Pai KF. 2016. First report of *Aurifilum marmelostoma* causing cankers on *Terminalia mantaly* in Taiwan. *J. Gen. Plant Pathol.* 82:216–19

119. Silva GA, Moreira SI, Ferreira MA. 2023. New Melastomataceae hosts of *Chrysosporthe* species in Brazil. *J. Phytopathol.* 171:716–22
120. Silva GA, Oliveira MES, Rêgo GMS, Wingfield BD, Wingfield MJ, Ferreira MA. 2024. *Chrysosporthe brasiliensis* sp. nov. pathogenic to Melastomataceae in southeast Brazil. *Fungal Biol.* 128:1790–99
121. Slippers B, Stenlid J, Wingfield MJ. 2005. Emerging pathogens: fungal host jumps following anthropogenic introduction. *Trends Ecol. Evol.* 20:420–21
122. Soares TPF, Ferreira MA, Mafia RG, Oliveira LSS, Hodges CS, Alfenas AC. 2018. Canker disease caused by *Chrysosporthe doradensis* and *C. cubensis* on *Eucalyptus* sp. and *Tibouchina* spp. in Brazil. *Tropical Plant Pathol.* 43:314–22
123. Soewarto J, Giblin FR, Carnegie AJ. 2019. *Austropuccinia psidii* (myrtle rust) Global Host List. Version 4. Canberra, ACT: Australian Network for Plant Conservation. https://www.anpc.asn.au/wp-content/uploads/2019/09/MyrtleRust_GlobalHostList_final_4.xlsx
124. Stewart JE, Ross-Davis AL, Graça RN, Alfenas AC, Peever TL, et al. 2018. Genetic diversity of the myrtle rust pathogen (*Austropuccinia psidii*) in the Americas and Hawaii: global implications for invasive threat assessments. *For. Pathol.* 48:e12378
125. Suzuki H, Marincowitz S, Rodas CA, Wingfield BD, Wingfield MJ. 2023. First report of two *Chrysosporthe* species, *Chrysosporthe doradensis* and *Chrysosporthe colombiana* sp. nov. from *Henriettea seemannii* pathogenic to *Eucalyptus* in Colombia. *Mycol. Prog.* 22:44
126. Suzuki H, Marincowitz S, Roux J, Paap T, Wingfield BD, Wingfield MJ. 2024. A new genus and species of Cryphonectriaceae causing stem cankers on plantation eucalypts in South Africa. *Plant Pathol.* 73:1180–93
127. Suzuki H, Marincowitz S, Roux J, Wingfield BD, Wingfield MJ. 2023. First report of canker caused by *Chrysosporthe austroafricana* on the plantation-grown eucalypt *Corymbia henryi* in South Africa. *Forestry* 96:509–17
128. Suzuki H, Marincowitz S, Wingfield BD, Wingfield MJ. 2022. Genetic diversity and population structure of *Chrysosporthe deuterocubensis* isolates from *Melastoma* and *Eucalyptus* in Malaysia and Indonesia. *For. Pathol.* 52:e12762
129. Swart WJ, Conradie E, Wingfield MJ. 1991. *Cryphonectria cubensis*, a potential pathogen of *Psidium guajava* in South Africa. *Eur. J. For. Pathol.* 21:424–29
130. Tarigan M, Wingfield MJ, Jami F, Marpaung YMAN, Durán A, Pham NQ. 2024. Pathogenicity of *Chrysosporthe deuterocubensis* on eucalypts in Indonesia. *South. For. J. For. Sci.* 86:42–52
131. Tian L, Xu J, Zhao D, Qiu H, Yang H, Qin C. 2020. New records of *Celoportha guangdongensis* and *Cytospora rhizophorae* on mangrove apple in China. *Biodivers. Data J.* 8:e55251
132. Ulloa Ulloa C, Almeda F, Goldenberg R, Kadereit G, Michelangeli FA, et al. 2022. Melastomataceae: global diversity, distribution, and endemism. In *Systematics, Evolution, and Ecology of Melastomataceae*, ed. R Goldenberg, FA Michelangeli, F Almeda, pp. 3–28. Springer
133. van der Merwe NA, Gryzenhout M, Steenkamp ET, Wingfield BD, Wingfield MJ. 2010. Multigene phylogenetic and population differentiation data confirm the existence of a cryptic species within *Chrysosporthe cubensis*. *Fungal Biol.* 114:966–79
134. van der Merwe NA, Myburg H, Wingfield BD, Rodas C, Wingfield MJ. 2001. Identification of *Cryphonectria cubensis* from Colombia based on rDNA sequence data. *South Afr. J. Sci.* 97:295–96
135. van der Merwe NA, Phakalatsane T, Wilken PM. 2023. The unique homothallic mating-type loci of the fungal tree pathogens *Chrysosporthe syzygiicola* and *Chrysosporthe zambiensis* from Africa. *Genes* 14:1158
136. van der Merwe NA, Steenkamp ET, Rodas C, Wingfield BD, Wingfield MJ. 2013. Host switching between native and non-native trees in a population of the canker pathogen *Chrysosporthe cubensis* from Colombia. *Plant Pathol.* 62:642–48
137. van der Merwe NA, Wingfield BD, Wingfield MJ. 2003. Primers for the amplification of sequence-characterized loci in *Cryphonectria cubensis* populations. *Mol. Ecol. Notes* 3:494–97
138. van der Nest MA, Chávez R, De Vos L, Duong TA, Gil-Durán C, et al. 2021. IMA genome - F14. *IMA Fungus* 12:5
139. van Heerden SW, Amerson HV, Preisig O, Wingfield BD, Wingfield MJ. 2005. Relative pathogenicity of *Cryphonectria cubensis* on *Eucalyptus* clones differing in their resistance to *C. cubensis*. *Plant Dis.* 89:659–62

140. van Heerden SW, Geletka LM, Preisig O, Nuss DL, Wingfield BD, Wingfield MJ. 2001. Characterization of South African *Cryphonectria cubensis* isolates infected with a *C. parasitica* hypovirus. *Phytopathology* 91:628–32
141. van Heerden SW, Wingfield MJ. 2001. Genetic diversity of *Cryphonectria cubensis* isolates in South Africa. *Mycol. Res.* 105:94–99
142. van Zyl LM, Wingfield MJ. 1999. Wound response of *Eucalyptus* clones after inoculation with *Cryphonectria cubensis*. *Eur. J. For. Pathol.* 29:161–67
143. van Zyl LM, Wingfield MJ, Alfenas AC, Crous PW. 1998. Population diversity among Brazilian isolates of *Cryphonectria cubensis*. *For. Ecol. Manag.* 112:41–47
144. Venter M, Myburg H, Wingfield BD, Coutinho TA, Wingfield MJ. 2002. A new species of *Cryphonectria* from South Africa and Australia, pathogenic to *Eucalyptus*. *Sydowia* 54:98–117
145. Vermeulen M, Gryzenhout M, Wingfield MJ, Roux J. 2011. New records of the Cryphonectriaceae from southern Africa including *Latruncellus aurorae* gen. sp. nov. *Mycologia* 103:554–69
146. Vermeulen M, Gryzenhout M, Wingfield MJ, Roux J. 2013. Population structure of *Chrysosporthe austroafricana* in southern Africa determined using vegetative compatibility groups (VCGs). *For. Pathol.* 43:124–31
147. Vermeulen M, Gryzenhout M, Wingfield MJ, Roux J. 2013. Species delineation in the tree pathogen genus *Celoporthe* (Cryphonectriaceae) in southern Africa. *Mycologia* 105:297–311
148. Wang W, Chen SF. 2023. Identification and pathogenicity of *Aurifilum* species (Cryphonectriaceae, Diaporthales) on *Terminalia* species in Southern China. *MycKeys* 98:37–58
149. Wang W, Li GQ, Liu QL, Chen SF. 2020. Cryphonectriaceae on Myrtales in China: phylogeny, host range, and pathogenicity. *Persoonia* 45:101–31
150. Wang W, Liu QL, Li GQ, Liu FF, Chen SF. 2018. Phylogeny and pathogenicity of *Celoporthe* species from plantation *Eucalyptus* in southern China. *Plant Dis.* 102:1915–27
151. Wardlaw TJ. 1999. *Endothia gyrosa* associated with severe stem cankers on plantation grown *Eucalyptus nitens* in Tasmania, Australia. *Eur. J. For. Pathol.* 29:199–208
152. Wingfield BD, Ades PK, Al-Naemi FA, Beirn LA, Bihon W, et al. 2015. Draft genome sequences of *Chrysosporthe austroafricana*, *Diplodia scrobiculata*, *Fusarium nygamai*, *Leptographium lundbergii*, *Limonomyces culmigenus*, *Stagonosporopsis tanacetii*, and *Thielaviopsis punctulata*. *IMA Fungus* 6:233–48
153. Wingfield BD, Barnes I, de Beer ZW, De Vos L, Duong TA, et al. 2015. Draft genome sequences of *Ceratocystis eucalypticola*, *Chrysosporthe cubensis*, *C. deuterocubensis*, *Davidsoniella virescens*, *Fusarium temperatum*, *Graphilbum fragrans*, *Penicillium nordicum*, and *Thielaviopsis musarum*. *IMA Fungus* 6:493–506
154. Wingfield MJ. 2003. 2003 Daniel McAlpine Memorial Lecture. Increasing threat of diseases to exotic plantation forests in the Southern Hemisphere: lessons from *Cryphonectria* canker. *Australas. Plant Pathol.* 32:133–39
155. Wingfield MJ, Brouckhoff EG, Wingfield BD, Slippers B. 2015. Planted forest health: the need for a global strategy. *Science* 349:832–36
156. Wingfield MJ, Rodas C, Myburg H, Venter M, Wright J, Wingfield BD. 2001. *Cryphonectria* canker on *Tibouchina* in Colombia. *For. Pathol.* 31:297–306
157. Wingfield MJ, Swart WJ, Abear BJ. 1989. First record of *Cryphonectria* canker of *Eucalyptus* in South Africa. *Phytophylactica* 21:311–13
158. Yuan ZQ, Mohammed C. 2000. The pathogenicity of isolates of *Endothia gyrosa* to *Eucalyptus nitens* and *E. globulus*. *Australas. Plant Pathol.* 29:29–35