

A critique of “Asexually Produced Cape Honeybee Queens (*Apis mellifera capensis*) Reproduce Sexually” authors: MADELEINE BEEKMAN, MICHAEL H. ALLSOPP, JULIANNE LIM, FRANCES GOUDIE, AND BENJAMIN P. OLDROYD *Journal of Heredity* 2011:102(5):562–566

Reproductive biology of the Cape honeybee: A critique of Beekman et al.

Christian W.W. Pirk¹, H. Michael G. Lattorff², Robin F.A. Moritz^{1,2}, Catherine L. Sole¹, Sarah E. Radloff³, Peter Neumann^{4,5}, H Randall Hepburn⁴, Robin M. Crewe¹

¹ Department of Zoology and Entomology, University of Pretoria, Pretoria 0002, South Africa

² Institut für Biologie, Molekulare Ökologie, Martin-Luther-Universität Halle-Wittenberg, Hoher Weg 4, D-06099 Halle (Saale), Germany.

³ Department of Statistics, Rhodes University, Grahamstown 6140, South Africa.

⁴ Department of Zoology and Entomology, Rhodes University, Grahamstown 6140, South Africa

⁵ Swiss Bee Research Centre, Agroscope Liebefeld-Posieux Research Station ALP, CH-3033 Bern, Switzerland

*A critique of “Asexually Produced Cape Honeybee Queens (*Apis mellifera capensis*) Reproduce Sexually”: Laying workers of the Cape honeybee parthenogenetically produce female offspring whereas queens typically produce males. Beekman et al. confirm this observation, which has repeatedly been reported over the last 100 years including the notion that natural selection should favor asexual reproduction in *A. m. capensis*. They attempt to support their arguments with an exceptionally surprising finding that *A. m. capensis* queens can parthenogenetically produce diploid homozygous queen offspring (homozygous diploid individuals develop into diploid males in the honeybee). Beekman et al. suggest that these homozygous queens are not viable because they did not find any homozygous individuals beyond the third larval instar. Even if this were true, such a lethal trait should be quickly eliminated by natural selection. The identification of sex (both with molecular and morphological markers) is possible but notoriously difficult in honeybees at the early larval stages. Ploidy is however a reliable indicator and we therefore suggest that these “homozygous” larvae found in queen cells are actually drones reared from unfertilized eggs, a phenomenon well known by honeybee queen breeders.*

The Cape honeybee, *Apis mellifera capensis*, shows several unique features in relation to reproduction and pheromonal dominance, which makes it an excellent model system for understanding the evolution of reproductive dominance in social hymenoptera. This is partly due to the fact that workers of the Cape honeybee are able to produce diploid female offspring from unfertilized eggs (Onions 1912; Anderson 1963), a process called thelytoky (Crozier 1975). These female offspring can then develop either into workers or queens (Anderson 1963; Ruttner 1977). The underlying cytological mechanism is well understood and is based on automictic parthenogenesis with a central fusion (Verma and Ruttner 1983) that prevents meiotic recombination (Moritz and Haberl 1994). Even recombination due to crossing-over is heavily reduced (Baudry et al. 2004).

Recently, Beekman et al (2011) reported in this journal that asexually produced Cape honeybee queens reproduce sexually. This is correct, but far from being a novel finding. Over the past **100 years** it has been repeatedly reported that new queens produced in *A. m. capensis* colonies were laying worker offspring (Onions 1912; Anderson 1963; Ruttner 1977; Moritz et al. 1996; Allsopp and Hepburn 1997; Moritz et al. 2011) and these queens behave no differently from queens produced from fertilized eggs (Jack 1917; Hepburn and Guye 1993; Tribe and Allsopp 2001).

Thelytokous parthenogenesis in the Cape honeybee results from the automictic fusion of the two central of the four meiotic products to restore diploidy (Tucker 1958; Verma and Ruttner 1983). Therefore the question in Beekman et al (2011) should not be whether mated queens produce diploid worker offspring from fertilized eggs or reproduce asexually; but, rather whether a mated Cape

honeybee queen of laying worker origin can produce male offspring. To address this question, the authors should have sampled larvae from drone cells rather than worker cells (Beekman et al. 2011). However even such a finding would have had only rather limited novelty since again it has been shown repeatedly over the past 100 years that impaternal female progeny of virgin queens can occur (Onions 1912; Mackensen 1943; Tryasko 1969; Tryasko 1975; Velthuis et al. 1990; Oldroyd et al. 2008). However, the conditional use of sex when producing workers or queens reported from two ant species (Pearcy et al. 2004; Fournier et al. 2005) might also occur in Cape honeybees, but has not been addressed experimentally by Beekman et al (2011). For this purpose they should have sampled both larvae from queen and worker cells. Nevertheless, recently it has been shown that queens fertilize eggs when producing queens and workers (Moritz et al. 2011).

Beekman et al (2011) finally argue that natural selection should favour asexual reproduction in *A. m. capensis*. Indeed this is true for a wide region of the fitness parameter space as had been noted several decades ago (Moritz 1989; Greeff 1996a; Greeff 1996b). However, Beekman et al (2011) attempt to support that contention with the results of Jordan et al (2008) who claimed to have found parthenogenetically produced homozygous female offspring. The evidence for that exceptionally surprising finding (homozygous diploid individuals develop into diploid males in the honeybee that are cannibalized by workers, Mackensen 1951) was however weak. Given the fundamental discrepancy to the basic knowledge on both sex determination and the cytogenetics of thelytokous parthenogenetics in honeybees, the data should have been supported by a cytogenetic confirmation of previous studies on the thelytokous parthenogenetic

mechanism in the Cape honeybee. Since central fusion prevents any recombination of chromosomes, the identical genotype of the mother is restored apart from rare crossing over events (Moritz and Haberl 1994; Neumann and Moritz 2002; Baudry et al. 2004). Unless a robust analysis confirms that the sampled larvae were female, it seems more parsimonious to explain the observed homozygosity by Jordan et al (2008) as male eggs being laid accidentally in queen cells which is a common phenomenon in apicultural queen rearing (Fell and Morse 1984).

We feel Beekman et al (2011) do not produce an evolutionary plausible explanation by stating that they could not find any homozygous individuals beyond the third larval instar, because they are not viable. Clearly, any lethal trait is exposed to strong selection and if this were the result of the thelytoky trait in *A. m. capensis* we would simply not be able to use this honeybee as a model for studying the evolution of social behaviour today.

References

- Allsopp MH, Hepburn HR (1997) Swarming, supersedure and mating system of a natural population of honey bees (*Apis mellifera capensis*). *Journal of Apicultural Research* 1:41-48
- Anderson RH (1963) The laying worker in the Cape honeybee *Apis mellifera capensis* *Journal of Apicultural Research* 2:85-92
- Baudry E, Kryger P, Allsopp MH, Koeniger N, Vautrin D, Mougél F, Cornuet J-M, Solognac M (2004) Whole-Genome Scan in Thelytokous-Laying Workers of the Cape Honeybee (*Apis mellifera capensis*): Central Fusion, reduced Recombination Rates and Centromere Mapping Using Half-Tetrad Analysis *Genetics* 167:243-252
- Beekman M, Allsopp MH, Lim J, Goudie F, Oldroyd BP (2011) Asexually Produced Cape Honeybee Queens (*Apis mellifera capensis*) Reproduce Sexually. *Journal of Heredity* 102:562-566
- Crozier RH (1975) *Animal Cytogenetics*. Bornträger Berlin, Germany

- Fell R, Morse R (1984) Emergency queen cell production in the honey bee colony. *Insectes Sociaux* 31:221-237
- Fournier D, Estoup A, Orivel J, Foucaud J, Jourdan H, Le Breton J, Keller L (2005) Clonal reproduction by males and females in the little fire ant. *Nature* 435:1230-1234
- Greeff JM (1996a) Effects of thelytokous worker reproduction on kin-selection and conflict in the Cape honeybee *Apis mellifera capensis* *Philosophical Transactions of the Royal Society of London B Biological-Sciences* 351:617-625
- Greeff JM (1996b) Thelytokous versus arrhenotokous worker reproduction in the Cape honeybee and other eusocial Hymenoptera. *Hereditas* 124:99-103
- Hepburn HR, Guye SG (1993) An annotated bibliography of the Cape honeybee, *Apis mellifera capensis* Eschscholtz (Hymenoptera: Apidae). *African Entomology* 1:235-252
- Jack RW (1917) XXI. Parthenogenesis amongst the Workers of the Cape Honey-Bee: Mr. G. W. Onions' Experiments. *Transactions of the Royal Entomological Society of London* 64:396-403
- Jordan LA, Allsopp MH, Oldroyd BP, Wossler TC, Beekman M (2008) Cheating honeybee workers produce royal offspring. *Proceedings of the Royal Society B: Biological Sciences* 275:345-351
- Mackensen O (1943) The Occurrence of Parthenogenetic Females in Some Strains of Honeybees. *Journal of Economic Entomology* 36:465-467
- Mackensen O (1951) Viability and Sex determination in the honey bee (*Apis mellifera* L.) *Genetics* 36:500-509
- Moritz R, Lattorff H, Crous K, Hepburn R (2011) Social parasitism of queens and workers in the Cape honeybee (*Apis mellifera capensis*). *Behavioral Ecology and Sociobiology* 65:735-740
- Moritz RFA (1989) Colony Level and within Colony Level Selection in Honeybees: A Two Allele Population Model for *Apis mellifera capensis*. *Behavioral Ecology and Sociobiology* 25:437-444
- Moritz RFA, Haberl M (1994) Lack of meiotic recombination in thelytokous parthenogenesis of laying workers of *Apis mellifera capensis* (the Cape honeybee). *Heredity* 73:98-102
- Moritz RFA, Kryger P, Allsopp MH (1996) Competition for royalty in bees. *Nature* 384 31
- Neumann P, Moritz RFA (2002) The Cape honeybee phenomenon: the sympatric evolution of a social parasite in real time? *Behavioral Ecology and Sociobiology* 52:271-281
- Oldroyd BP, Allsopp MH, Gloag RS, Lim J, Jordan LA, Beekman M (2008) Thelytokous parthenogenesis in unmated queen honey bees (*Apis mellifera capensis*): central fusion and high recombination rates. *Genetics* 180:359-366
- Onions GW (1912) South African 'fertile worker bees'. *Agricultural Journal of the Union of South Africa* 3:720-728
- Pearcy M, Aron S, Doums C, Keller L (2004) Conditional Use of Sex and Parthenogenesis for Worker and Queen Production in Ants. *Science* 306:1780-1783

- Ruttner F (1977) The problem of Cape Bee (*Apis mellifera capensis* Escholtz) - Parthenogenesis - Size of Population - Evolution. *Apidologie* 8:281-294
- Tribe GD, Allsopp MH (2001) Honeybee Reproduction. In: Johannsmeier MF (ed) *Beekeeping in South Africa*. ARC-Plant Protection Research Institute, Pretoria, South Africa, pp 40-46
- Tryasko VV (1969) Study of spontaneous female parthenogenesis in honeybees *Proceedings of the Apidmondia Congress München*:599
- Tryasko VV (1975) Cytologic mechanism of female parthenogenesis in honeybees. In: *Proceedings of the 25th Apimondia Congress*, vol 25, Grenoble, pp 318
- Tucker KW (1958) Automictic parthenogenesis in the honey bee. *Genetics* 43:299-316
- Velthuis HHW, Ruttner F, Crewe RM, Engels W (1990) Differentiation in Reproductive Physiology and Behaviour during the Development of laying worker honeybees. In: *Social Insects: an evolutionary approach to castes and reproduction*. Springer Verlag, Berlin, pp 231-243
- Verma S, Ruttner F (1983) Cytological analysis of the thelytokous parthenogenesis in the Cape honeybee (*Apis mellifera capensis* Escholtz). *Apidologie* 14:41-57