


Supplemental online material: The role of co-creation tools in the evaluation and redesign of healthcare services

Supplement 1: Citation network



(created with VOSviewer)

Supplement 2: Main path



(created with Pajek)

Supplement 3: Main path documents

Identification	Reference
R1	Rethans JJ, Drop R, Sturmans F, van der Vleuten C. A method for introducing standardized (simulated) patients into general practice consultations. <i>Br J Gen Pract.</i> 1991 Mar;41(344):94-6. PMID: 2031766; PMCID: PMC1371619.
R2	Carney PA, Dietrich AJ, Freeman DH Jr, Mott LA. The periodic health examination provided to asymptomatic older women: an assessment using standardized patients. <i>Ann Intern Med.</i> 1993 Jul 15;119(2):129-35. doi: 10.7326/0003-4819-119-2-199307150-00007. PMID: 8512162.
R3	McLeod PJ, Tamblyn RM, Gayton D, Grad R, Snell L, Berkson L, Abrahamowicz M. Use of standardized patients to assess between-physician variations in resource utilization. <i>JAMA.</i> 1997 Oct 8;278(14):1164-8. PMID: 9326476.
R4	Woodward CA, McConvey GA, Neufeld V, Norman GR, Walsh A. Measurement of physician performance by standardized patients. Refining techniques for undetected entry in physicians' offices. <i>Med Care.</i> 1985 Aug;23(8):1019-27. doi: 10.1097/00005650-198508000-00009. PMID: 4021578.
R5	Peabody JW, Luck J, Glassman P, Dresselhaus TR, Lee M. Comparison of vignettes, standardized patients, and chart abstraction: a prospective validation study of 3 methods for measuring quality. <i>JAMA.</i> 2000 Apr 5;283(13):1715-22. doi: 10.1001/jama.283.13.1715. PMID: 10755498.
R6	Luck J, Peabody JW, Dresselhaus TR, Lee M, Glassman P. How well does chart abstraction measure quality? A prospective comparison of standardized patients

	with the medical record. Am J Med. 2000 Jun 1;108(8):642-9. doi: 10.1016/s0002-9343(00)00363-6. PMID: 10856412.
R7	Fiscella K, Franks P, Srinivasan M, Kravitz RL, Epstein R. Ratings of physician communication by real and standardized patients. Ann Fam Med. 2007 MarApr;5(2):151-8. doi: 10.1370/afm.643. PMID: 17389540; PMCID: PMC1838677.
R8	Zabar S, Ark T, Gillespie C, Hsieh A, Kalet A, Kachur E, Manko J, Regan L. Can unannounced standardized patients assess professionalism and communication skills in the emergency department? Acad Emerg Med. 2009 Sep;16(9):915-8. doi: 10.1111/j.1553-2712.2009.00510.x. Epub 2009 Aug 10. PMID: 19673703.
R9	Siminoff LA, Rogers HL, Waller AC, Harris-Haywood S, Epstein RM, Carrio FB, Gliva-McConvey G, Longo DR. The advantages and challenges of unannounced standardized patient methodology to assess healthcare communication. Patient Educ Couns. 2011 Mar;82(3):318-24. doi: 10.1016/j.pec.2011.01.021. PMID: 21316182; PMCID: PMC3064426.
R10	Sylvia S, Xue H, Zhou C, Shi Y, Yi H, Zhou H, Rozelle S, Pai M, Das J. Tuberculosis detection and the challenges of integrated care in rural China: A cross-sectional standardized patient study. PLoS Med. 2017 Oct 17;14(10):e1002405. doi: 10.1371/journal.pmed.1002405. PMID: 29040263; PMCID: PMC5644979.
R11	Kwan A, Daniels B, Bergkvist S. Use of standardised patients for healthcare quality research in low- and middle-income countries. BMJ Global Health. 2019;4:e001669. doi: 10.1136/bmjgh-2019-001669. PMID: 31565413; PMCID: PMC6747906.

R12	Sulis G, Daniels B, Kwan A. Antibiotic overuse in the primary health care setting: a secondary data analysis of standardised patient studies from India, China and Kenya. <i>BMJ Global Health</i> . 2020;5:e003393.
R13	Xu DR, Hu M, He W, Liao J, Cai Y, Sylvia S, Hanson K, Chen Y, Pan J, Zhou Z, Zhang N, Tang C, Wang X, Rozelle S, He H, Wang H, Chan G, Melipillán ER, Zhou W, Gong W. Assessing the quality of primary healthcare in seven Chinese provinces with unannounced standardised patients: protocol of a cross-sectional survey. <i>BMJ Open</i> . 2019;9:e023997. doi: 10.1136/bmjopen-2018-023997. PMID: 30765399; PMCID: PMC6398795.
R14	Xu D, Pan J, Dai X, Hu M, Cai Y, He H, Zhang Y, Liao J, Chen Y, Gong W, Zhou Z, Zhang N, Wang X, Chan KCG, Ying X, Cai Y, Wang R, Xue Q, Yip CW. Comparing quality of primary healthcare between public and private providers in China: study protocol of a cross-sectional study using unannounced standardised patients in seven provinces of China. <i>BMJ Open</i> . 2021 Jan 12;11(1):e040792. doi: 10.1136/bmjopen-2020-040792. PMID: 33436467; PMCID: PMC7805374.
R15	Kwan A, Daniels B, Saria V, Satyanarayana S, Subbaraman R, McDowell A, Bergkvist S, Das RK, Das V, Das J, Pai M. Variations in the quality of tuberculosis care in urban India: A cross-sectional, standardized patient study in two cities. <i>PLoS Med</i> . 2018 Sep 25;15(9):e1002653. doi: 10.1371/journal.pmed.1002653. PMID: 30252849; PMCID: PMC6155454.
R16	Wiseman V, Lagarde M, Kovacs R, Wulandari LPL, Powell-Jackson T, King J, Goodman C, Hanson K, Miller R, Xu D, Liverani M, Yeung S, Hompashe D, Khan M, Burger R, Christian CS, Blaauw D. Using unannounced standardised patients to obtain data on quality of care in low-income and middle-income countries:

	key challenges and opportunities. <i>BMJ Glob Health</i> . 2019 Sep 12;4(5):e001908. doi: 10.1136/bmjgh-2019-001908. PMID: 31565422; PMCID: PMC6747897
R17	Rutter PM, Horsley E, Brown DT. Evaluation of community pharmacists' recommendations to standardized patient scenarios. <i>Ann Pharmacother</i> . 2004 Jun;38(6):1080-5. doi: 10.1345/aph.1D519. Epub 2004 Apr 27. PMID: 15113984.
R18	Watson M, Norris P, Granas A. A systematic review of the use of simulated patients and pharmacy practice research. <i>International Journal of Pharmacy Practice</i> . 2006 14: 83-93. doi: https://doi.org/10.1211/ijpp.14.2.0002 .
R19	Watson MC, Cleland JA, Bond CM. Simulated patient visits with immediate feedback to improve the supply of over-the-counter medicines: a feasibility study. <i>Fam Pract</i> . 2009 Dec;26(6):532-42. doi: 10.1093/fampra/cmp061. Epub 2009 Oct 14. PMID: 19828574.
R20	Xu T, de Almeida Neto AC, Moles RJ. A systematic review of simulated-patient methods used in community pharmacy to assess the provision of non-prescription medicines. <i>Int J Pharm Pract</i> . 2012 Oct;20(5):307-19. doi: 10.1111/j.2042-7174.2012.00201.x. Epub 2012 Mar 13. PMID: 22953770.
R21	Satyanarayana S, Kwan A, Daniels B, Subbaraman R, McDowell A, Bergkvist S, Das RK, Das V, Das J, Pai M. Use of standardised patients to assess antibiotic dispensing for tuberculosis by pharmacies in urban India: a cross-sectional study. <i>Lancet Infect Dis</i> . 2016 Nov;16(11):1261-1268. doi: 10.1016/S1473-3099(16)30215-8. Epub 2016 Aug 25. PMID: 27568359; PMCID: PMC5067371.
R22	Franz CE, Epstein R, Miller KN, Brown A, Song J, Feldman M, Franks P, Kelly-Reif S, Kravitz RL. Caught in the act? Prevalence, predictors, and consequences of physician detection of unannounced standardized patients. <i>Health Serv Res</i> .

	2006 Dec;41(6):2290-302. doi: 10.1111/j.1475-6773.2006.00560.x. PMID: 17116121; PMCID: PMC1955318.
R23	Daniels B, Dolinger A, Bedoya G. Use of standardised patients to assess quality of healthcare in Nairobi, Kenya: a pilot, cross-sectional study with international comparisons. <i>BMJ Global Health</i> . 2017;2:e000333.
R24	Das J, Kwan A, Daniels B, Satyanarayana S, Subbaraman R, Bergkvist S, Das RK, Das V, Pai M. Use of standardised patients to assess quality of tuberculosis care: a pilot, cross-sectional study. <i>Lancet Infect Dis</i> . 2015 Nov;15(11):1305-13. PMID: 26268690; PMCID: PMC4633317.
R25	Rosen GM, Phillips WR. A cautionary lesson from simulated patients. <i>J Am Acad Psychiatry Law</i> . 2004;32(2):132-3. PMID: 15281413.