

## **Sexual reproduction in populations of *Austropuccinia psidii***

Alistair R. McTaggart<sup>1,2\*</sup>, Esna du Plessis<sup>1\*</sup>, Jolanda Roux<sup>1</sup>, Irene Barnes<sup>3</sup>, Stuart Fraser<sup>1,4</sup>,  
Ginna M. Granados<sup>3</sup>, Wellcome W. H. Ho<sup>5</sup>, Louise S. Shuey<sup>3</sup>, André Drenth<sup>2</sup>

\* The first two authors contributed equally to the study

<sup>1</sup>Department of Plant and Soil Sciences, Tree Protection Co-operative Programme (TPCP),  
Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa.

<sup>2</sup>Queensland Alliance for Agriculture and Food Innovation, The University of Queensland,  
Ecosciences Precinct, GPO Box 267, Brisbane, Queensland 4001, Australia.

<sup>3</sup>Department of Biochemistry, Genetics and Microbiology, TPCP, FABI, University of Pretoria,  
South Africa.

<sup>4</sup>Forest Protection, Scion, Rotorua, New Zealand.

<sup>5</sup>Plant Health & Environment Laboratory, Diagnostic and Surveillance Services, Ministry for  
Primary Industries P.O. Box 2095, Auckland, 1140, New Zealand.

email: a.mctaggart@uq.edu.au

## **Abstract**

*Austropuccinia psidii* is a rust fungus that has expanded its known geographic distribution and host range on Myrtaceae. Invasions by rust fungi are often caused by asexual urediniospores that give rise to populations with low genotypic diversity. Recently it was shown that basidiospores, the gametic spores of *A. psidii*, were able to infect species of Myrtaceae under controlled conditions. The present study tested the hypothesis that sexual reproduction occurs through infection of Myrtaceae by basidiospores of *A. psidii* in recently invasive populations from New Zealand and South Africa. We provided three lines of evidence to test this hypothesis, i) presence of a sexual stage, ii) high genotypic diversity within an invasive population and iii) no genetic linkage between microsatellite markers in multilocus genotypes. Our results provide evidence that invasions of *A. psidii* are caused by both urediniospores that spread clonal genotypes, and teliospores that produce recombinant basidiospores, which infect Myrtaceae. We reject the hypothesis that field infections of *A. psidii* are only caused by asexual urediniospores, and support that sexual reproduction occurs in invasive populations and may accelerate adaptation to environmental change.

## **Running Title**

Recombinant populations of *A. psidii*

**Keywords:** Biological invasions, clonal fungi, life cycle, mixed mating system, myrtle rust, Pucciniales

## Introduction

The rust fungus *Austropuccinia psidii* (syn. *Puccinia psidii*, Sphaerophragmiaceae, Pucciniales) causes disease on species of Myrtaceae. Winter (1884) described *A. psidii* on a species of *Psidium* from Brazil, and it was soon found on other native genera in South America, such as *Eugenia* and *Plinia* (Rangel 1916). It was a new encounter pathogen on introduced species of *Eucalyptus* in Brazil (Joffily 1944), which foreshadowed its potential to cause disease on a wider host range of the Myrtaceae. *Austropuccinia psidii* spread from its hypothesized native range in Central and South America (Coutinho et al. 1998) and is currently present in at least 27 countries (Carnegie and Pegg 2018).

The life cycle stages of *A. psidii* include urediniospores, a dikaryotic clonal spore stage, and diploid teliospores that produce basidiospores through meiosis. Mature basidiospores contain two nuclei, but it is unknown whether these are hetero- or homokaryotic (Morin et al. 2014; McTaggart et al. 2018). Two different life cycles of *A. psidii* have been hypothesized. Coutinho et al. (1998) suggested completion of its life cycle on a single host without production of spermogonia and aecia. Ruiz et al. (1989) postulated that spermogonia and aecia were formed on an alternate, but unknown host, similar to other heteroecious rust fungi.

The role of basidiospores has been investigated to resolve the life cycle of *A. psidii*. Figueiredo et al. (1984) and Figueiredo (2001) inoculated *Syzygium jambos* and a species of *Eucalyptus* with basidiospores, which infected and produced uredinia on both hosts. However, Morin et al. (2014) observed that basidiospores did not penetrate the cuticle of inoculated *Agonis flexuosa*, and despite production of uredinia from inoculations with basidiospores, there were no signs of recombination measured by genotypic diversity. Morin et al. (2014) suggested urediniospores had contaminated inocula in their own, and earlier studies. The lack of evidence for sexual recombination in populations of *A. psidii* sampled from cultivated hosts and invasion events led to the conclusion that it reproduces clonally (Graça et al. 2013; Machado et al. 2015). The observed genotypic diversity in populations of *A. psidii* was treated

as the product of mutation rather than sexual reproduction (Graça et al. 2013; Machado et al. 2015).

McTaggart et al. (2018) queried the assumption that *A. psidii* only reproduces asexually and provided evidence for the alternative hypothesis that genotypic diversity is a product of recombination and infection of plants by basidiospores. They filtered urediniospores from their inocula and used microsatellite markers to confirm pustules that developed after inoculations with basidiospores on *S. jambos* were the product of sexual reproduction. These experiments showed *A. psidii* completes the asexual and sexual parts of its life cycle on a single host (autoecious) and that meiotic recombination produces basidiospores that are able to infect species of Myrtaceae under controlled conditions (McTaggart et al. 2018).

Fungal populations acquire genetic diversity in different ways (Milgroom 1996). A population that reproduces sexually may produce sexual structures, have increased genotypic diversity and many possible recombinant genotypes, and different regions of the genome may have different histories of descent (Milgroom 1996). Alleles are randomly distributed among individuals after sexual reproduction, which can be measured in diploid and dikaryotic organisms using the inbreeding coefficient (De Meeûs et al. 2006). Clonal populations will have low genotypic diversity, show linkage disequilibrium between markers, and be mostly either homozygous or heterozygous at individual loci (De Meeûs et al. 2006; Grunwald et al. 2017; Milgroom 1996). Clonal populations from a single invasion event may also have low allelic diversity when a single genotype infects, colonizes and spreads (Gladieux et al. 2015; McDonald and Linde 2002).

The recent incursions of *A. psidii* in South Africa (Roux et al. 2013) and New Zealand (Ho et al. 2019) provided us with an opportunity to test the hypothesis that sexual reproduction of this rust occurs in the field as part of, or immediately following, an invasion event. Roux et al. (2016) reported one genotype was present in South Africa and noted that telia, the

precursors to sexual reproduction, were abundant. In a separate invasion event of *A. psidii* in New Zealand, several genotypes were identified (du Plessis et al. 2019) and telia were present (Ho et al. 2019). We specifically sought to test if (i) basidiospores of *A. psidii* play a role in the epidemiology of invasive populations, and (ii) recombination gives rise to a random association of genetic markers in the invading population. Insight into the reproductive biology of invasive species such as rusts, and understanding their disease cycle and reproductive biology, including the generation of genetic diversity through recombination, are important in efforts to either eradicate, contain or manage these invasions.

## **Materials and methods**

### *Sampling and DNA extraction*

We used an *ad hoc* sampling strategy for specimens collected in South Africa as part of an ongoing study to monitor the distribution of *A. psidii*. We sampled the first five leaf specimens of *A. psidii* from the disease incursion in New Zealand, sent to South Africa under import permit P0079592, as described in the study of du Plessis et al. (2019). The presence or absence of telia was recorded by visual inspection for all specimens from New Zealand and South Africa. All specimens used in the present study are listed in Table 1.

Two to four discrete pustules were sampled from single leaves of each specimen to detect genotypic diversity expected from infections caused by basidiospores. If four or more of the five microsatellite loci amplified, the pustule was included in the analyses of genotypic diversity and recombination. These analyses included 35 (out of 62) pustules of *A. psidii* from South Africa and nine (out of 22) pustules from New Zealand (Table 1).

DNA was extracted from single pustules using either the Genra Puregene Yeast/Bacteria Kit (Qiagen, Germany) or the Ultraclean® Microbial DNA Isolation Kit (MoBio Laboratories, USA). The protocols supplied by the manufacturer were followed, with the exception that DNA was eluted in a final volume of 15 µL.

### *Selection, amplification and analysis of microsatellite loci*

We tested whether sexual reproduction occurred in populations of *A. psidii* from South Africa and New Zealand with five microsatellite markers (PpSSR012, PpSSR014, PpSSR018, PpSSR102, PpSSR161) originally developed by Zhong et al. (2008), and modified by Graça et al. (2013). The markers were labeled with FAM™, NED™, PET® or VIC® fluorescent dye. These markers had two alleles each in the invasive population of *A. psidii* in South Africa (Roux et al. 2016), and had two alleles each in the population from New Zealand, with the exception of PpSSR102 that had one allele (du Plessis et al. 2019).

PCR mixtures included 1× PCR FastStart™ Taq Buffer with MgCl<sub>2</sub> (Sigma-Aldrich, USA), 200 μM dNTPs, 0.1 μM fluorescent primers, 1 unit FastStart™ Taq DNA polymerase and DNA template in 12.5 μL reaction volumes. PCR products were amplified with the following conditions: one cycle at 95°C for 10 min, followed by 10 cycles at 95°C for 30s, 52–56°C (depending on the primer pair) for 45s, 72°C for 60s, 30 cycles on an auto-delta step at 95°C for 30s, 52–56°C (depending on the primer pair) for 45s plus 5s added after each cycle, and 72°C for 60s, one cycle at 60°C for 30 min. Primers PpSSR 012, PpSSR 014, PpSSR 018, PpSSR 161 were amplified at 56°C, and PpSSR 102 at 52°C. An ABI® Applied Biosystems Veriti 96 Well Thermal Cycler (ThermoFisher Scientific, USA) was used for PCRs.

PCR fragments were analysed on an ABI Applied Biosystems PRISM® 3500xl Autosequencer (ThermoFisher Scientific, USA) at the Sequencing Facility of the Faculty of Natural and Agricultural Sciences, University of Pretoria. Samples were run with GeneScan™ 500 LIZ® Size Standard (ThermoFisher Scientific, USA) and scored with GeneMapper® Software 5 (Thermo Fischer Scientific, USA).

**Table 1.** Collection information and allele sizes of *Austropuccinia psidii* at five microsatellite loci used in this study. NA is used if microsatellite locus was not amplified.

Herbarium number (PREM <sup>a</sup> )	Origin	Host	Telia	Allele sizes				
				PpSSR 12	PpSSR 14	PpSSR 18	PpSSR 102	PpSSR 161
62050	LIM <sup>b</sup> , RSA <sup>c</sup>	<i>Eugenia natalitia</i>	Yes	237, 237	205, 205	165, 174	140, 140	288, 288
62051	LIM, RSA	<i>E. natalitia</i>	Yes	NA	205, 205	165, 174	140, 142	270, 288
62052	WC <sup>d</sup> , RSA	<i>Myrtus communis</i>	No	233, 237	205, 213	165, 174	140, 142	270, 288
62053	KZN <sup>e</sup> , RSA	<i>E. erythrophylla</i>	Yes	NA	213, 213	174, 174	140, 142	288, 288
62054	KZN, RSA	<i>E. verdoorniae</i>	Yes	233, 233	213, 213	NA	140, 140	270, 288
62055	KZN, RSA	<i>E. umtamvunensis</i>	Yes	237, 237	205, 205	165, 174	142, 142	288, 288
62056	KZN, RSA	<i>E. natalitia</i>	Yes	237, 237	205, 213	165, 165	140, 142	288, 288
62057	KZN, RSA	<i>E. natalitia</i>	Yes	233, 237	205, 213	NA	140, 140	270, 288
62058	KZN, RSA	<i>E. natalitia</i>	No	233, 237	205, 213	165, 174	140, 142	270, 288
62059	KZN, RSA	<i>E. simii</i>	No	233, 237	205, 213	165, 174	142, 142	NA
62060	KZN, RSA	<i>E. simii</i>	No	233, 237	205, 213	165, 174	NA	270, 288
62061	KZN, RSA	<i>E. erythrophylla</i>	Yes	237, 237	205, 213	165, 165	140, 142	270, 270
62062	KZN, RSA	<i>E. verdoorniae</i>	Yes	233, 237	205, 213	165, 174	140, 142	270, 288
62063	KZN, RSA	<i>E. umtamvunensis</i>	Yes	237, 237	205, 205	174, 174	NA	270, 270
62064	KZN, RSA	<i>E. erythrophylla</i>	No	237, 237	205, 213	165, 165	140, 140	270, 288
62065	KZN, RSA	<i>E. verdoorniae</i>	Yes	233, 237	205, 205	172, 172	142, 142	270, 288
62066	KZN, RSA	<i>E. umtamvunensis</i>	No	NA	213, 213	174, 174	140, 140	270, 270
62067	KZN, RSA	<i>E. umtamvunensis</i>	No	237, 237	NA	174, 174	142, 142	288, 288
62068	MPU <sup>f</sup> , RSA	<i>Melaleuca viminalis</i>	Yes	233, 233	213, 213	165, 174	140, 142	288, 288
62069	KZN, RSA	<i>E. capensis</i>	Yes	233, 237	213, 213	165, 174	142, 142	270, 270
62070	LIM, RSA	<i>E. natalitia</i>	No	233, 233	205, 213	165, 165	140, 142	288, 288
62071	KZN, RSA	<i>My. communis</i>	No	233, 237	205, 213	165, 174	140, 142	270, 288
62072	MPU, RSA	<i>Mel. viminalis</i>	Yes	233, 237	213, 213	174, 174	140, 142	288, 288
62073	KZN, RSA	<i>E. capensis</i>	Yes	233, 233	205, 205	165, 174	142, 142	270, 288
62074	EC <sup>g</sup> , RSA	<i>My. communis</i>	No	NA	205, 213	165, 174	140, 142	270, 288
62075	KZN, RSA	<i>E. capensis</i>	No	237, 237	205, 213	165, 174	140, 142	270, 288
62076	KZN, RSA	<i>E. capensis</i>	No	233, 237	213, 213	165, 174	142, 142	270, 288
62077	KZN, RSA	<i>E. capensis</i>	No	233, 233	213, 213	174, 174	140, 140	NA
62078	KZN, RSA	<i>E. capensis</i>	No	233, 237	205, 213	NA	140, 142	270, 288
62079	KZN, RSA	<i>E. verdoorniae</i>	No	233, 233	205, 213	165, 172	140, 142	NA
62080	KZN, RSA	<i>E. verdoorniae</i>	No	233, 237	205, 213	165, 174	140, 142	270, 288
62081	KZN, RSA	<i>E. erythrophylla</i>	No	233, 233	205, 213	165, 174	140, 142	270, 288
62082	MPU, RSA	<i>Mel. viminalis</i>	No	233, 237	205, 213	NA	140, 142	270, 270
62083	KZN, RSA	<i>E. capensis</i>	No	233, 237	205, 213	174, 174	140, 142	270, 288
62084	KZN, RSA	<i>E. capensis</i>	Yes	233, 237	205, 213	165, 174	140, 142	NA
62096 – 1	RI <sup>h</sup> , NZ <sup>i</sup>	<i>Metrosideros kermadecensis</i>	Yes	230, 236	207, 211	172, 172	140, 140	276, 276
62096 – 2	RI, NZ	<i>Met. kermadecensis</i>	Yes	230, 236	211, 211	170, 172	NA	290, 290
62099 – 1	NI <sup>j</sup> , NZ	<i>Met. excelsa</i>	Yes	230, 236	211, 211	170, 172	NA	276, 290
62099 – 2	NI, NZ	<i>Met. excelsa</i>	Yes	230, 236	207, 207	170, 170	140, 140	NA
62099 – 3	NI, NZ	<i>Met. excelsa</i>	Yes	230, 236	207, 207	170, 172	140, 140	276, 290
62099 – 4	NI, NZ	<i>Met. excelsa</i>	Yes	236, 236	207, 211	170, 172	140, 140	NA
62102	NI, NZ	<i>Lophomyrtus bullata</i>	Yes	236, 236	207, 211	170, 170	140, 140	NA
62103	NI, NZ	<i>Lophomyrtus bullata</i>	Yes	236, 236	207, 211	170, 172	NA	276, 290
62104	NI, NZ	<i>Lophomyrtus bullata</i>	Yes	230, 230	207, 211	170, 172	140, 140	NA

<sup>a</sup>Herbarium of the South African National Fungus Collection

<sup>b</sup>Limpopo  
<sup>c</sup>Republic of South Africa  
<sup>d</sup>Western Cape  
<sup>e</sup>KwaZulu-Natal  
<sup>f</sup>Mpumalanga  
<sup>g</sup>Eastern Cape  
<sup>h</sup>Raoul Island  
<sup>i</sup>New Zealand  
<sup>j</sup>North Island

Alleles for isolates that could not be amplified were treated as a result of limited DNA available per extraction rather than PCR bias or null alleles, as discussed by McTaggart et al. (2018). The allele profiles for single-pustule isolates are shown in Supplementary Figure 1 and the scored alleles are listed in Table 1.



**Figure 1:** Telia (T) and uredinia (U) of *Austropuccinia psidii* on *Eugenia natalitia* in Dlinza Forest Reserve, KwaZulu Natal, South Africa (PREM 62056). Telia (meiotic stage) and uredinia (mitotic stage) were distinguished by their colour, where uredinia are yellow and telia are brown.

### *Genotypic diversity, gene diversity and tests for sexual reproduction*

We tested whether basidiospores played a role in the life cycle through analyses of genotypic diversity. If the population had high genotypic diversity there was evidence that recombinant spores (basidiospores) caused infection, rather than clonal urediniospores. Genotypic diversity of the populations was assessed using the genotypic evenness and Simpson Index, calculated from 1000 permutations of the *poppr* function in *poppr* (Kamvar et al. 2014; R Core Team 2014). All commands and data are provided in Supplementary File 2.

We used the level of gene diversity (heterozygosity) to determine whether sexual reproduction occurred in the invading population or if there were multiple sequential invasions, as a higher level of gene diversity may indicate multiple introductions. We used *GenoDive* (Meirmans and Van Tienderen 2004) to determine overall diversity for populations by the frequency of alleles across all five loci.

We used the Index of Association ( $I_A$ ) to test whether there was evidence of sexual reproduction within populations from South Africa and New Zealand, calculated with the *ia* function in *poppr*. This tested a null hypothesis of random mating and showed whether the observed multi-locus genotypes were best explained by clonal propagation (if there was linkage between markers) or recombination (if loci were not linked). The  $I_A$  was tested on clone-corrected and non-clone corrected data. We tested whether there was a random distribution of alleles among individuals using the inbreeding coefficient calculated in *GenoDive*.

We visualized recombination events using a Neighbour-Net in *SplitsTree4* v4.14.8 (Huson and Bryant 2005). Allelic data were converted to a distance matrix with *PopGenReport* (Adamack and Gruber 2014) using the genetic distance measure of Kosman and Leonard (2005), appropriate for organisms with mixed mating methods (Kosman and Leonard 2007). Reticulations within the Neighbour-Net were treated as putative recombination events in

populations (Huson and Bryant 2005).

## Results

### *Presence of sexual stage*

Telia were present on 16 out of the 35 (~46%) specimens of *A. psidii* from South Africa used in this study (Table 1), which supported the observation by Roux et al. (2016). Telia were observed on all five specimens from New Zealand. Telia were brown and compact, whereas uredinia were bright yellow and powdery (Figure 1).

### *Genotypic diversity, gene diversity and tests for sexual reproduction*

Genotypic diversity was high in both populations. The genotypic evenness of populations from New Zealand and South Africa showed a single (clonal) genotype was not dominant (Table 2). The Simpson Indices showed a high probability that two randomly selected samples from each population had a different genotype (Table 2).

**Table 2.** Genotypic diversity in populations of *Austropuccinia psidii* from New Zealand and South Africa. Genotypic evenness and the Simpson index show high genotypic diversity in the two populations of *A. psidii*.

Population	Number of pustules	Number of MLGs <sup>a</sup>	Expected number of MLGs	Standard error	Genotypic evenness	Simpson index
New Zealand	9	8	9	0	1	0.89
South Africa	35	31	31	0	0.79	0.96

<sup>a</sup>Multilocus genotype

Gene diversity showed both populations had one or two alleles for each locus. The exception was marker PpSSR018, which had three alleles, one of which was unique in two pustules of the South African population (Table 3). There were different alleles in the populations from New Zealand and South Africa, which shows they are not connected.

**Table 3.** Gene diversity in populations of *Austropuccinia psidii* from New Zealand and South Africa.


Population	Pustules	Frequency of alleles at each microsatellite locus					Gene diversity	Inbreeding coefficient (P value)
		PpSSR 012	PpSSR 014	PpSSR 018	PpSSR 102	PpSSR 161		
New Zealand	9	A 0.39	A 0.5	A 0.56	A 1	A 0.5	0.45	-0.069 (0.97)
		B 0.61	B 0.5	B 0.44		B 0.5		
South Africa	35	<b>C 0.45</b>	<b>C 0.47</b>	<b>C 0.42</b>	<b>A 0.45</b>	<b>C 0.47</b>	<b>0.58</b>	<b>-0.138 (0.55)</b>
		<b>D 0.55</b>	<b>D 0.53</b>	<b>D 0.53</b>	<b>B 0.55</b>	<b>D 0.53</b>		
				<b>B 0.05</b>				

The clone-corrected  $I_A$  values for the New Zealand and South African populations were not significant for linkage disequilibrium ( $P < 0.05$ ), which suggests random mating and provided evidence for sexual reproduction in the two populations (Table 4). The inbreeding coefficient of populations from New Zealand and South Africa was close to zero, which showed alleles were randomly distributed among the population, indicative of sexual reproduction (Table 4). The Neighbour-Net (Fig. 2) was reticulated within the populations, a sign of potential recombination events in the two examined populations.

**Table 4.** Test for linkage disequilibrium in populations of *Austropuccinia psidii* from New Zealand and South Africa. The clone corrected tests supported that loci were in linkage equilibrium ( $P I_A, P \bar{r}_d \geq 0.01$ ).

Population	Number of pustules	Number of MLGs	Non-clone corrected				Clone corrected			
			$I_A^a$	$P I_A$	$\bar{r}_d$	$P \bar{r}_d$	$I_A$	$P I_A$	$\bar{r}_d$	$P \bar{r}_d$
New Zealand	9	8	-0.50	0.998	-0.170	0.991	-0.50	0.998	-0.170	0.993
South Africa	35	31	0.22	0.001	0.055	0.001	0.066	0.053	0.017	0.053

<sup>a</sup> $I_A$  = Index of Association



**Figure 2:** Neighbour-Net made in SplitsTree based on Kosman genetic distance between multilocus genotypes of *A. psidii* from New Zealand and South Africa. Reticulations indicate possible recombinant relationships between the multilocus genotypes.

## Discussion

Our results support the hypothesis that invasive populations of *A. psidii* in New Zealand and South Africa reproduce sexually. This is based on three lines of evidence: (i) telia, the sexual stage, were present in the field, (ii) there was a high level of genotypic diversity within these invasive populations, and (iii) there were many recombinant multilocus genotypes with unlinked loci; all of which are putative signs of sexual reproduction (Milgroom 1996; De Meeûs et al. 2006). Teliospores, the precursors of sexual recombination in rust fungi, were identified in 46% (n=35) of the South African samples and in 100% (n=5) of the New Zealand samples. There was high genotypic diversity, despite low allelic diversity, in invasive populations from New Zealand and South Africa, as measured by the genotypic evenness and Simpson index. There was also no linkage between individual microsatellite markers, there was a random distribution of alleles, and the Neighbour-Net was reticulate, all of which are expected in recombinant populations.

There are three competing hypotheses to explain high genotypic diversity in an invasive population with low allelic diversity, namely sexual or parasexual reproduction, or

parallel evolution. Graça et al. (2013) treated *A. psidii* as clonal and hypothesized genotypic diversity was a result of parallel evolution. Wu et al. (2019) showed a species of rust reproduced by mitotic recombination (parasexual reproduction) when unable to complete its sexual life cycle. We cannot reject parasexual reproduction and parallel evolution as hypotheses based on our current data. However, McTaggart et al. (2018) resolved the sexual life cycle of *A. psidii* under controlled conditions; basidiospores infected Myrtaceae and dikaryotic, recombinant uredinia and telia were produced. With knowledge that *A. psidii* reproduces sexually, the most logical explanation from our findings is that basidiospores of *A. psidii* infect species of Myrtaceae to complete its sexual life cycle under field conditions.

An alternative hypothesis to explain high genotypic diversity is that observed multilocus genotypes were part of genetically different and genotypically diverse invasive populations. This alternative hypothesis is unlikely because there was only one locus that had a third allele, and agrees with findings that incursions in New Zealand and South Africa were caused by single strains (du Plessis et al. 2019; Roux et al. 2016). The third allele in the locus from South Africa was attributed to mutation based on its low frequency and that stepwise mutations are a characteristic of microsatellite loci (Valdes et al. 1993).

Our results, analyses and conclusions are in contrast to studies that found low genotypic diversity and clonal spread of *A. psidii* in Australia (Machado et al. 2015; Sandhu et al. 2016), Brazil (Graça et al. 2013), Colombia (Granados et al. 2017), Hawaii (Stewart et al. 2017), New Caledonia (Soewarto et al. 2017) and South Africa (Roux et al. 2016). Those studies sampled plants grown in monoculture, such as *Eucalyptus* and guava, sampled at different geographic scales, or were from recently invasive populations in which a founder effect may have led to low sampled genotypic diversity (Drenth et al. 2019; Gladioux et al. 2015; Taylor et al. 2015). Future studies need to investigate whether these populations are truly clonal or whether there is evidence of recombination, as shown in New Zealand and South Africa.

Our results demonstrate that *A. psidii* has a mixed mating system. Urediniospores are formed in abundance, spread quickly and maintain desirable genotypes by clonal reproduction. Teliospores and basidiospores play a role in dispersal of the pathogen and infection of Myrtaceae in addition to the long-term benefits of sexual recombination. Gladioux et al. (2015) proposed that mixed mating systems, with asexual and sexual reproduction, benefit fungal invasions. Clonal reproduction allows propagation without losing advantageous gene combinations (McDonald and Linde 2002), and recombination through sexual reproduction allows populations to increase diversity after a bottleneck (Barrett 2015).

Different genotypes of *A. psidii* could potentially recombine if their basidiospores infected the same host, based on the life cycle hypothesized by McTaggart et al. (2018). Recombination within or between genotypes enables pathogens to increase their speed of adaptation (reviewed by Nieuwenhuis and James 2016) and overcome resistance of host plants in fewer generations (McDonald and Linde 2002). Another species of rust, *Hemileia vastatrix*, the cause of coffee rust, undergoes cryptic sexual reproduction, and recombinant genotypes with new combinations of alleles overcame resistance in coffee plants (Carvalho et al. 2011). New combinations of pathogenicity and virulence may occur through sexual recombination, which typically reduces the lifespan of selective breeding for resistance.

Our findings show that invasive populations of *A. psidii* in New Zealand and South Africa reproduce sexually, which may increase its ability to more rapidly adapt to changes in the environment. *Austropuccinia psidii* is capable of quickly building inocula through clonal reproduction in the form of urediniospores, and of overcoming population bottlenecks and host resistance through recombination and potential outcrossing between genotypes.

## **Acknowledgements**

We are grateful to Kate and Graham Grieve (Custodians of Rare and Endangered Wildflowers, CREW), as well as Jackie Cossey, Izette Greyling, Giovanni Sale, Christo van

Zyl and Henk du Plessis for providing or assisting in collection of samples for this study. This work is based on research partially supported by the Tree Protection Co-operative Programme (TPCP) and the National Research Foundation of South Africa (Grant specific unique reference numbers UID 78566 and UID 83924) and the DST-NRF Centre of Excellence in Tree Health Biotechnology (CTHB). The grant holders acknowledge that opinions, findings and conclusions or recommendations expressed in any publication generated by NRF supported research are that of the authors and that the NRF accepts no liability whatsoever in this regard. ARM acknowledges the University of Queensland Development Fellowships (UQFEL1718905) and support from the Department of the Environment and Energy under the Australian Biological Resources Study (grant number RG18-43). This work was partly funded by the New Zealand Ministry of Business, Innovation and Employment (C09X1806), and the Australian Plant Biosecurity Science Foundation (PBSF018). We thank Luke Barrett and eight anonymous reviewers who have helped improve previous submissions of this manuscript.

### **Compliance with Ethical Standards**

The authors declare that there are no potential conflicts of interest and that the study did not involve human participants or animals.

### **Literature cited**

- Adamack, A. T., & Gruber, B. (2014). PopGenReport: simplifying basic population genetic analyses in R. *Methods in Ecology and Evolution*, 5(4), 384–387, doi:10.1111/2041-210X.12158.
- Barrett, S. C. (2015). Foundations of invasion genetics: the Baker and Stebbins legacy. *Molecular Ecology*, 24(9), 1927–1941, doi:10.1111/mec.13014.
- Carnegie, A. J., & Pegg, G. S. (2018). Lessons from the incursion of myrtle rust in Australia. *Annual Review of Phytopathology*, 56(1), 457–478, doi:10.1146/annurev-phyto-080516-035256.
- Carvalho, C. R., Fernandes, R. C., Carvalho, G. M. A., Barreto, R. W., & Evans, H. C. (2011). Cryptosexuality and the genetic diversity paradox in coffee rust, *Hemileia vastatrix*. *PLoS ONE*, 6(11), e26387, doi:10.1371/journal.pone.0026387.
- Coutinho, T. A., Wingfield, M. J., Alfenas, A. C., & Crous, P. W. (1998). Eucalyptus rust: a disease with the potential for serious international implications. *Plant Disease*, 82(7), 819–825, doi:10.1094/pdis.1998.82.7.819.
- De Meeûs, T., Lehmann, L., & Balloux, F. (2006). Molecular epidemiology of clonal diploids: A quick overview and a short DIY (do it yourself) notice. *Infection, Genetics and Evolution*, 6(2), 163–170, doi:<https://doi.org/10.1016/j.meegid.2005.02.004>.

- Drenth, A., McTaggart, A. R., & Wingfield, B. D. (2019). Fungal clones win the battle, recombination wins the war. *IMA Fungus*, *10*(18), <https://doi.org/10.1186/s43008-019-0020-8>.
- du Plessis, E., Granados, G. M., Barnes, I., Ho, W. H., Alexander, B. J. R., Roux, J., et al. (2019). The pandemic strain of *Austropuccinia psidii* causes myrtle rust in New Zealand and Singapore. *Australasian Plant Pathology*, *48*, 253–256, doi:10.1007/s13313-019-0624-x.
- Figueiredo, M. B. (2001). Life cycle and ecology of *Puccinia psidii*. *O Biologico*, *63*, 69–71.
- Figueiredo, M. B., Coutinho, L. N., & Hennen, J. F. (1984). Studies on the determination the life cycle of *Puccinia psidii* Winter. *Summa Phytopathologica*, *10*, 53–54.
- Gladieux, P., Feurtey, A., Hood, M. E., Snirc, A., Clavel, J., Dutech, C., et al. (2015). The population biology of fungal invasions. *Molecular Ecology*, *24*(9), 1969–1986, doi:10.1111/mec.13028.
- Graça, R. N., Ross-Davis, A. L., Klopfenstein, N. B., Kim, M.-S., Peever, T. L., Cannon, P. G., et al. (2013). Rust disease of eucalypts, caused by *Puccinia psidii*, did not originate via host jump from guava in Brazil. *Molecular Ecology*, *22*(24), 6033–6047, doi:10.1111/mec.12545.
- Granados, G. M., McTaggart, A. R., Barnes, I., Rodas, C. A., Roux, J., & Wingfield, M. J. (2017). The pandemic biotype of *Austropuccinia psidii* discovered in South America. *Australasian Plant Pathology*, *46*(3), 267–275, doi:10.1007/s13313-017-0488-x.
- Grunwald, N. J., Everhart, S. E., Knaus, B. J., & Kamvar, Z. N. (2017). Best practices for population genetic analyses. *Phytopathology*, *107*(9), 1000–1010, doi:10.1094/phyto-12-16-0425-rvw.
- Ho, W. H., Baskarathevan, J., Griffin, R. L., Quinn, B. D., Alexander, B. J. R., Havell, D., et al. (2019). First report of myrtle rust caused by *Austropuccinia psidii* on *Metrosideros kermadecensis* on Raoul Island and on *M. excelsa* in Kerikeri, New Zealand. *Plant Disease*, PDIS-12-18-2243-PDN, doi:10.1094/PDIS-12-18-2243-PDN.
- Huson, D. H., & Bryant, D. (2005). Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution*, *23*(2), 254–267, doi:10.1093/molbev/msj030.
- Joffily, J. (1944). Ferrugem do eucalipto. *Bragantia*, *4*, 475–487.
- Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*, e281, doi:10.7717/peerj.281.
- Kosman, E., & Leonard, K. J. (2005). Similarity coefficients for molecular markers in studies of genetic relationships between individuals for haploid, diploid, and polyploid species. *Molecular Ecology*, *14*(2), 415–424, doi:10.1111/j.1365-294X.2005.02416.x.
- Kosman, E., & Leonard, K. J. (2007). Conceptual analysis of methods applied to assessment of diversity within and distance between populations with asexual or mixed mode of reproduction. *New Phytologist*, *174*(3), 683–696, doi:10.1111/j.1469-8137.2007.02031.x.
- Machado, P. d. S., Alfenas, A. C., Alfenas, R. F., Mohammed, C. L., & Glen, M. (2015). Microsatellite analysis indicates that *Puccinia psidii* in Australia is mutating but not recombining. *Australasian Plant Pathology*, *44*(4), 455–462, doi:10.1007/s13313-015-0364-5.

- McDonald, B. A., & Linde, C. (2002). The population genetics of plant pathogens and breeding strategies for durable resistance. [journal article]. *Euphytica*, 124(2), 163–180, doi:10.1023/a:1015678432355.
- McTaggart, A. R., Shuey, L. S., Granados, G. M., du Plessis, E., Fraser, S., Barnes, I., et al. (2018). Evidence that *Austropuccinia psidii* may complete its sexual life cycle on Myrtaceae. *Plant Pathology*, 67, 729–734, doi:10.1111/ppa.12763.
- Meirmans, P. G., & Van Tienderen, P. H. (2004). GenoType and GenoDive: two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes*, 4(4), 792–794, doi:10.1111/j.1471-8286.2004.00770.x.
- Milgroom, M. G. (1996). Recombination and the multilocus structure of fungal populations. *Annual Review of Phytopathology*, 34, 457–477, doi:10.1146/annurev.phyto.34.1.457.
- Morin, L., Talbot, M. J., & Glen, M. (2014). Quest to elucidate the life cycle of *Puccinia psidii* sensu lato. *Fungal Biology*, 118, 253–263.
- Nieuwenhuis, B. P. S., & James, T. Y. (2016). The frequency of sex in fungi. [10.1098/rstb.2015.0540]. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150540.
- R Core Team (2014). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rangel, E. (1916). Contribuição para o estudo dos Puccinias das Myrtáceas. *Archivos do Museu Nacional do Rio de Janeiro*, 18, 147–156.
- Roux, J., Granados, G. M., Shuey, L., Barnes, I., Wingfield, M. J., & McTaggart, A. R. (2016). A unique genotype of the rust pathogen, *Puccinia psidii*, on Myrtaceae in South Africa. *Australasian Plant Pathology*, 45, 645–652, doi:10.1007/s13313-016-0447-y.
- Roux, J., Greyling, I., Coutinho, T. A., Verleur, M., & Wingfield, M. J. (2013). The myrtle rust pathogen, *Puccinia psidii*, discovered in Africa. *IMA Fungus*, 4(1), 155–159.
- Ruiz, R. A. R., Alfenas, A. C., Ferreira, F. A., & Vale, F. X. R. (1989). Influência da temperatura, do tempo de molhamento foliar, fotoperíodo e intensidade de luz sobre a infecção de *Puccinia psidii* em eucalipto. *Fitopatologia Brasileira*, 14, 55–61.
- Sandhu, K. S., Karaoglu, H., Zhang, P., & Park, R. F. (2016). Simple sequence repeat markers support the presence of a single genotype of *Puccinia psidii* in Australia. *Plant Pathology*, 65, 1084–1094.
- Soewarto, J., Carriconde, F., Hugot, N., Bocs, S., Hamelin, C., & Maggia, L. (2017). Impact of *Austropuccinia psidii* in New Caledonia, a biodiversity hotspot. *Forest Pathology*, 48(2), e12402, doi:10.1111/efp.12402.
- Stewart, J. E., Ross-Davis, A. L., Graça, R. N., Alfenas, A. C., Peever, T. L., Hanna, J. W., et al. (2017). Genetic diversity of the myrtle rust pathogen (*Austropuccinia psidii*) in the Americas and Hawaii: Global implications for invasive threat assessments. *Forest Pathology*, 48, e12378, doi:10.1111/efp.12378.
- Taylor, J. W., Hann-Soden, C., Branco, S., Sylvain, I., & Ellison, C. E. (2015). Clonal reproduction in fungi. *Proceedings of the National Academy of Sciences*, 112(29), 8901–8908, doi:10.1073/pnas.1503159112.
- Valdes, A. M., Slatkin, M., & Freimer, N. B. (1993). Allele frequencies at microsatellite loci: the stepwise mutation model revisited. *Genetics*, 133(3), 737.
- Winter, G. (1884). *Dr. L. Rabenhorst's Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz. Erster Band: Pilze* (Vol. 1, pt. 1). Leipzig: Verlag von Eduard Kummer.

- Wu, J. Q., Dong, C., Song, L., Cuomo, C. A., & Park, R. F. (2019). Dissecting the first phased dikaryotic genomes of the wheat rust pathogen *Puccinia triticina* reveals the mechanisms of somatic exchange in nature. *bioRxiv*, doi: <https://doi.org/10.1101/705475>
- Zhong, S., Yang, B., & Alfenas, A. C. (2008). Permanent genetic resources: Development of microsatellite markers for the guava rust fungus, *Puccinia psidii*. *Molecular Ecology Resources*, 8(2), 348–350, doi:10.1111/j.1471-8286.2007.01952.x.